

(610)

10/752342

403

11/01/04

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number
WO 02/068406 A2(51) International Patent Classification⁷: C07D 401/00

(21) International Application Number: PCT/US02/03064

(22) International Filing Date: 11 January 2002 (11.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/261,882 12 January 2001 (12.01.2001) US

60/323,808 19 September 2001 (19.09.2001) US

10/046,622 10 January 2002 (10.01.2002) US

(71) Applicant: AMGEN INC. [US/US]; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).

(72) Inventors: CHEN, Guoqing; 515 Oakbury Court, Thousand Oaks, CA 91360 (US). ADAMS, Jeffrey; 2915 Venezia Lane, Thousand Oaks, CA 91362 (US). BEMIS, Jean; 256 Appleton Street, Arlington, MA 02476 (US). CROGHAN, Michael; 1108 Dover Lane, Ventura, CA 93001 (US). DIPIETRO, Lucian; 37 Centennial Avenue, Gloucester, MA 01930 (US). DOMINGUEZ, Celia; 963 Cedarcliff Court, Thousand Oaks, CA 91362 (US). ELBAUM, Daniel; 25 Sherrin Road, Newton, MA 02462 (US). GERMAIN, Julie; 28 Woods Avenue, Somerville, MA 02144 (US). HUANG, Qi; 12308 Willow Forest Drive, Moorpark, CA 93021 (US). KIM, Joseph, L.; 20 Green Way, Wayland, MA 01778 (US). OUYANG, Xiaohu; 132-20 Maple Avenue, Flushing, NY 11355 (US). PATEL, Vinod, F.; 3 Mossy Lane, Acton, MA 01720 (US). SMITH, Leon, M.; 33 Julie Court, Sommerset, NJ 08873 (US). TASKER, Andrew; 561 Granite Hills Street,

Simi Valley, CA 93065 (US). XI, Ning; 565 Timberwood Avenue, Thousand Oaks, CA 91360 (US). XU, Shimin; 1890 West Hillcrest Drive, Number 406, Newbury Park, CA 91320 (US). YUAN, Chester, Chenguang; 1416 Geyser Court, Newbury Park, CA 91320 (US). KIM, Tae-Seong; 1580 Dover Avenue, Thousand Oaks, CA 91360 (US).

(74) Agents: ODRE, Steven, M. et al.; Amgen Inc., One Amgen Center Drive, M/S 27-4-A, Thousand Oaks, CA 91320-1799 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/068406 A2

(54) Title: SUBSTITUTED AMINE DERIVATIVES AND METHODS OF USE

(57) Abstract: Selected amines are effective for prophylaxis and treatment of diseases, such as angiogenesis mediated diseases. The invention encompasses novel compounds, analogs, prodrugs and pharmaceutically acceptable salts thereof, pharmaceutical compositions and methods for prophylaxis and treatment of diseases and other maladies or conditions involving, cancer and the like. The subject invention also relates to processes for making such compounds as well as to intermediates useful in such processes.

- 1 -

SUBSTITUTED AMINE DERIVATIVES AND METHODS OF USE

FIELD OF THE INVENTION

5

This invention is in the field of pharmaceutical agents and specifically relates to compounds, compositions, uses and methods for treating cancer and angiogenesis-related disorders.

10

BACKGROUND OF THE INVENTION

Protein kinases represent a large family of proteins which play a central role in the regulation of a wide variety of cellular processes, maintaining control over cellular function. A partial list of such kinases includes ab1, Akt, bcr-ab1, Blk, Brk, Btk, c-kit, c-met, c-src, CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8, CDK9, CDK10, cRaf1, CSF1R, CSK, EGFR, ErbB2, ErbB3, ErbB4, Erk, Fak, fes, FGFR1, FGFR2, FGFR3, FGFR4, FGFR5, Fgr, flt-1, Fps, Frk, Fyn, Hck, IGF-1R, INS-R, Jak, KDR, Lck, Lyn, MEK, p38, PDGFR, PIK, PKC, PYK2, ros, tie, tie2, TRK, Yes, and Zap70. Inhibition of such kinases has become an important therapeutic target.

25 Certain diseases are known to be associated with deregulated angiogenesis, for example ocular neovascularisation, such as retinopathies (including diabetic retinopathy), age-related macular degeneration, psoriasis, hemangioblastoma, hemangioma, arteriosclerosis, inflammatory disease, such as a rheumatoid or rheumatic
30 inflammatory disease, especially arthritis (including rheumatoid arthritis), or other chronic inflammatory disorders, such as chronic asthma, arterial or post-transplantational atherosclerosis, endometriosis, and
35 neoplastic diseases, for example so-called solid tumors and liquid tumors (such as leukemias).

- 2 -

At the center of the network regulating the growth and differentiation of the vascular system and its components, both during embryonic development and normal growth, and in a wide number of pathological anomalies and diseases, lies the angiogenic factor known as Vascular Endothelial Growth Factor" (VEGF; originally termed 'Vascular Permeability Factor', VPF), along with its cellular receptors (see G. Breier et al., Trends in Cell Biology, 6, 454-6 (1996)).

VEGF is a dimeric, disulfide-linked 46-kDa glycoprotein related to "Platelet-Derived Growth Factor" (PDGF); it is produced by normal cell lines and tumor cell lines; is an endothelial cell-specific mitogen; shows angiogenic activity in *in vivo* test systems (e.g. rabbit cornea); is chemotactic for endothelial cells and monocytes; and induces plasminogen activators in endothelial cells, which are involved in the proteolytic degradation of extracellular matrix during the formation of capillaries. A number of isoforms of VEGF are known, which show comparable biological activity, but differ in the type of cells that secrete them and in their heparin-binding capacity. In addition, there are other members of the VEGF family, such as "Placenta Growth Factor" (PlGF) and VEGF-C.

VEGF receptors (VEGFR) are transmembranous receptor tyrosine kinases. They are characterized by an extracellular domain with seven immunoglobulin-like domains and an intracellular tyrosine kinase domain. Various types of VEGF receptor are known, e.g. VEGFR-1 (also known as flt-1), VEGFR-2 (also known as KDR), and VEGFR-3.

A large number of human tumors, especially gliomas and carcinomas, express high levels of VEGF and its receptors. This has led to the hypothesis that the VEGF released by tumor cells stimulates the growth of blood capillaries and the proliferation of tumor endothelium in a paracrine manner and through the improved blood supply, accelerate tumor

- 3 -

growth. Increased VEGF expression could explain the occurrence of cerebral edema in patients with glioma. Direct evidence of the role of VEGF as a tumor angiogenesis factor *in vivo* is shown in studies in which VEGF expression or VEGF activity was inhibited. This was achieved with anti-VEGF antibodies, with dominant-negative VEGFR-2 mutants which inhibited signal transduction, and with antisense-VEGF RNA techniques. All approaches led to a reduction in the growth of glioma cell lines or other tumor cell lines *in vivo* as a result of inhibited tumor angiogenesis.

Angiogenesis is regarded as an absolute prerequisite for tumors which grow beyond a diameter of about 1-2 mm; up to this limit, oxygen and nutrients may be supplied to the tumor cells by diffusion. Every tumor, regardless of its origin and its cause, is thus dependent on angiogenesis for its growth after it has reached a certain size.

Three principal mechanisms play an important part in the activity of angiogenesis inhibitors against tumors: 1) Inhibition of the growth of vessels, especially capillaries, into avascular resting tumors, with the result that there is no net tumor growth owing to the balance that is achieved between cell death and proliferation; 2) Prevention of the migration of tumor cells owing to the absence of blood flow to and from tumors; and 3) Inhibition of endothelial cell proliferation, thus avoiding the paracrine growth-stimulating effect exerted on the surrounding tissue by the endothelial cells which normally line the vessels. See R. Connell and J. Beebe, *Exp. Opin. Ther. Patents*, 11, 77-114 (2001).

VEGF's are unique in that they are the only angiogenic growth factors known to contribute to vascular hyperpermeability and the formation of edema. Indeed, vascular hyperpermeability and edema that is associated with

- 4 -

the expression or administration of many other growth factors appears to be mediated via VEGF production.

Inflammatory cytokines stimulate VEGF production.

Hypoxia results in a marked upregulation of VEGF in numerous
5 tissues, hence situations involving infarct, occlusion, ischemia, anemia, or circulatory impairment typically invoke VEGF/VPF-mediated responses. Vascular hyperpermeability, associated edema, altered transendothelial exchange and macromolecular extravasation, which is often accompanied by
10 diapedesis, can result in excessive matrix deposition, aberrant stromal proliferation, fibrosis, etc. Hence, VEGF-mediated hyperpermeability can significantly contribute to disorders with these etiologic features. As such, regulators of angiogenesis have become an important therapeutic target.

15 Schipper US patent 3,226,394, issued Dec. 28, 1965, describes anthranilamides as CNS depressants. Japanese patent JP2000256358 describes pyrazole derivatives that block the calcium release-activated calcium channel. EP application 9475000, published 6 October 1999, describes
20 compounds as PGE₂ antagonists. PCT publication WO96/41795, published 27 December 1996, describes benzamides as vasopressin antagonists. WO01/29009 describes aminopyridines as KDR inhibitors. WO01/30745 describes anthranilic acids as cGMP phosphodiesterase inhibitors.
25 WO00/02851, published 20 Jan 2000 describes arylsulfonylaminoaryl amides as guanylate cyclase activators. WO98/45268 describes nicotinamide derivatives as PDE4 inhibitors. WO98/24771 describes benzamides as vasopressin antagonists.

30 US Patent No. 5,532,358, issued July 2, 1996, describes the preparation of 2-(cyclopropylamino)-N-(2-methoxy-4-methyl-3-pyridinyl)-3-pyridinecarboxamide as an intermediate for HIV inhibitors. Triazine-substituted amines are described for their aggregating ability (J. Amer.


- 5 -

Chem. Soc., 115, 905-16 (1993). Substituted imidazolines were tested for their antidepressant activity in Ind. J. Het. Chem., 2, 129-32 (1992). N-(4-Pyridyl)anthranilic amides were described in Chem Abstr. 97:109837 (1981). PCT publication WO99/32477, published 1 July 1999, describes anthranilamides as anti-coagulants. US patent 6,140,351 describes anthranilamides as anti-coagulants. PCT publication WO99/62885, published 9 December 1999, describes 1-(4-aminophenyl)pyrazoles as antiinflammatories. PCT publication WO00/39111, published 6 July 2000, describes amides as factor Xa inhibitors. PCT publication WO00/39117, published 6 July 2000, describes heteroaromatic amides as factor Xa inhibitors. PCT publication WO00/27819, published 18 May 2000, describes anthranilic acid amides as VEGF inhibitors. PCT publication WO00/27820 published 18 May 2000, describes N-aryl anthranilic acid amides as VEGF inhibitors. 7-Chloroquinolinylamines are described in FR2168227 as antiinflammatories. WO01/55114, published 2 Aug. 2001, describes nicotinamides for the treatment of cancer. WO01/55115, published 2 Aug. 2001, describes nicotinamides for the treatment of apoptosis. WO01/85715, published 15 November 2001, describes substituted pyridines and pyrimidines as anti-angiogenesis agents. PCT publication WO01/85691 published 15 November 2001, describes anthranilic amides as VEGF inhibitors. PCT publication WO01/85671 published 15 November 2001, describes anthranilic amides as VEGF inhibitors. PCT publication WO01/81311 published 1 November 2001, describes anthranilic amides as VEGF inhibitors. However, compounds of the current invention have not been described as inhibitors of angiogenesis such as for the treatment of cancer.

- 6 -

DESCRIPTION OF THE INVENTION

A class of compounds useful in treating cancer and
 5 angiogenesis is defined by Formula I



wherein each of A¹ and A² is independently C, CH or N;

10 wherein ring A is selected from


a) 5- or 6-membered partially saturated heterocyclyl,
 preferably dihydropyran, dihydrothienyl,
 dihydrofuryl, oxo-dihydrofuryl, pyrrolinyl,
 dihydrothiazolyl, dihydro-oxazolyl, dihydro-
 15 isothiazolyl, dihydro-isoxazolyl, imidazolinyl
 and pyrazolinyl,

b) 5- or 6-membered heteroaryl,
 preferably

1) 5-membered heteroaryl selected from
 20 thienyl, furanyl, pyrrolyl, thiazolyl,
 oxazolyl, imidazolyl, pyrazolyl, isoxazolyl,
 triazolyl and isothiazolyl,
 even more preferably 5-membered heteroaryl
 selected from

25


- 7 -



5


specifically

A)





10

B)




- 8 -




5

II) preferably 6-membered heteroaryl selected
from pyridyl, pyrazinyl, pyrimidinyl,
pyridazinyl, and triazinyl,
even more preferably 6-membered heteroaryl
selected from

10



more specifically





- c) 9-, 10- or 11-membered fused partially saturated
heterocyclyl,
preferably tetrahydroquinolinyl,
d) 9- or 10-membered fused heteroaryl,
preferably

15

- 9 -

- i) fused 9-membered fused heteroaryl selected from benzothienyl, benzothiazolyl, indolyl, benzimidazolyl, benzoxazolyl, benzofuryl, indazolyl and isoindolyl, and
- 5 ii) fused 10-membered heteroaryl selected from quinolyl, isoquinolyl, naphthpyridinyl, quinoxalinyl and quinazolinyl,
- e) aryl, and
- f) 4-, 5- or 6-membered cycloalkenyl,
- 10 preferably 5-membered cycloalkenyl, more preferably cyclopentadienyl or cyclopentenyl;



wherein Z is oxygen or sulfur;
wherein R is selected from

- a) substituted or unsubstituted 4-6 membered
- 20 heterocyclyl,
preferably substituted or unsubstituted 5-6 membered heteroaryl comprising one or more nitrogen atoms, more preferably pyrazolyl, triazolyl, pyridyl, pyrimidinyl, and pyridazinyl,

- 10 -

even more preferably 4-pyridyl, 3-pyridyl, 2-pyridyl, triazolyl, 4-pyrimidinyl and 4-pyridazinyl,
most preferably 4-pyridyl,

- 5 b) substituted aryl,
 preferably substituted phenyl,
 more preferably optionally substituted
 (heterocyclyl-substituted phenyl), and
- 10 c) substituted or unsubstituted 9-14-membered bicyclic or
 tricyclic heterocyclyl,
 preferably substituted or unsubstituted 9-10 membered
 bicyclic or 13-14 membered tricyclic heterocyclyl,
 more preferably substituted or unsubstituted 9-10
 membered fused heterocyclyl,
15 even more preferably indazolyl, indolyl,
 isoindolyl, quinoliny, isoquinoliny,
 benzotriazolyl, 2,3-dihydrobenzofuryl, 1,2-
 dihydroquinolyl, naphthyridinyl and
 quinazoliny,
20 even more preferably 5-indazolyl, 6-indazolyl,
 4-quinolyl, 5-quinolyl, 6-quinolyl,
 indolyl, isoindolyl, benzotriazolyl, 2,3-
 dihydrobenzofuryl, 1,2-dihydroquinolyl,
 quinoxaliny, 4-isoquinolyl, 5-isoquinolyl,
25 naphthyridinyl and 6-isoquinolyl,
 especially preferred is 6-indazolyl;

wherein substituted R is substituted with one or more
substituents independently selected from halo, -OR³,
-SR³, -SO₂R³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -SO₂NR³R³,
30 -NR³C(O)OR³, -NR³C(O)R³, -NR³C(O)NR³R³, cycloalkyl,
 optionally substituted 4-6 membered heterocyclyl,
 optionally substituted phenyl, nitro, oxo,
 alkylaminoalkoxyalkoxy, cyano, alkylaminoalkoxy,
 lower alkyl substituted with R², lower alkenyl

- 11 -

substituted with R^2 , and lower alkynyl substituted with R^2 ;
preferably halo, $-OR^3$, $-SR^3$, $-SO_2R^3$, $-CO_2R^3$, $-CONR^3R^3$,
5 $-COR^3$, $-NR^3R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$,
 C_{3-6} -cycloalkyl, optionally substituted 4-6
membered heterocyclyl, optionally substituted
phenyl, nitro, C_{1-4} -alkylamino- C_{1-4} -alkoxy- C_{1-4} -
alkoxy, cyano, C_{1-4} -alkylamino- C_{1-4} -alkoxy, C_{1-2} -
alkyl substituted with R^2 , C_{2-3} -alkenyl
10 substituted with R^2 , and C_{2-3} -alkynyl substituted
with R^2 ,
more preferably halo, hydroxy, C_{1-4} -alkyl, C_{1-2} -
alkoxy, optionally substituted 4-6 membered
heterocyclyl- C_{1-2} -alkoxy, amino, C_{1-2} -
15 alkylamino, aminosulfonyl, $-NR^3C(O)OR^3$, $-$
 $NR^3C(O)R^3$, C_{3-6} -cycloalkyl, optionally
substituted 4-6 membered heterocyclyl,
optionally substituted phenyl, nitro, C_{1-2} -
alkylamino- C_{1-2} -alkoxy- C_{1-2} -alkoxy, cyano, C_{1-2} -
20 alkylamino- C_{1-2} -alkoxy, C_{1-2} -alkylamino- C_{1-2} -
alkyl, C_{1-2} -alkylamino- C_{2-3} -alkynyl, C_{1-2} -
hydroxyalkyl, C_{1-2} -aminoalkyl, C_{1-2} -haloalkyl,
optionally substituted 4-6 membered
heterocyclyl- C_{2-3} -alkenyl, and optionally
25 substituted 4-6 membered heterocyclyl- C_{2-3} -
alkynyl,
even more preferably chloro, fluoro, bromo,
hydroxy, methoxy, ethoxy, amino,
dimethylamino, diethylamino, 1-
30 methylpiperidinylmethoxy, aminosulfonyl,
cyclohexyl, dimethylaminopropynyl,
dimethylaminoethoxy, 3-(4-
morpholinyl)propyn-1-yl,
dimethylaminoethoxyethoxy, optionally

- 12 -

- substituted piperidinyl, morpholinyl,
optionally substituted piperazinyl,
optionally substituted phenyl, methyl,
ethyl, propyl, cyano, hydroxymethyl,
5 aminomethyl, nitro and trifluoromethyl;
wherein R¹ is selected from
- a) substituted or unsubstituted 6-10 membered aryl,
preferably phenyl, naphthyl, indanyl, indenyl and
tetrahydronaphthyl,
10 more preferably phenyl, tetrahydronaphthyl, and
naphthyl,
- b) substituted or unsubstituted 4-6 membered
heterocyclyl,
preferably 5-6 membered heteroaryl,
15 more preferably isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl,
pyridazinyl, imidazolyl, oxazolyl, furyl and
pyrrolyl,
- c) substituted or unsubstituted 9-14 membered bicyclic or
20 tricyclic heterocyclyl,
preferably 9-10 membered bicyclic or 13-14 membered
tricyclic heterocyclyl,
more preferably indazolyl, indolyl, isoindolyl,
2,3-dihydro-1H-indolyl, naphthyridinyl, 2,1,3-
25 benzothiadiazolyl, isoquinolyl, quinolyl, 1,2-
dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl,
benzo[d]isothiazolyl, benzothienyl,
30 tetrahydroquinolyl, benzofuryl,
dihydrobenimidazolyl, benzimidazolyl,
benzoxazolyl, benzthiazolyl, benzodioxanyl and
quinazolinyl,
d) cycloalkyl,

- 13 -

preferably C₃₋₆-cycloalkyl,

more preferably cyclohexyl, and

e) cycloalkenyl,


wherein substituted R¹ is substituted with one or more

5 substituents independently selected from halo, oxo, -OR³, -SR³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -NH(C₁-C₄ alkylenylR⁴), -SO₂R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, -NR³C(O)NR³R³, optionally substituted cycloalkyl, optionally substituted 4-6 membered

10 heterocyclyl, optionally substituted phenyl, halosulfonyl, cyano, alkylaminoalkoxy, alkylaminoalkoxyalkoxy, nitro, lower alkyl substituted with R², lower alkenyl substituted with R², and lower alkynyl substituted with R²,

15 preferably halo, -OR³, oxo, -SR³, -SO₂R³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -NH(C₁-C₄ alkylenylR³), - (C₁-C₄ alkylenyl)NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₁-C₆-alkylamino-C₁-C₆-alkoxy, C₁-C₆-alkylamino-C₁-C₆-alkoxy-C₁-C₆-alkoxy, halosulfonyl,

20 optionally substituted 4-6 membered heterocyclylcarbonylalkyl, C₁₋₄-


alkoxycarbonylamino-C₁₋₆-alkyl, , optionally substituted C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₆-alkylenyl, optionally substituted 4-6 membered heterocyclyl-C₁-C₆-alkylenyl, 4-6 membered heterocyclyl-C₂-C₆-alkenylenyl, C₁₋₄-alkyl, cyano, C₁₋₄-hydroxyalkyl, nitro and C₁₋₄-haloalkyl,

25 more preferably halo, C₁₋₄-alkyl, optionally substituted C₃₋₆-cycloalkyl, optionally substituted phenyl, optionally substituted

30

- 14 -

phenyl-C₁-C₄-alkylenyl, C₁₋₂-haloalkoxy,
 optionally substituted phenyloxy, optionally
 substituted 4-6 membered heterocyclyl-C₁-C₄-
 alkylenyl, optionally substituted 4-6 membered
 5 heterocyclyl-C₂-C₄-alkenylenyl, optionally
 substituted 4-6 membered heterocyclyl,
 optionally substituted 4-6 membered
 heterocyclyloxy, optionally substituted 4-6
 membered heterocyclylsulfonyl, optionally
 10 substituted 4-6 membered heterocyclylamino,
 optionally substituted 4-6 membered
 heterocyclylcarbonyl, optionally substituted
 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl,
 C₁₋₂-haloalkyl, C₁₋₄-aminoalkyl, nitro, amino,
 15 hydroxy, oxo, cyano, aminosulfonyl, C₁₋₂-
 alkylsulfonyl, halosulfonyl, C₁₋₄-
 alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
 alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-
 alkoxy-C₁₋₃-alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-
 20 alkoxycarbonylamino-C₁₋₄-alkyl, C₁₋₄-

hydroxyalkyl,  and C₁₋₄-alkoxy, and

even more preferably bromo, chloro, fluoro, iodo,
 nitro, amino, cyano, aminoethyl, Boc-
 aminoethyl, hydroxy, oxo, aminosulfonyl, 4-
 25 methylpiperazinylsulfonyl, cyclohexyl, phenyl,
 phenylmethyl, morpholinylmethyl, 1-
 methylpiperazin-4-ylmethyl, 1-methylpiperazin-
 4-ylpropyl, morpholinylpropyl, piperidin-1-
 ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-
 30 methyl-2-(1-methylpiperidin-4-yl)ethyl,
 morpholinylethyl, 1-(4-morpholinyl)-2,2-
 dimethylpropyl, piperidin-4-ylethyl, 1-Boc-
 piperidin-4-ylethyl, piperidin-1-ylethyl, 1-

- 15 -

Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl,
1-Boc-piperidin-4-ylmethyl, piperidin-4-
ylpropyl, 1-Boc-piperidin-4-ylpropyl,
piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl,
5 pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-
ylpropyl, pyrrolidin-1-ylmethyl, pyrrolidin-2-
ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl,
pyrrolidinylpropenyl, pyrrolidinylbutenyl,
fluorosulfonyl, methylsulfonyl,
10 methylcarbonyl, Boc; piperidin-1-
ylmethylcarbonyl, 4-methylpiperazin-1-
ylcarbonylethyl, methoxycarbonyl,
aminomethylcarbonyl,
dimethylaminomethylcarbonyl, 3-ethoxycarbonyl-
15 2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-
methyl-1-piperidyl, 1-Boc-4-piperidyl,
piperidin-4-yl, 1-methylpiperidin-4-yl, 1-
methyl-(1,2,3,6-tetrahydropyridyl),
imidazolyl, morpholinyl, 4-trifluoromethyl-1-
20 piperidinyl, hydroxybutyl, methyl, ethyl,
propyl, isopropyl, butyl, tert-butyl, sec-
butyl, trifluoromethyl, pentafluoroethyl,
nonafluorobutyl, dimethylaminopropyl, 1,1-
di(trifluoromethyl)-1-hydroxymethyl, 1,1-
25 di(trifluoromethyl)-1-
(piperidinylethoxy)methyl, 1,1-
di(trifluoromethyl)-1-
(methoxyethoxyethoxy)methyl, 1-hydroxyethyl,
2-hydroxyethyl, trifluoromethoxy, 1-
30 aminoethyl, 2-aminoethyl, 1-(N-
isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-
chlorophenoxy, phenyloxy, azetidin-3-
ylmethoxy, 1-Boc-azetidin-3-ylmethoxy, pyrrol-

- 16 -

2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-
1-ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-
isopropyl-pyrrol-2-ylmethoxy, 1-Boc-piperdin-
4-ylmethoxy, piperdin-4-ylmethoxy, 1-
5 methylpiperdin-4-yloxy, isopropoxy, methoxy
and ethoxy;

wherein R² is one or more substituents independently selected
from H, halo, -OR³, oxo, -SR³, -CO₂R³, -COR³, -CONR³R³, -
NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, cycloalkyl,
10 optionally substituted phenylalkyl, optionally
substituted 4-6 membered heterocyclyl, optionally
substituted heteroarylalkyl, optionally substituted
phenyl, lower alkyl, cyano, lower hydroxyalkyl, lower
carboxyalkyl, nitro, lower alkenyl, lower alkynyl, C₁₋₄-
15 alkoxy-C₁₋₄-alkoxy and C₁₋₄-alkoxy-C₁₋₄-alkoxy-C₁₋₄-alkoxy,
lower aminoalkyl, lower alkylaminoalkyl and lower
haloalkyl,
preferably H, halo, -OR³, -SR³, -CO₂R³, -CONR³R³, -COR³, -
NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₃₋₆-cycloalkyl,
20 optionally substituted 4-6 membered heterocyclyl,
optionally substituted phenyl, C₁₋₆-alkyl, cyano, C₁₋₄-
hydroxyalkyl, C₁₋₄-carboxyalkyl, nitro, C₁₋₃-alkoxy-C₁₋₂-
alkoxy, C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₂₋₃-alkenyl, C₂₋₃-
alkynyl and C₁₋₄-haloalkyl,
25 more preferably H, halo, hydroxy, C₁₋₂-alkoxy, C₁₋₂-
haloalkoxy, amino, C₁₋₂-alkylamino, optionally
substituted 4-6 membered heterocyclyl-C₁₋₂-
alkylamino, aminosulfonyl, C₃₋₆-cycloalkyl,
optionally substituted 4-6 membered heterocyclyl,
30 optionally substituted phenyl, C₁₋₄-alkyl, cyano, C₁₋₂-
hydroxyalkyl, C₁₋₃-carboxyalkyl, nitro, C₂₋₃-
alkenyl, C₂₋₃-alkynyl and C₁₋₂-haloalkyl, and
even more preferably H, chloro, fluoro, bromo,
hydroxy, methoxy, ethoxy, trifluoromethoxy,

- 17 -

amino, dimethylamino, aminosulfonyl,
carboxymethyl, cyclopropyl, optionally
substituted phenyl, methyl, ethyl, propyl, cyano,
hydroxymethyl, nitro, propenyl, propynyl and
5 trifluoromethyl and unsubstituted or substituted
heteroaryl selected

from thienyl, furanyl, pyridyl, imidazolyl,
and pyrazolyl;

wherein R³ is selected from H, lower alkyl, optionally
10 substituted phenyl, optionally substituted 3-6 membered
heterocyclyl, optionally substituted C₃-C₆-cycloalkyl,
optionally substituted phenylalkyl, optionally
substituted 3-6 membered heterocyclylalkyl, optionally
substituted C₃-C₆ cycloalkylalkyl, and lower haloalkyl,
15 preferably H, C₁₋₄-alkyl, optionally substituted phenyl,
optionally substituted phenyl-C₁₋₄-alkyl, optionally
substituted 4-6 membered heterocyclyl, optionally
substituted 4-6 membered heterocyclyl-C₁₋₄-alkyl,
optionally substituted C₃-C₆ cycloalkyl and C₁₋₂-
20 haloalkyl,
more preferably H, methyl, phenyl, cyclopropyl,
cyclohexyl, benzyl, morpholinylmethyl, 4-
methylpiperazinylmethyl, azetidinyll,
azetidinyllmethyl, 4-methylpiperdinyllmethyl, 4-
25 morpholinylmethyl, 4-morpholinylethyl, 1-(4-
morpholinyl)-2,2-dimethylpropyl, 1-piperdinyllmethyl,
1-piperdinyllpropyl, 1-pyrrolidinylpropyl and
trifluoromethyl;

wherein R⁴ is independently selected from a direct bond, C₂-
30 ₄-alkylenyl, C₂₋₄-alkenylenyl and C₂₋₄-alkynylenyl, where
one of the CH₂ groups may be replaced with an oxygen atom
or an -NH-, wherein R⁴ is optionally substituted with
hydroxy,
preferably a direct bond or R^{4a};

- 18 -

wherein R^{4a} is selected from C_{2-4} -alkylenyl where one of the CH_2 groups may be replaced with an oxygen atom or an $-NH-$, wherein R^{4a} is optionally substituted with hydroxy;



- 5 wherein R^5 is selected from H, lower alkyl, optionally substituted phenyl and optionally substituted lower aralkyl,
preferably H, methyl or ethyl,
more preferably H;
- 10 wherein R^{5a} is selected from H, lower alkyl, optionally substituted phenyl and lower aralkyl,
preferably H, methyl or ethyl,
more preferably H;
- wherein R^7 is selected from H, C_{1-6} -alkyl, optionally substituted phenyl, optionally substituted phenyl- C_{1-6} -alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl- C_{1-6} -alkyl, C_{1-4} -alkoxy- C_{1-4} -alkyl
- 15 and C_{1-4} -alkoxy- C_{1-4} -alkoxy- C_{1-4} -alkyl,
preferably H, C_{1-3} -alkyl, optionally substituted phenyl, optionally substituted phenyl- C_{1-3} -alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl- C_{1-3} -alkyl, C_{1-3} -alkoxy- C_{1-2} -alkyl and C_{1-3} -alkoxy- C_{1-3} -alkoxy- C_{1-3} -alkyl;
- 20 wherein R^c is selected from H, methyl and optionally substituted phenyl; and
- wherein R^e and R^f are independently selected from H and C_{1-2} -haloalkyl, preferably $-CF_3$;
- 30 wherein R^{14} is selected from H, optionally substituted phenyl, optionally substituted 4-6 membered heterocyclyl and optionally substituted C_3-C_6 cycloalkyl;

- 19 -


provided A is not pyridyl when X is -C(O)NH- and when R¹ is 4-[3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl]phenyl when R⁵ is methyl and when R is 4-methylpiperidyl; further provided A is not pyridyl when X is -C(O)NH-, when
5 R⁵ is H, when R² is 6-methyl and when R is indazolyl; further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl, 4-bromophenyl, 2-methylphenyl, 4-methoxyphenyl, when R⁵ is H and when R is 4-pyridyl; further provided A is not phenyl when X is -C(O)NH-, when R¹
10 is phenyl, when R⁵ is H and when R is 2-oxobenzopyran-4-yl; further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl, 4-chlorophenyl, 3-nitrophenyl, 4-methoxyphenyl, when R⁵ is H and when R is 4-imidazolinyll; further provided A is not phenyl when X is -C(O)NH-, when R⁵
15 is H, when R⁵ is substituted benzyl and when R is substituted triazinyl; further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl or 2-chlorophenyl, when R⁵ is H and when R is 4-quinazolinyl; further provided A is not phenyl when X is -C(O)NH-, when R¹
20 is phenyl, when R⁵ is H and when R is isoquinolin-1-yl; further provided A is not phenyl when X is -C(O)NH-, when R¹ is 2-chlorophenyl or 4-chlorophenyl, when R⁵ is H and when R is 3-chloroisoquinolin-1-yl; further provided A is not phenyl when X is -C(O)NH-, when R¹
25 is 1-ethylpiperid-3-yl or 1-ethylpiperid-4-yl, when R⁵ is H and when R is 8-trifluoromethylquinolin-4-yl; further provided A is not phenyl when X is -C(O)NH-, when R¹ is 1-ethylpiperid-3-yl, when R⁵ is H and when R is 8-chloroquinolin-4-yl; further provided A is not phenyl when X is -C(O)NH-, when R¹
30 is halo substituted phenyl, 1-butylpiperid-4-yl, 1-

- 20 -

ethylpiperid-3-yl or 1-ethylpiperid-4-yl, when R⁵ is H
and when R is 7-chloroquinolin-4-yl; and
further provided R is not unsubstituted 2-thienyl,
unsubstituted 2-pyridyl or unsubstituted 3-pyridyl.

5

The invention also relates to compounds of Formula II




wherein R is selected from unsubstituted or substituted 9-
10 or 10-membered fused nitrogen-containing heteroaryl,
preferably indolyl, isoindolyl, indazolyl,
quinolyl, isoquinolyl, naphthyridinyl and
quinoxalinyl,
more preferably 6-indazolyl,
15 where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
alkoxy-C₁₋₆-alkoxy, optionally substituted
20 heterocyclyl-C₂₋₄-alkynyl, and optionally substituted
heterocyclylalkoxy,
preferably chloro, fluoro, amino, hydroxy, methyl,
ethyl, propyl, trifluoromethyl,
dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
25 dimethylaminoethoxyethoxy, methoxy and ethoxy;
wherein R¹ is selected from unsubstituted or substituted
aryl,
preferably phenyl, tetrahydronaphthyl, indanyl,
indenyl and naphthyl,

- 21 -

cycloalkyl,
preferably cyclohexyl,
4-6 membered heterocyclyl,
preferably isoxazolyl, pyrazolyl, thiazolyl,
5 thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and
9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclyl,
preferably 1,2-dihydroquinolyl, 1,2,3,4-
10 tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
1H-indolyl, naphthyridinyl, quinoxalinyl,
benzo[d]isothiazolyl, 2,3,4,4a,9,9a-
hexahydro-1H-3-aza-fluorenyl, 5,6,7-
15 trihydro-1,2,4-triazolo[3,4-a]isoquinolyl,
tetrahydroquinolinyl, indazolyl, 2,1,3-
benzothiadiazolyl, benzodioxanyl,
benzothienyl, benzofuryl, benzimidazolyl,
benzoxazolyl and benzthiazolyl;
20 wherein substituted R¹ is substituted with one or more
substituents selected from halo, C₁₋₄-alkyl, optionally
substituted C₃₋₆-cycloalkyl, optionally substituted
phenyl, optionally substituted phenyl-C₁-C₄-alkylenyl,
C₁₋₂-haloalkoxy, optionally substituted phenyloxy,
25 optionally substituted 4-6 membered heterocyclyl-C₁-C₄-
alkylenyl, optionally substituted 4-6 membered
heterocyclyl-C₂-C₄-alkenylenyl, optionally substituted
4-6 membered heterocyclyl, optionally substituted 4-6
membered heterocycliloxy, optionally substituted 4-6
30 membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
optionally substituted 4-6 membered heterocyclylamino,
optionally substituted 4-6 membered
heterocyclylcarbonyl, optionally substituted 4-6

- 22 -

membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
 C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
 5 alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
 alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋₄-
 alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy,
 preferably bromo, chloro, fluoro, iodo, nitro, amino,
 cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo,
 10 aminosulfonyl, 4-methylpiperazinylsulfonyl,
 cyclohexyl, phenyl, phenylmethyl,
 morpholinylmethyl, 1-methylpiperazin-4-ylmethyl, 1-
 methylpiperazin-4-ylpropyl, morpholinylpropyl,
 piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl,
 15 2-methyl-2-(1-methylpiperidin-4-yl)ethyl,
 morpholinylethyl, 1-(4-morpholinyl)-2,2-
 dimethylpropyl, piperidin-4-ylethyl, 1-Boc-
 piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-
 piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-
 20 piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-
 piperidin-4-ylpropyl, piperidin-1-ylpropyl,
 pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-
 Boc-pyrrolidin-2-ylpropyl, pyrrolidin-1-ylmethyl,
 pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl,
 25 pyrrolidinylpropenyl, pyrrolidinylbutenyl,
 fluorosulfonyl, methylsulfonyl, methylcarbonyl,
 Boc, piperidin-1-ylmethylcarbonyl, 4-
 methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,
 aminomethylcarbonyl, dimethylaminomethylcarbonyl,
 30 3-ethoxycarbonyl-2-methyl-fur-5-yl, 4-
 methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-
 4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-
 yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),

- 23 -


imidazolyl, morpholinyl, 4-trifluoromethyl-1-
piperidinyl, hydroxybutyl, methyl, ethyl, propyl,
isopropyl, butyl, tert-butyl, sec-butyl,
trifluoromethyl, pentafluoroethyl, nonafluorobutyl,
5 dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-
hydroxymethyl, 1,1-di(trifluoromethyl)-1-
(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-
1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-
hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-
10 aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-
chlorophenoxy, phenyloxy, azetidin-3-ylmethoxy, 1-
Boc-azetidin-3-ylmethoxy, pyrrol-2-ylmethoxy, 1-
Boc-pyrrol-2-ylmethoxy, pyrrol-1-ylmethoxy, 1-
15 methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-2-
ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-
ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy,
methoxy and ethoxy;


wherein R² is one or more substituents independently
20 selected from

H,
halo,
hydroxy,
amino,
25 C₁₋₆-alkyl,
C₁₋₆-haloalkyl,
C₁₋₆-alkoxy,
C₁₋₂-alkylamino,
aminosulfonyl,
30 C₃₋₆-cycloalkyl,
cyano,
C₁₋₂-hydroxyalkyl,
nitro,
C₂₋₃-alkenyl,

- 24 -

C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 5 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 heterocyclyl;
 preferably H, chloro, fluoro, bromo, amino, hydroxy,
 methyl, ethyl, propyl, oxo, dimethylamino,
 10 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
 nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
 trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 substituted phenyl and unsubstituted or
 15 substituted heteroaryl selected from
 thienyl, furanyl, pyridyl, imidazolyl, and
 pyrazolyl;
 wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and




20 preferably direct bond, ethyl, butyl, and ;
 wherein R^e and R^f are independently selected from H and C₁₋₂-
 haloalkyl,
 preferably -CF₃; and
 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally
 25 substituted phenyl, optionally substituted phenyl-C₁₋₃-
 alkyl, optionally substituted 4-6 membered
 heterocyclyl, optionally substituted 4-6 membered
 heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-
 alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl.

30

The invention also relates to compounds of Formula III

- 25 -



III

wherein R is selected from

unsubstituted or substituted 9- or 10-membered fused

5 nitrogen-containing heteroaryl,

preferably indolyl, isoindolyl, indazolyl,
quinolyl, isoquinolyl, naphthyridinyl and
quinoxalinyl,

more preferably 5-indazolyl and 6-indazolyl,

10 even more preferably 6-indazolyl,

where R is substituted with one or more substituents

selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-

15 alkoxy-C₁₋₆-alkoxy, optionally substituted

heterocyclyl-C₂₋₄-alkynyl, and optionally substituted
heterocyclylalkoxy,

preferably chloro, fluoro, amino, hydroxy, methyl,
ethyl, propyl, trifluoromethyl,

20 dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
dimethylaminoethoxyethoxy, methoxy and ethoxy;

wherein R¹ is selected from unsubstituted or substituted
aryl,

preferably phenyl, tetrahydronaphthyl, indanyl,

25 indenyl and naphthyl,

cycloalkyl,

preferably cyclohexyl,

4-6 membered heterocyclyl,

- 26 -

preferably isoxazolyl, pyrazolyl, thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl and pyridazinyl, and

9-10 membered bicyclic and 13-14 membered

5 tricyclic heterocyclyl,

preferably 1,2-dihydroquinolyl, 1,2,3,4-

tetrahydro-isoquinolyl, isoquinolyl,

quinolyl, indolyl, isoindolyl, 2,3-dihydro-

1H-indolyl, naphthyridinyl, quinoxalinyl,

10 benzo[d]isothiazolyl, 2,3,4,4a,9,9a-

hexahydro-1H-3-aza-fluorenyl, 5,6,7-

trihydro-1,2,4-triazolo[3,4-a]isoquinolyl,

tetrahydroquinolinyl, indazolyl, 2,1,3-

benzothiadiazolyl, benzodioxanyl,

15 benzothienyl, benzofuryl, benzimidazolyl,

benzoxazolyl and benzthiazolyl;

wherein R¹ is substituted with one or more substituents

selected from halo, C₁₋₄-alkyl, optionally substituted

C₃₋₆-cycloalkyl, optionally substituted phenyl,

20 optionally substituted phenyl-C₁-C₄-alkylenyl, C₁₋₂-

haloalkoxy, optionally substituted phenyloxy,

optionally substituted 4-6 membered heterocyclyl-C₁-C₄-

alkyl, optionally substituted 4-6 membered

heterocyclyl-C₂-C₄-alkenylenyl, optionally substituted

25 4-6 membered heterocyclyl, optionally substituted 4-6

membered heterocycliloxy, optionally substituted 4-6

membered heterocyclyl-C₁₋₄-alkoxy, optionally

substituted 4-6 membered heterocyclylsulfonyl,

optionally substituted 4-6 membered heterocyclylamino,

30 optionally substituted 4-6 membered

heterocyclylcarbonyl, optionally substituted 4-6


membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-

haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,

cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,

- 27 -

C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy,

- 5 preferably bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, methylpiperazinylmethyl, 10 methylpiperazinylpropyl, morpholinylpropyl, methylpiperidinylmethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidinylethyl, piperidinylmethyl, piperidinylpropyl, pyrrolidinylpropyl, pyrrolidinylpropenyl, 15 pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, piperidinylmethylcarbonyl, methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, 20 methylpiperazinyl, methylpiperidyl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, 25 pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, 30 hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-

- 28 -

chlorophenoxy, phenyloxy, 1-methylpiperidin-4-yloxy,
isopropoxy, methoxy and ethoxy;


wherein R² is one or more substituents independently
selected from

- 5 H,
 halo,
 hydroxy,
 amino,
 C₁₋₆-alkyl,
10 C₁₋₆-haloalkyl,
 C₁₋₆-alkoxy,
 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
15 cyano,
 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
20 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
25 heterocyclyl;
 preferably H, chloro, fluoro, bromo, amino, hydroxy,
 methyl, ethyl, propyl, oxo, dimethylamino,
 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
 nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
30 trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 substituted phenyl and unsubstituted or
 substituted heteroaryl selected from


- 29 -

thienyl, furanyl, pyridyl, imidazolyl, and
pyrazolyl;

wherein R^4 is selected from a direct bond, C_{1-4} -alkyl, and



;




;

5 preferably direct bond, ethyl, butyl, and
wherein R^e and R^f are independently selected from H and C_{1-2} -
haloalkyl,
preferably $-CF_3$; and

wherein R^7 is selected from H, C_{1-3} -alkyl, optionally
10 substituted phenyl- C_{1-3} -alkyl, 4-6 membered
heterocyclyl, optionally substituted 4-6 membered
heterocyclyl- C_{1-3} -alkyl, C_{1-3} -alkoxy- C_{1-2} -alkyl and C_{1-3} -
alkoxy- C_{1-3} -alkoxy- C_{1-3} -alkyl.

15 The invention also relates to compounds of Formula IV



IV

wherein A^3 is selected from CR^2 and N;

wherein A^4 is selected from CR^2 and N; provided one of A^3 and

20 A^4 is not CR^2 ;

wherein R is selected from

unsubstituted or substituted 9- or 10-membered fused
nitrogen-containing heteroaryl,

preferably indolyl, isoindolyl, indazolyl,

25 quinolyl, isoquinolyl, naphthyridinyl and
quinoxalinyl,

more preferably 5-indazolyl and 6-indazolyl,


- 30 -

even more preferably 6-indazolyl,
where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
5 C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
alkoxy-C₁₋₆-alkoxy, optionally substituted
heterocyclyl-C₂₋₄-alkynyl, and optionally substituted
heterocyclylalkoxy,
preferably chloro, fluoro, amino, hydroxy, methyl,
10 ethyl, propyl, trifluoromethyl,
dimethylaminopropynyl, 1-methylpiperdinylmethoxy,
dimethylaminoethoxyethoxy, methoxy and ethoxy;
wherein R¹ is selected from unsubstituted or substituted
aryl,
15 preferably phenyl, tetrahydronaphthyl, indanyl,
indenyl and naphthyl,
cycloalkyl,
preferably cyclohexyl,
4-6 membered heterocyclyl,
20 preferably isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and
9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclyl,
25 preferably 1,2-dihydroquinolyl, 1,2,3,4-
tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
1H-indolyl, naphthyridinyl, quinoxalinyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
30 5,6,7-trihydro-1,2,4-triazolo[3,4-
a]isoquinolyl, tetrahydroquinolinyl,
indazolyl, benzo[d]isothiazolyl, 2,1,3-
benzothiadiazolyl, benzodioxanyl,

- 31 -

benzothienyl, benzofuryl, benzimidazolyl,
benzoxazolyl and benzthiazolyl;

wherein R¹ is substituted with one or more substituents
selected from halo, C₁₋₄-alkyl, optionally substituted
5 C₃₋₆-cycloalkyl, optionally substituted phenyl,
optionally substituted phenyl-C₁₋₄-alkylenyl, C₁₋₂-
haloalkoxy, optionally substituted phenyloxy,
optionally substituted 4-6 membered heterocyclyl-C₁₋₄-
alkylenyl, optionally substituted 4-6 membered
10 heterocyclyl-C₂₋₄-alkenylenyl, optionally substituted
4-6 membered heterocyclyl, optionally substituted 4-6
membered heterocyclyloxy, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
15 optionally substituted 4-6 membered heterocyclylamino,
optionally substituted 4-6 membered
heterocyclylcarbonyl, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
20 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy,

25 preferably bromo, chloro, fluoro, iodo, nitro, amino,
cyano, aminoethyl, Boc-aminoethyl, hydroxy,
aminosulfonyl, 4-methylpiperazinylsulfonyl,
cyclohexyl, phenyl, phenylmethyl,
morpholinylmethyl, methylpiperazinylmethyl,
30 methylpiperazinylpropyl, morpholinylpropyl,
methylpiperidinylmethyl, morpholinylethyl, 1-(4-
morpholinyl)-2,2-dimethylpropyl, piperidinylethyl,
piperidinylmethyl, piperidinylpropyl,



- 32 -

pyrrolidinylpropyl, pyrrolidinylpropenyl,
pyrrolidinylbutenyl, fluorosulfonyl,
methylsulfonyl, methylcarbonyl,
piperidinylmethylcarbonyl,
5 methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-
ethoxycarbonyl-2-methyl-fur-5-yl,
methylpiperazinyl, methylpiperidyl, 1-methyl-
(1,2,3,6-tetrahydropyridyl), imidazolyl,
morpholinyl, 4-trifluoromethyl-1-piperidinyl,
10 hydroxybutyl, methyl, ethyl, propyl, isopropyl,
butyl, tert-butyl, sec-butyl, trifluoromethyl,
pentafluoroethyl, nonafluorobutyl,
dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-
hydroxymethyl, 1,1-di(trifluoromethyl)-1-
15 (piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-
1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-
hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-
aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-
20 chlorophenoxy, phenyloxy, 1-methylpiperidin-4-yloxy,
isopropoxy, methoxy and ethoxy;

wherein R² is one or more substituents independently
selected from


H,
25 halo,
hydroxy,
amino,
C₁₋₆-alkyl,
C₁₋₆-haloalkyl,
30 C₁₋₆-alkoxy,
C₁₋₂-alkylamino,
aminosulfonyl,
C₃₋₆-cycloalkyl,
cyano,

- 33 -

C_{1-2} -hydroxyalkyl,
 nitro,
 C_{2-3} -alkenyl,
 C_{2-3} -alkynyl,
 5 C_{1-6} -haloalkoxy,
 C_{1-6} -carboxyalkyl,
 5-6-membered heterocyclyl- C_{1-6} -alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 10 heterocyclyl;
 preferably H, chloro, fluoro, bromo, amino, hydroxy,
 methyl, ethyl, propyl, oxo, dimethylamino,
 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
 nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
 15 trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 substituted phenyl and unsubstituted or
 substituted heteroaryl selected from
 thienyl, furanyl, pyridyl, imidazolyl, and
 20 pyrazolyl;
 wherein R^4 is selected from a direct bond, C_{1-4} -alkyl, and
 ;
 preferably direct bond, ethyl, butyl, and  ;
 wherein R^e and R^f are independently selected from H and C_{1-2} -
 25 haloalkyl,
 preferably $-CF_3$; and
 wherein R^7 is selected from H, C_{1-3} -alkyl, optionally
 substituted phenyl- C_{1-3} -alkyl, 4-6 membered
 heterocyclyl, optionally substituted 4-6 membered
 30 heterocyclyl- C_{1-3} -alkyl, C_{1-3} -alkoxy- C_{1-2} -alkyl and C_{1-3} -
 alkoxy- C_{1-3} -alkoxy- C_{1-3} -alkyl.

- 34 -

The invention also relates to compounds of Formula V



wherein A⁵ is selected from S, O and NR⁶;

5 wherein R is selected from

unsubstituted or substituted 9- or 10-membered fused
nitrogen-containing heteroaryl,
preferably indolyl, isoindolyl, indazolyl,
quinolyl, isoquinolyl, naphthyridinyl and
10 quinozaliny, more preferably 5-indazolyl and 6-indazolyl,
even more preferably 6-indazolyl,

where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
15 haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
alkoxy-C₁₋₆-alkoxy, optionally substituted
heterocyclyl-C₂₋₄-alkynyl, and optionally substituted
heterocyclylalkoxy,

20 preferably chloro, fluoro, amino, hydroxy, methyl,
ethyl, propyl, trifluoromethyl,
dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
dimethylaminoethoxyethoxy, methoxy and ethoxy;

wherein R¹ is selected from unsubstituted or substituted
25 aryl,

preferably phenyl, tetrahydronaphthyl, indanyl,
indenyl and naphthyl,
cycloalkyl,
preferably cyclohexyl,


- 35 -


4-6 membered heterocyclyl,
preferably isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and

5 9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclyl,
preferably 1,2-dihydroquinolyl, 1,2,3,4-
tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
10 1H-indolyl, naphthyridinyl, quinoxalyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
5,6,7-trihydro-1,2,4-triazolo[3,4-
a]isoquinolyl, benzo[d]isothiazolyl,
tetrahydroquinolinyl, indazolyl, 2,1,3-
15 benzothiadiazolyl, benzodioxanyl,
benzothienyl, benzofuryl, benzimidazolyl,
benzoxazolyl and benzthiazolyl;

wherein R¹ is substituted with one or more substituents
selected from halo, C₁₋₄-alkyl, optionally substituted
20 C₃₋₆-cycloalkyl, optionally substituted phenyl,
optionally substituted phenyl-C₁-C₄-alkylenyl, C₁₋₂-
haloalkoxy, optionally substituted phenyloxy,
optionally substituted 4-6 membered heterocyclyl-C₁-C₄-
alkylenyl, optionally substituted 4-6 membered
25 heterocyclyl-C₂-C₄-alkenylenyl, optionally substituted
4-6 membered heterocyclyl, optionally substituted 4-6
membered heterocyclyoxy, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
30 optionally substituted 4-6 membered heterocyclylamino,
optionally substituted 4-6 membered
heterocyclylcarbonyl, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,

cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋



5 4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy, preferably bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, methylpiperazinylmethyl, methylpiperazinylpropyl, morpholinylpropyl, methylpiperidinylmethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidinylethyl, piperidinylmethyl, piperidinylpropyl, pyrrolidinylpropyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, piperidinylmethylcarbonyl, methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, methylpiperazinyl, methylpiperidyl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-

- 37 -

chlorophenoxy, phenyloxy, 1-methylpiperidin-4-yloxy,
isopropoxy, methoxy and ethoxy;


wherein R² is one or more substituents independently
selected from


- 5 H,
 halo,
 hydroxy,
 amino,
 C₁₋₆-alkyl,
10 C₁₋₆-haloalkyl,
 C₁₋₆-alkoxy,
 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
15 cyano,
 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
20 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
25 heterocyclyl;
 preferably H, chloro, fluoro, bromo, amino, hydroxy,
 methyl, ethyl, propyl, oxo, dimethylamino,
 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
 nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
30 trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 substituted phenyl and unsubstituted or
 substituted heteroaryl selected from

- 38 -

thienyl, furanyl, pyridyl, imidazolyl, and
pyrazolyl;

wherein R^4 is selected from a direct bond, C_{1-4} -alkyl, and




5 preferably direct bond, ethyl, butyl, and  ;
wherein R^e and R^f are independently selected from H and C_{1-2} -
haloalkyl,
preferably $-CF_3$;

wherein R^6 is H or C_{1-2} -alkyl, preferably H or methyl; and

10 wherein R^7 is selected from H, C_{1-3} -alkyl, optionally
substituted phenyl- C_{1-3} -alkyl, 4-6 membered
heterocyclyl, optionally substituted 4-6 membered
heterocyclyl- C_{1-3} -alkyl, C_{1-3} -alkoxy- C_{1-2} -alkyl and C_{1-3} -
alkoxy- C_{1-3} -alkoxy- C_{1-3} -alkyl.

15

The invention also relates to compounds of Formula VI



VI

wherein A^5 is selected from S, O and NR^6 ;

20 wherein R is selected from

unsubstituted or substituted 9- or 10-membered fused
nitrogen-containing heteroaryl,
preferably indolyl, isoindolyl, indazolyl,
quinolyl, isoquinolyl, naphthyridinyl and
25 quinoxalinyl,
more preferably 5-indazolyl and 6-indazolyl,


- 39 -

even more preferably 6-indazolyl,
where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
5 C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
alkoxy-C₁₋₆-alkoxy, optionally substituted
heterocyclyl-C₂₋₄-alkynyl, and optionally substituted
heterocyclylalkoxy,
preferably chloro, fluoro, amino, hydroxy, methyl,
10 ethyl, propyl, trifluoromethyl,
dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
dimethylaminoethoxyethoxy, methoxy and ethoxy;
wherein R¹ is selected from unsubstituted or substituted
aryl,
15 preferably phenyl, tetrahydronaphthyl, indanyl,
indenyl and naphthyl,
cycloalkyl,
preferably cyclohexyl,
4-6 membered heterocyclyl,
20 preferably isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and
9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclyl,
25 preferably 1,2-dihydroquinolyl, 1,2,3,4-
tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
1H-indolyl, naphthyridinyl, quinoxalinyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
30 5,6,7-trihydro-1,2,4-triazolo[3,4-
a]isoquinolyl, benzo[d]isothiazolyl,
tetrahydroquinolinyl, indazolyl, 2,1,3-
benzothiadiazolyl, benzodioxanyl,

- 40 -

benzothienyl, benzofuryl, benzimidazolyl,
benzoxazolyl and benzthiazolyl;

wherein R¹ is substituted with one or more substituents
selected from halo, C₁₋₄-alkyl, optionally substituted
5 C₃₋₆-cycloalkyl, optionally substituted phenyl,
optionally substituted phenyl-C_{1-C4}-alkylenyl, C₁₋₂-
haloalkoxy, optionally substituted phenyloxy,
optionally substituted 4-6 membered heterocyclyl-C_{1-C4}-
alkylenyl, optionally substituted 4-6 membered
10 heterocyclyl-C_{2-C4}-alkenylenyl, optionally substituted
4-6 membered heterocyclyl, optionally substituted 4-6
membered heterocyclyloxy, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
15 optionally substituted 4-6 membered heterocyclylamino,
optionally substituted 4-6 membered
heterocyclylcarbonyl, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
20 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy,

25 preferably bromo, chloro, fluoro, iodo, nitro, amino,
cyano, aminoethyl, Boc-aminoethyl, hydroxy,
aminosulfonyl, 4-methylpiperazinylsulfonyl,
cyclohexyl, phenyl, phenylmethyl,
morpholinylmethyl, methylpiperazinylmethyl,
30 methylpiperazinylpropyl, morpholinylpropyl,
methylpiperidinylmethyl, morpholinylethyl, 1-(4-
morpholinyl)-2,2-dimethylpropyl, piperidinylethyl,
piperidinylmethyl, piperidinylpropyl,

- 41 -

pyrrolidinylpropyl, pyrrolidinylpropenyl,
pyrrolidinylbutenyl, fluorosulfonyl,
methylsulfonyl, methylcarbonyl,
piperidinylmethylcarbonyl,
5 methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-
ethoxycarbonyl-2-methyl-fur-5-yl,
methylpiperazinyl, methylpiperidyl, 1-methyl-
(1,2,3,6-tetrahydropyridyl), imidazolyl,
morpholinyl, 4-trifluoromethyl-1-piperidinyl,
10 hydroxybutyl, methyl, ethyl, propyl, isopropyl,
butyl, tert-butyl, sec-butyl, trifluoromethyl,
pentafluoroethyl, nonafluorobutyl,
dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-
hydroxymethyl, 1,1-di(trifluoromethyl)-1-
15 (piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-
1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-
hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-
aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-
20 chlorophenoxy, phenyloxy, 1-methylpiperidin-4-yloxy,
isopropoxy, methoxy and ethoxy;


wherein R² is one or more substituents independently
selected from


H,
25 halo,
hydroxy,
amino,
C₁₋₆-alkyl,
C₁₋₆-haloalkyl,
30 C₁₋₆-alkoxy,
C₁₋₂-alkylamino,
aminosulfonyl,
C₃₋₆-cycloalkyl,
cyano,

- 42 -

C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 5 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 10 heterocyclyl;
 preferably H, chloro, fluoro, bromo, amino, hydroxy,
 methyl, ethyl, propyl, oxo, dimethylamino,
 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
 nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
 15 trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 substituted phenyl and unsubstituted or
 substituted heteroaryl selected from
 thienyl, furanyl, pyridyl, imidazolyl, and
 20 pyrazolyl;

wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and




preferably direct bond, ethyl, butyl, and  ;
 wherein R^e and R^f are independently selected from H and C₁₋₂-
 25 haloalkyl,
 preferably -CF₃;

wherein R⁶ is H or C₁₋₂-alkyl, preferably H or methyl; and
 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally
 substituted phenyl-C₁₋₃-alkyl, 4-6 membered
 30 heterocyclyl, optionally substituted 4-6 membered
 heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-
 alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl.

- 43 -

The invention also relates to compounds of Formulas VIIa and VIIb



wherein A⁵ is selected from S, O and NR⁶;

wherein R is selected from

unsubstituted or substituted 9- or 10-membered fused

nitrogen-containing heteroaryl,

preferably indolyl, isoindolyl, indazolyl,

quinolyl, isoquinolyl, naphthyridinyl and

quinoxalinyl,

more preferably 5-indazolyl and 6-indazolyl,

even more preferably 6-indazolyl,

where R is substituted with one or more substituents

selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-

haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,

C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-

alkoxy-C₁₋₆-alkoxy, optionally substituted

heterocyclyl-C₂₋₄-alkynyl, and optionally substituted

heterocyclylalkoxy,

preferably chloro, fluoro, amino, hydroxy, methyl,

ethyl, propyl, trifluoromethyl,

dimethylaminopropynyl, 1-methylpiperidinylmethoxy,

dimethylaminoethoxyethoxy, methoxy and ethoxy;

wherein R¹ is selected from unsubstituted or substituted

aryl,

preferably phenyl, tetrahydronaphthyl, indanyl,


indenyl and naphthyl,

- 44 -

cycloalkyl,
preferably cyclohexyl,
4-6 membered heterocyclyl,
preferably isoxazolyl, pyrazolyl, thiazolyl,
5 thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and
9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclyl,
preferably 1,2-dihydroquinolyl, -1,2,3,4-
10 tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
1H-indolyl, naphthyridinyl, quinoxalinyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
5,6,7-trihydro-1,2,4-triazolo[3,4-
15 a]isoquinolyl, tetrahydroquinolinyl,
indazolyl, 2,1,3-benzothiadiazolyl,
benzo[d]isothiazolyl, benzodioxanyl,
benzothienyl, benzofuryl, benzimidazolyl,
benzoxazolyl and benzthiazolyl;
20 wherein R¹ is substituted with one or more substituents
selected from halo, C₁₋₄-alkyl, optionally substituted
C₃₋₆-cycloalkyl, optionally substituted phenyl,
optionally substituted phenyl-C_{1-C4}-alkylenyl, C₁₋₂-
haloalkoxy, optionally substituted phenyloxy,
25 optionally substituted 4-6 membered heterocyclyl-C_{1-C4}-
alkyl, optionally substituted 4-6 membered
heterocyclyl-C_{2-C4}-alkenyl, optionally substituted 4-6
membered heterocyclyl, optionally substituted 4-6
membered heterocyclyloxy, optionally substituted 4-6
30 membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
optionally substituted 4-6 membered heterocyclylamino,
optionally substituted 4-6 membered
heterocyclylcarbonyl, optionally substituted 4-6

- 45 -

membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

₄-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy, preferably bromo, chloro, fluoro, iodo, nitro, amino,

cyano, aminoethyl, Boc-aminoethyl, hydroxy, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, methylpiperazinylmethyl, methylpiperazinylpropyl, morpholinylpropyl, methylpiperidinylmethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidinylethyl, piperidinylmethyl, piperidinylpropyl, pyrrolidinylpropyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, piperidinylmethylcarbonyl, methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, methylpiperazinyl, methylpiperidyl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-

- 46 -

aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy, phenyloxy, 1-methylpiperidin-4-yloxy, isopropoxy, methoxy and ethoxy;


5 wherein R² is one or more substituents independently selected from

H,
halo,
hydroxy,
10 amino,
C₁₋₆-alkyl,
C₁₋₆-haloalkyl,
C₁₋₆-alkoxy,
C₁₋₂-alkylamino,
15 aminosulfonyl,
C₃₋₆-cycloalkyl,
cyano,
C₁₋₂-hydroxyalkyl,
nitro,
20 C₂₋₃-alkenyl,
C₂₋₃-alkynyl,
C₁₋₆-haloalkoxy,
C₁₋₆-carboxyalkyl,
5-6-membered heterocyclyl-C₁₋₆-alkylamino,
25 unsubstituted or substituted phenyl and
unsubstituted or substituted 4-6 membered
heterocyclyl;
preferably H, chloro, fluoro, bromo, amino, hydroxy,
methyl, ethyl, propyl, oxo, dimethylamino,
30 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
trifluoromethoxy, carboxymethyl,
morpholinylethylamino, propynyl, unsubstituted or


- 47 -

substituted phenyl and unsubstituted or
substituted heteroaryl selected from
thienyl, furanyl, pyridyl, imidazolyl, and
pyrazolyl;

5 wherein R^4 is selected from a direct bond, C_{1-4} -alkyl, and



preferably direct bond, ethyl, butyl, and



wherein R^5 and R^6 are independently selected from H and C_{1-2} -
haloalkyl,

10 preferably $-CF_3$;

wherein R^6 is H or C_{1-2} -alkyl, preferably H or methyl; and

wherein R^7 is selected from H, C_{1-3} -alkyl, optionally


substituted phenyl- C_{1-3} -alkyl, 4-6 membered

heterocyclyl, optionally substituted 4-6 membered

15 heterocyclyl- C_{1-3} -alkyl, C_{1-3} -alkoxy- C_{1-2} -alkyl and C_{1-3} -
alkoxy- C_{1-3} -alkoxy- C_{1-3} -alkyl.


The invention also relates to compounds of Formulas
VIIIa and VIIIb

20



VIIIa

and



VIIIb

wherein A^5 is selected from S, O and NR^6 ;

wherein R is selected from

25 unsubstituted or substituted 9- or 10-membered fused
nitrogen-containing heteroaryl,


- 48 -

preferably indolyl, isoindolyl, indazolyl,
quinolyl, isoquinolyl, naphthyridinyl and
quinoxalinyl,
more preferably 5-indazolyl and 6-indazolyl,
5 even more preferably 6-indazolyl,
where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
10 alkoxy-C₁₋₆-alkoxy, optionally substituted
heterocyclyl-C₂₋₄-alkynyl, and optionally substituted
heterocyclylalkoxy,
preferably chloro, fluoro, amino, hydroxy, methyl,
ethyl, propyl, trifluoromethyl,
15 dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
dimethylaminoethoxyethoxy, methoxy and ethoxy;
wherein R¹ is selected from unsubstituted or substituted
aryl,
preferably phenyl, tetrahydronaphthyl, indanyl,
20 indenyl and naphthyl,
cycloalkyl,
preferably cyclohexyl,
4-6 membered heterocyclyl,
preferably isoxazolyl, pyrazolyl, thiazolyl,
25 thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and
9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclyl,
preferably 1,2-dihydroquinolyl, 1,2,3,4-
30 tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
1H-indolyl, naphthyridinyl, quinoxalinyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
5,6,7-trihydro-1,2,4-triazolo[3,4-

- 49 -

a]isoquinolyl, tetrahydroquinolyl,
indazolyl, benzo[d]isothiazolyl, 2,1,3-
benzothiadiazolyl, benzodioxanyl,
benzothienyl, benzofuryl, benzimidazolyl,
5 benzoxazolyl and benzthiazolyl;

wherein R¹ is substituted with one or more substituents
selected from halo, C₁₋₄-alkyl, optionally substituted
C₃₋₆-cycloalkyl, optionally substituted phenyl,
optionally substituted phenyl-C_{1-C4}-alkylenyl, C₁₋₂-
10 haloalkoxy, optionally substituted phenoxy,
optionally substituted 4-6 membered heterocyclyl-C_{1-C4}-
alkyl, optionally substituted 4-6 membered
heterocyclyl-C_{2-C4}-alkenylenyl, optionally substituted
4-6 membered heterocyclyl, optionally substituted 4-6
15 membered heterocyclloxy, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
optionally substituted 4-6 membered heterocyclylamino,
optionally substituted 4-6 membered
20 heterocyclylcarbonyl, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
25 alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy,
preferably bromo, chloro, fluoro, iodo, nitro, amino,
cyano, aminoethyl, Boc-aminoethyl, hydroxy,
30 aminosulfonyl, 4-methylpiperazinylsulfonyl,
cyclohexyl, phenyl, phenylmethyl,
morpholinylmethyl, methylpiperazinylmethyl,
methylpiperazinylpropyl, morpholinylpropyl,

- 50 -

methylnpiperidinylmethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidinylethyl, piperidinylmethyl, piperidinylpropyl, pyrrolidinylpropyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, piperidinylmethylcarbonyl, methylnpiperazinylcarbonylethyl, methoxycarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, methylnpiperazinyl, methylnpiperidyl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy, phenyloxy, 1-methylnpiperidin-4-yloxy, isopropoxy, methoxy and ethoxy;

wherein R² is one or more substituents independently selected from


H,
halo,
hydroxy,
amino,
C₁₋₆-alkyl,
C₁₋₆-haloalkyl,
C₁₋₆-alkoxy,
C₁₋₂-alkylamino,

- 51 -

aminosulfonyl,
 C₃₋₆-cycloalkyl,
 cyano,
 C₁₋₂-hydroxyalkyl,
 5 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 10 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 heterocyclyl;
 preferably H, chloro, fluoro, bromo, amino, hydroxy,
 15 methyl, ethyl, propyl, oxo, dimethylamino,
 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
 nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
 trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 20 substituted phenyl and unsubstituted or
 substituted heteroaryl selected from
 thienyl, furanyl, pyridyl, imidazolyl, and
 pyrazolyl;

wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and




preferably direct bond, ethyl, butyl, and ;
 wherein R^e and R^f are independently selected from H and C₁₋₂-
 haloalkyl,
 preferably -CF₃;

30 wherein R⁵ is H or C₁₋₂-alkyl, preferably H or methyl; and
 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally
 substituted phenyl-C₁₋₃-alkyl, 4-6 membered

- 52 -

heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁-C₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl.

5 The invention also relates to compounds of Formula IX



IX

wherein A⁵ is selected from S, O and NR⁶;

wherein R is selected from


- 10 unsubstituted or substituted 9- or 10-membered fused nitrogen-containing heteroaryl, preferably indolyl, isoindolyl, indazolyl, quinolyl, isoquinolyl, naphthyridinyl and quinoxalinyl,
- 15 more preferably 5-indazolyl and 6-indazolyl, even more preferably 6-indazolyl,
- where R is substituted with one or more substituents selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl, C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-alkoxy-C₁₋₆-alkoxy, optionally substituted
- 20 heterocyclyl-C₂₋₄-alkynyl, and optionally substituted heterocyclylalkoxy, preferably chloro, fluoro, amino, hydroxy, methyl,
- 25 ethyl, propyl, trifluoromethyl, dimethylaminopropynyl, 1-methylpiperidinylmethoxy, dimethylaminoethoxyethoxy, methoxy and ethoxy;
- wherein R¹ is selected from unsubstituted or substituted aryl,

- 53 -

preferably phenyl, tetrahydronaphthyl, indanyl,
indenyl and naphthyl,
cycloalkyl,
preferably cyclohexyl,
5 4-6 membered heterocyclyl,
preferably isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and
9-10 membered bicyclic and 13-14 membered
10 tricyclic heterocyclyl,
preferably 1,2-dihydroquinolyl, 1,2,3,4-
tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
1H-indolyl, naphthyridinyl, quinoxalyl,
15 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
5,6,7-trihydro-1,2,4-triazolo[3,4-
a]isoquinolyl, tetrahydroquinolinyl,
indazolyl, 2,1,3-benzothiadiazolyl,
benzo[d]isothiazolyl, benzodioxanyl,
20 benzothienyl, benzofuryl, benzimidazolyl,
benzoxazolyl and benzthiazolyl;
wherein R¹ is substituted with one or more substituents
selected from halo, C₁₋₄-alkyl, optionally substituted
C₃₋₆-cycloalkyl, optionally substituted phenyl,
25 optionally substituted phenyl-C₁-C₄-alkylenyl, C₁₋₂-
haloalkoxy, optionally substituted phenyloxy,
optionally substituted 4-6 membered heterocyclyl-C₁-C₄-
alkyl, optionally substituted 4-6 membered
heterocyclyl-C₂-C₄-alkenylenyl, optionally substituted
30 4-6 membered heterocyclyl, optionally substituted 4-6
membered heterocycliloxy, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
optionally substituted 4-6 membered heterocyclylamino,

- 54 -

optionally substituted 4-6 membered
 heterocyclylcarbonyl, optionally substituted 4-6
 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
 5 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
 C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
 alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
 alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy,

10 preferably bromo, chloro, fluoro, iodo, nitro, amino,
 cyano, aminoethyl, Boc-aminoethyl, hydroxy,
 aminosulfonyl, 4-methylpiperazinylsulfonyl,
 cyclohexyl, phenyl, phenylmethyl,
 morpholinylmethyl, methylpiperazinylmethyl,
 15 methylpiperazinylpropyl, morpholinylpropyl,
 methylpiperidinylmethyl, morpholinylethyl, 1-(4-
 morpholinyl)-2,2-dimethylpropyl, piperidinylethyl,
 piperidinylmethyl, piperidinylpropyl,
 pyrrolidinylpropyl, pyrrolidinylpropenyl,
 20 pyrrolidinylbutenyl, fluorosulfonyl,
 methylsulfonyl, methylcarbonyl,
 piperidinylmethylcarbonyl,
 methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-
 ethoxycarbonyl-2-methyl-fur-5-yl,
 25 methylpiperazinyl, methylpiperidyl, 1-methyl-
 (1,2,3,6-tetrahydropyridyl), imidazolyl,
 morpholinyl, 4-trifluoromethyl-1-piperidinyl,
 hydroxybutyl, methyl, ethyl, propyl, isopropyl,
 butyl, tert-butyl, sec-butyl, trifluoromethyl,
 30 pentafluoroethyl, nonafluorobutyl,
 dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-
 hydroxymethyl, 1,1-di(trifluoromethyl)-1-
 (piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-

- 55 -

1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy, phenyloxy, 1-methylpiperidin-4-yloxy, isopropoxy, methoxy and ethoxy;


wherein R² is one or more substituents independently selected from

H,
halo,
hydroxy,
amino,
C₁₋₆-alkyl,
C₁₋₆-haloalkyl,
C₁₋₆-alkoxy,
C₁₋₂-alkylamino,
aminosulfonyl,
C₃₋₆-cycloalkyl,
cyano,
C₁₋₂-hydroxyalkyl,
nitro,
C₂₋₃-alkenyl,
C₂₋₃-alkynyl,
C₁₋₆-haloalkoxy,
C₁₋₆-carboxyalkyl,
5-6-membered heterocyclyl-C₁₋₆-alkylamino,
unsubstituted or substituted phenyl and
unsubstituted or substituted 4-6 membered
heterocyclyl;
preferably H, chloro, fluoro, bromo, amino, hydroxy,
methyl, ethyl, propyl, oxo, dimethylamino,
aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
trifluoromethoxy, carboxymethyl,


- 56 -

morpholinylethylamino, propynyl, unsubstituted or substituted phenyl and unsubstituted or substituted heteroaryl selected from thienyl, furanyl, pyridyl, imidazolyl, and pyrazolyl;

wherein R^4 is selected from a direct bond, C_{1-4} -alkyl, and



preferably direct bond, ethyl, butyl, and




wherein R^a and R^f are independently selected from H and C_{1-2} -haloalkyl, preferably $-CF_3$;

wherein R^6 is H or C_{1-2} -alkyl, preferably H or methyl; and

wherein R^7 is selected from H, C_{1-3} -alkyl, optionally substituted phenyl- C_{1-3} -alkyl, 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl- C_{1-3} -alkyl, C_{1-3} -alkoxy- C_{1-2} -alkyl and C_{1-3} -alkoxy- C_{1-3} -alkoxy- C_{1-3} -alkyl.

The invention also relates to compounds of Formula X



X

wherein A^5 is selected from S, O and NR^6 ;

wherein A^6 is selected from N and CR^2 ;

wherein R is selected from


- 57 -

unsubstituted or substituted 9- or 10-membered fused
nitrogen-containing heteroaryl,
preferably indolyl, isoindolyl, indazolyl,
quinolyl, isoquinolyl, naphthyridinyl and
5 quinozaliny, more preferably 5-indazolyl and 6-indazolyl,
even more preferably 6-indazolyl,
where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
10 haloalkyl, C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
alkoxy-C₁₋₆-alkoxy, optionally substituted
heterocyclyl-C₂₋₄-alkynyl, and optionally substituted
heterocyclylalkoxy,
15 preferably chloro, fluoro, amino, hydroxy, methyl,
ethyl, propyl, trifluoromethyl,
dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
dimethylaminoethoxyethoxy, methoxy and ethoxy;
wherein R¹ is selected from unsubstituted or substituted
20 aryl,
preferably phenyl, tetrahydronaphthyl, indanyl,
indenyl and naphthyl,
cycloalkyl,
preferably cyclohexyl,
25 4-6 membered heterocyclyl,
preferably isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl
and pyridazinyl, and
9-10 membered bicyclic and 13-14 membered
30 tricyclic heterocyclyl,
preferably 1,2-dihydroquinolyl, 1,2,3,4-
tetrahydro-isoquinolyl, isoquinolyl,
quinolyl, indolyl, isoindolyl, 2,3-dihydro-
1H-indolyl, naphthyridinyl, quinozaliny,

- 58 -

2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl,
 5,6,7-trihydro-1,2,4-triazolo[3,4-
 a]isoquinolyl, tetrahydroquinoliny,
 indazolyl, 2,1,3-benzothiadiazolyl,
 benzo[d]isothiazolyl, benzodioxanyl,
 benzothienyl, benzofuryl, benzimidazolyl,
 benzoxazolyl and benzthiazolyl;

wherein R¹ is substituted with one or more substituents
 selected from halo, C₁₋₄-alkyl, optionally substituted
 C₃₋₆-cycloalkyl, optionally substituted phenyl,
 optionally substituted phenyl-C₁-C₄-alkylenyl, C₁₋₂-
 haloalkoxy, optionally substituted phenyloxy,
 optionally substituted 4-6 membered heterocyclyl-C₁-C₄-
 alkylenyl, optionally substituted 4-6 membered
 heterocyclyl-C₂-C₄-alkenylenyl, optionally substituted
 4-6 membered heterocyclyl, optionally substituted 4-6
 membered heterocycliloxy, optionally substituted 4-6
 membered heterocyclyl-C₁₋₄-alkoxy, optionally
 substituted 4-6 membered heterocyclylsulfonyl,
 optionally substituted 4-6 membered heterocyclylamino,
 optionally substituted 4-6 membered
 heterocyclylcarbonyl, optionally substituted 4-6
 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
 C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
 alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
 alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy,
 preferably bromo, chloro, fluoro, iodo, nitro, amino,
 cyano, aminoethyl, Boc-aminoethyl, hydroxy,
 aminosulfonyl, 4-methylpiperazinylsulfonyl,
 cyclohexyl, phenyl, phenylmethyl,

- 59 -

morpholinylmethyl, methylpiperazinylmethyl,
methylpiperazinylpropyl, morpholinylpropyl,
methylpiperidinylmethyl, morpholinylethyl, 1-(4-
morpholinyl)-2,2-dimethylpropyl, piperidinylethyl,
5 piperidinylmethyl, piperidinylpropyl,
pyrrolidinylpropyl, pyrrolidinylpropenyl,
pyrrolidinylbutenyl, fluorosulfonyl,
methylsulfonyl, methylcarbonyl,
piperidinylmethylcarbonyl,
10 methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-
ethoxycarbonyl-2-methyl-fur-5-yl,
methylpiperazinyl, methylpiperidyl, 1-methyl-
(1,2,3,6-tetrahydropyridyl), imidazolyl,
morpholinyl, 4-trifluoromethyl-1-piperidinyl,
15 hydroxybutyl, methyl, ethyl, propyl, isopropyl,
butyl, tert-butyl, sec-butyl, trifluoromethyl,
pentafluoroethyl, nonafluorobutyl,
dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-
hydroxymethyl, 1,1-di(trifluoromethyl)-1-
20 (piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-
1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-
hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-
aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-
25 chlorophenoxy, phenyloxy, 1-methylpiperidin-4-yloxy,
isopropoxy, methoxy and ethoxy;


wherein R² is one or more substituents independently
selected from


30 H,
halo,
hydroxy,
amino,
C₁₋₆-alkyl,
C₁₋₆-haloalkyl,

- 60 -

C₁₋₆-alkoxy,
 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 5 cyano,
 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 10 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 15 heterocyclyl;
 preferably H, chloro, fluoro, bromo, amino, hydroxy,
 methyl, ethyl, propyl, oxo, dimethylamino,
 aminosulfonyl, cyclopropyl, cyano, hydroxymethyl,
 nitro, propenyl, trifluoromethyl, methoxy, ethoxy,
 20 trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 substituted phenyl and unsubstituted or
 substituted heteroaryl selected from
 thienyl, furanyl, pyridyl, imidazolyl, and
 25 pyrazolyl;

wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and



preferably direct bond, ethyl, butyl, and  ;
 wherein R^e and R^f are independently selected from H and C₁₋₂-

30 haloalkyl,








preferably -CF₃;

wherein R⁶ is H or C₁₋₂-alkyl, preferably H or methyl;

- 61 -

wherein R^7 is selected from H, C_{1-3} -alkyl, optionally substituted phenyl- C_{1-3} -alkyl, 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl- C_{1-3} -alkyl, C_{1-3} -alkoxy- C_{1-2} -alkyl and C_{1-3} -alkoxy- C_{1-3} -alkoxy- C_{1-3} -alkyl; and

wherein



A family of specific compounds of particular interest within Formula I consists of compounds and pharmaceutically-acceptable derivatives thereof as follows:

N-(4-Chlorophenyl)[2-(6-quinolylamino)(3-pyridyl)]carboxamide;

- 62 -

- N-(4-Chlorophenyl) [2-(5-isoquinolylamino) (3-pyridyl)]carboxamide;
- N-(4-Chlorophenyl) [2-(1H-indazol-5-ylamino) (3-pyridyl)]carboxamide;
- 5 N-(4-Chlorophenyl) [2-(1H-indazol-6-ylamino) (3-pyridyl)]carboxamide;
- 2-(1H-Indazol-6-ylamino)-N-(4-isopropyl-phenyl)nicotinamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)]-N-[3-(methylethyl)phenyl]carboxamide;
- 10 [2-(1H-Indazol-6-ylamino) (3-pyridyl)]-N-[4-(methylpropyl)phenyl]carboxamide;
- N-[4-(tert-Butyl)phenyl] [2-(1H-indazol-6-ylamino) (3-pyridyl)]carboxamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)]-N-[3-(trifluoromethyl)phenyl]carboxamide;
- 15 N-[3-(tert-Butyl)phenyl] [2-(1H-indazol-6-ylamino) (3-pyridyl)]carboxamide;
- 2-(1H-Indazol-6-ylamino)-N-(4-{2,2,2-trifluoro-1-[2-(2-methoxy-ethoxy)-ethoxy]-1-trifluoromethyl-ethyl}-phenyl)-
- 20 nicotinamide trifluoroacetate;
- [2-(Benzotriazol-6-ylamino) (3-pyridyl)]-N-[4-(tert-butyl)phenyl]carboxamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)]-N-(3-phenylpyrazol-5-yl)carboxamide;
- 25 N-(4-Chloro-3-sulfamoylphenyl) [2-(1H-indazol-6-ylamino) (3-pyridyl)]carboxamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)]-N-(4-methyl-2-oxo-1,2-dihydroquinol-7-yl)carboxamide;
- N-[4-(Methylethyl)phenyl] {2-[(4-methyl-2-oxo(7-hydroquinolyl)) amino] (3-pyridyl)}carboxamide;
- 30 N-[5-(tert-Butyl)isoxazol-3-yl] [2-(1H-indazol-6-ylamino) (3-pyridyl)]carboxamide;
- N-[5-(tert-Butyl)-1-methylpyrazol-3-yl] [2-(1H-indazol-6-ylamino) (3-pyridyl)]carboxamide;

- 63 -

- N-[4-(tert-Butyl)(1,3-thiazol-2-yl)][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
N-(4-Chlorophenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide hydrochloride;
5 N-[5-(tert-Butyl)(1,3,4-thiadiazol-2-yl)][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(1,1,2,2,3,3,4,4,4-nonafluorobutyl)phenyl]carboxamide;
{2-[(1-Methyl(1H-indazol-6-yl))amino](3-pyridyl)}-N-[4-(methylethyl)phenyl]carboxamide;
10 N-[4-(tert-Butyl)phenyl]{2-[(7-bromo(1H-indazol-6-yl))amino](3-pyridyl)}carboxamide;
2-(1H-Indazol-6-ylamino)-N-[4-tert-butyl-3-(1,2,3,6-tetrahydropyridin-4-yl)phenyl]nicotinamide;
15 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]carboxamide;
N-[5-(tert-Butyl)-2-methoxyphenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
N-[4-(tert-Butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide hydrochloride;
20 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[6-[4-(trifluoromethyl)piperidyl](3-pyridyl)]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(1-oxo(7-2,3,4-trihydroisoquinolyl))carboxamide;
25 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylethoxy)phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]carboxamide;
N-(4-{(1S)-1-[(Methylethyl)amino]ethyl}phenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
30 N-[4-(tert-Butyl)-3-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(4-methylpiperazinyl)phenyl]carboxamide;

- 64 -

- N-[4-(tert-Butyl)-2-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- N-[2-[2-(Dimethylamino)ethoxy]-5-(tert-butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 5 N-[3-[2-(Dimethylamino)ethoxy]phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 2-(1H-Indazol-6-ylamino)-N-[3-(2-pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenyl]-nicotinamide hydrochloride;
- N-(3-Hydroxy-4-methoxyphenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 10 N-[3-[2-(Dimethylamino)ethoxy]-4-methoxyphenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-methoxy-3-(1-methyl(4-piperidyl)oxy)phenyl]carboxamide;
- 15 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolin-2-yl)carboxamide;
- [2-({3-[2-(Dimethylamino)ethoxy](1H-indazol-6-yl)}amino)(3-pyridyl)]-N-[4-(tert-butyl)phenyl]carboxamide;
- N-[3,3-Dimethyl-1-(4-piperidylmethyl)indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 20 2-(1H-Indazol-6-ylamino)-N-(3-morpholin-4-ylmethyl-4-pentafluoroethyl-phenyl)-nicotinamide hydrochloride;
- N-[3,3-Dimethyl-1-(1-methyl-piperidin-4-ylmethyl)-2,3-dihydro-1H-indol-6-yl]-2-(1H-indazol-6-ylamino)-
- 25 nicotinamide;
- 2-(1H-Indazol-6-ylamino)-N-(4-phenoxy-phenyl)-nicotinamide;
- [2-(1H-Indazol-5-ylamino)(3-pyridyl)]-N-(4-phenoxyphenyl)carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-phenylphenyl)carboxamide;
- 30 [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylsulfonyl)phenyl]carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(1-methyl(4-piperidyl))indolin-6-yl]carboxamide;

- 65 -

- N-[3,3-Dimethyl-1-(1-methyl(4-piperidyl))indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(1-methyl(4-piperidyl))indol-5-yl]carboxamide;
5 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(trifluoromethyl)phenyl]carboxamide;
N-[3-[3-(Dimethylamino)propyl]-5-(trifluoromethyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
2-(1H-Indazol-6-ylamino)-N-(2-oxo-4-trifluoromethyl-2H-chromen-7-yl)-nicotinamide hydrochloride;
10 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[5-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-3-(trifluoromethyl)phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(1-methyl(4-piperidyl))phenyl]carboxamide;
15 N-[4-(tert-Butyl)-3-(3-piperidylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
N-[3-((1E)-4-Pyrrolidinylbut-1-enyl)-4-(tert-butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
20 N-[4-(tert-Butyl)-3-(3-pyrrolidinylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
N-[4-(tert-Butyl)-3-(3-morpholin-4-ylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
25 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-[3-(4-methylpiperazinyl)-3-oxopropyl]phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-[3-(4-methylpiperazinyl)-3-oxopropyl]phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-[3-(4-methylpiperazinyl)propyl]phenyl]carboxamide;
30 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-[3-(4-methylpiperazinyl)propyl]phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(2-morpholin-4-ylethyl)indol-6-yl]carboxamide;

- 66 -

N-[4-(1,1-Dimethyl-3-morpholin-4-ylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
2-(1H-Indazol-6-ylamino)-N-(4-(2,2,2-trifluoro-1-[2-(2-methoxy-ethoxy)-ethoxy]-1-trifluoromethyl-ethyl)-phenyl)-
5 nicotinamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-[2,2,2-trifluoro-1-(2-piperidylethoxy)-1-(trifluoromethyl)ethyl]phenyl)carboxamide;
N-[4-(*tert*-Butyl)phenyl][6-fluoro-2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
10 [6-Fluoro-2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylethyl)phenyl]carboxamide;
[6-Fluoro-2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[3-(trifluoromethyl)phenyl]carboxamide; and
15 {2-[(1-(2-Pyridyl)pyrrolidin-3-yl)amino](3-pyridyl)}-N-[3-(trifluoromethyl)phenyl]carboxamide.

Indications

Compounds of the present invention would be useful
20 for, but not limited to, the prevention or treatment of angiogenesis related diseases. The compounds of the invention have kinase inhibitory activity, such as VEGFR/KDR inhibitory activity. The compounds of the invention are useful in therapy as antineoplasia agents or to minimize
25 deleterious effects of VEGF.

Compounds of the invention would be useful for the treatment of neoplasia including cancer and metastasis, including, but not limited to: carcinoma such as cancer of the bladder, breast, colon, kidney, liver, lung (including
30 small cell lung cancer), esophagus, gall-bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin (including squamous cell carcinoma); hematopoietic tumors of lymphoid lineage (including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-
35 cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma,

- 67 -

hairy cell lymphoma and Burkett's lymphoma); hematopoietic tumors of myeloid lineage (including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia); tumors of mesenchymal origin
5 (including fibrosarcoma and rhabdomyosarcoma, and other sarcomas, e.g. soft tissue and bone); tumors of the central and peripheral nervous system (including astrocytoma, neuroblastoma, glioma and schwannomas); and other tumors (including melanoma, seminoma, teratocarcinoma,
10 osteosarcoma, xenoderma pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma).

Preferably, the compounds are useful for the treatment of neoplasia selected from lung cancer, colon cancer and breast cancer.

15 The compounds also would be useful for treatment of ophthalmological conditions such as corneal graft rejection, ocular neovascularization, retinal neovascularization including neovascularization following injury or infection, diabetic retinopathy, retrolental fibroplasia and
20 neovascular glaucoma; retinal ischemia; vitreous hemorrhage; ulcerative diseases such as gastric ulcer; pathological, but non-malignant, conditions such as hemangiomas, including infantile hemangiomas, angiofibroma of the nasopharynx and avascular necrosis of bone; and disorders of the female
25 reproductive system such as endometriosis. The compounds are also useful for the treatment of edema, and conditions of vascular hyperpermeability.

The compounds of the invention are useful in therapy of proliferative diseases. These compounds can be used for
30 the treatment of an inflammatory rheumatoid or rheumatic disease, especially of manifestations at the locomotor apparatus, such as various inflammatory rheumatoid diseases, especially chronic polyarthritis including rheumatoid arthritis, juvenile arthritis or psoriasis arthropathy;

- 68 -

paraneoplastic syndrome or tumor-induced inflammatory diseases, turbid effusions, collagenosis, such as systemic Lupus erythematosus, poly-myositis, dermatomyositis, systemic scleroderma or mixed collagenosis; postinfectious
5 arthritis (where no living pathogenic organism can be found at or in the affected part of the body), seronegative spondylarthritis, such as spondylitis ankylosans; vasculitis, sarcoidosis, or arthrosis; or further any combinations thereof. An example of an inflammation related
10 disorder is (a) synovial inflammation, for example, synovitis, including any of the particular forms of synovitis, in particular bursal synovitis and purulent synovitis, as far as it is not crystal-induced. Such synovial inflammation may for example, be consequential to
15 or associated with disease, e.g. arthritis, e.g. osteoarthritis, rheumatoid arthritis or arthritis deformans. The present invention is further applicable to the systemic treatment of inflammation, e.g. inflammatory diseases or conditions, of the joints or locomotor apparatus in the
20 region of the tendon insertions and tendon sheaths. Such inflammation may be, for example, be consequential to or associated with disease or further (in a broader sense of the invention) with surgical intervention, including, in particular conditions such as insertion endopathy,
25 myofasciale syndrome and tendomyositis. The present invention is further especially applicable to the treatment of inflammation, e.g. inflammatory disease or condition, of connective tissues including dermatomyositis and myositis.

These compounds can be used as active agents against
30 such disease states as arthritis, atherosclerosis, psoriasis, hemangiomas, myocardial angiogenesis, coronary and cerebral collaterals, ischemic limb angiogenesis, wound healing, peptic ulcer Helicobacter related diseases, fractures, cat scratch fever, rubeosis, neovascular glaucoma

- 69 -

and retinopathies such as those associated with diabetic retinopathy or macular degeneration. In addition, some of these compounds can be used as active agents against solid tumors, malignant ascites, hematopoietic cancers and
5 hyperproliferative disorders such as thyroid hyperplasia (especially Grave's disease), and cysts (such as hypervascularity of ovarian stroma, characteristic of polycystic ovarian syndrome (Stein- Leventhal syndrome)) since such diseases require a proliferation of blood vessel
10 cells for growth and/or metastasis.

Further, some of these compounds can be used as active agents against burns, chronic lung disease, stroke, polyps, anaphylaxis, chronic and allergic inflammation, ovarian hyperstimulation syndrome, brain tumor-associated cerebral
15 edema, high-altitude, trauma or hypoxia induced cerebral or pulmonary edema, ocular and macular edema, ascites, and other diseases where vascular hyperpermeability, effusions, exudates, protein extravasation, or edema is a manifestation of the disease. The compounds will also be useful in
20 treating disorders in which protein extravasation leads to the deposition of fibrin and extracellular matrix, promoting stromal proliferation (e.g. fibrosis, cirrhosis and carpal tunnel syndrome).

The compounds of the present invention are also useful
25 in the treatment of ulcers including bacterial, fungal, Mooren ulcers and ulcerative colitis.

The compounds of the present invention are also useful in the treatment of conditions wherein undesired angiogenesis, edema, or stromal deposition occurs in viral
30 infections such as Herpes simplex, Herpes Zoster, AIDS, Kaposi's sarcoma, protozoan infections and toxoplasmosis, following trauma, radiation, stroke, endometriosis, ovarian hyperstimulation syndrome, systemic lupus, sarcoidosis, synovitis, Crohn's disease, sickle cell anaemia, Lyme

- 70 -

disease, pemphigoid, Paget's disease, hyperviscosity syndrome, Osler-Weber-Rendu disease, chronic inflammation, chronic occlusive pulmonary disease, asthma, and inflammatory rheumatoid or rheumatic disease. The compounds
5 are also useful in the reduction of sub-cutaneous fat and for the treatment of obesity.

The compounds of the present invention are also useful in the treatment of ocular conditions such as ocular and macular edema, ocular neovascular disease, scleritis, radial
10 keratotomy, uveitis, vitritis, myopia, optic pits, chronic retinal detachment, post-laser complications, glaucoma, conjunctivitis, Stargardt's disease and Eales disease in addition to retinopathy and macular degeneration.

The compounds of the present invention are also useful
15 in the treatment of cardiovascular conditions such as atherosclerosis, restenosis, arteriosclerosis, vascular occlusion and carotid obstructive disease.

The compounds of the present invention are also useful in the treatment of cancer related indications such as solid
20 tumors, sarcomas (especially Ewing's sarcoma and osteosarcoma), retinoblastoma, rhabdomyosarcomas, neuroblastoma, hematopoietic malignancies, including leukemia and lymphoma, tumor-induced pleural or pericardial effusions, and malignant ascites.

25 The compounds of the present invention are also useful in the treatment of diabetic conditions such as diabetic retinopathy and microangiopathy.

The compounds of this invention may also act as inhibitors of other protein kinases, e.g. p38, EGFR, CDK-2,
30 CDK-5, IKK, JNK3, and thus be effective in the treatment of diseases associated with other protein kinases.

Besides being useful for human treatment, these compounds are also useful for veterinary treatment of companion animals, exotic animals and farm animals,

- 71 -

including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.

As used herein, the compounds of the present invention include the pharmaceutically acceptable derivatives
5 thereof.

Definitions

The term "treatment" includes therapeutic treatment as well as prophylactic treatment (either preventing the onset
10 of disorders altogether or delaying the onset of a preclinically evident stage of disorders in individuals).

A "pharmaceutically-acceptable derivative " denotes any salt, ester of a compound of this invention, or any other compound which upon administration to a patient is
15 capable of providing (directly or indirectly) a compound of this invention, or a metabolite or residue thereof, characterized by the ability to inhibit angiogenesis.

The phrase "therapeutically-effective" is intended to qualify the amount of each agent, which will achieve the
20 goal of improvement in disorder severity and the frequency of incidence over treatment of each agent by itself, while avoiding adverse side effects typically associated with alternative therapies. For example, effective neoplastic therapeutic agents prolong the survivability of the patient,
25 inhibit the rapidly-proliferating cell growth associated with the neoplasm, or effect a regression of the neoplasm.

The term "H" denotes a single hydrogen atom. This radical may be attached, for example, to an oxygen atom to form a hydroxyl radical.

30 Where the term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "alkylamino", it embraces linear or branched radicals having one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about six carbon atoms.

- 72 -

Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, *sec*-butyl, *tert*-butyl, pentyl, isoamyl, hexyl and the like. Even more preferred are lower alkyl radicals having one or two carbon atoms. The term

5 "alkylenyl" embraces bridging divalent alkyl radicals such as methylenyl and ethylenyl. The term "lower alkyl substituted with R²" does not include an acetal moiety.

The term "alkenyl" embraces linear or branched radicals having at least one carbon-carbon double bond of

10 two to about twelve carbon atoms. More preferred alkenyl radicals are "lower alkenyl" radicals having two to about six carbon atoms. Most preferred lower alkenyl radicals are radicals having two to about four carbon atoms. Examples of alkenyl radicals include ethenyl, propenyl, allyl, propenyl,

15 butenyl and 4-methylbutenyl. The terms "alkenyl" and "lower alkenyl", embrace radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations.

The term "alkynyl" denotes linear or branched radicals having at least one carbon-carbon triple bond and having two

20 to about twelve carbon atoms. More preferred alkynyl radicals are "lower alkynyl" radicals having two to about six carbon atoms. Most preferred are lower alkynyl radicals having two to about four carbon atoms. Examples of such radicals include propargyl, butynyl, and the like.

25 The term "halo" means halogens such as fluorine, chlorine, bromine or iodine atoms.

The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl,

30 dihaloalkyl and polyhaloalkyl radicals including perhaloalkyl. A monohaloalkyl radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different

- 73 -

halo radicals. "Lower haloalkyl" embraces radicals having 1-6 carbon atoms. Even more preferred are lower haloalkyl radicals having one to three carbon atoms. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, 5 trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. "Perfluoroalkyl" means alkyl radicals having all hydrogen 10 atoms replaced with fluoro atoms. Examples include trifluoromethyl and pentafluoroethyl.

The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl 15 radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms and one or more hydroxyl radicals. Examples of such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl. Even more preferred are lower 20 hydroxyalkyl radicals having one to three carbon atoms.

The term "alkoxy" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals are "lower alkoxy" radicals having one to six carbon atoms. 25 Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and *tert*-butoxy. Even more preferred are lower alkoxy radicals having one to three carbon atoms. Alkoxy radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkoxy" 30 radicals. Even more preferred are lower haloalkoxy radicals having one to three carbon atoms. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy.

- 74 -

The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one or two rings wherein such rings may be attached together in a fused manner. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, indenyl, tetrahydronaphthyl, and indanyl. More preferred aryl is phenyl. Said "aryl" group may have 1 to 3 substituents such as lower alkyl, hydroxyl, halo, haloalkyl, nitro, cyano, alkoxy and lower alkylamino. Phenyl substituted with -O-CH₂-O- forms the aryl benzodioxolyl substituent.

The term "heterocyclyl" embraces saturated, partially saturated and unsaturated heteroatom-containing ring radicals, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. It does not include rings containing -O-O-, -O-S- or -S-S- portions. Said "heterocyclyl" group may have 1 to 3 substituents such as hydroxyl, Boc, halo, haloalkyl, cyano, lower alkyl, lower aralkyl, oxo, lower alkoxy, amino and lower alkylamino.

Examples of saturated heterocyclic radicals include saturated 3 to 6-membered heteromonocyclic groups containing 1 to 4 nitrogen atoms [e.g. pyrrolidinyl, imidazolidinyl, piperidinyl, pyrrolinyl, piperazinyl]; saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. morpholinyl]; saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., thiazolidinyl]. Examples of partially saturated heterocyclyl radicals include dihydrothienyl, dihydropyranyl, dihydrofuryl and dihydrothiazolyl.

Examples of unsaturated heterocyclic radicals, also termed "heteroaryl" radicals, include unsaturated 5 to 6 membered heteromonocyclyl group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, imidazolyl, pyrazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrimidyl, pyrazinyl,

- 75 -

pyridazinyl, triazolyl [e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl]; unsaturated 5- to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, 2-furyl, 3-furyl, etc.; unsaturated 5 to 6-membered heteromonocyclic group containing a sulfur atom, for example, 2-thienyl, 3-thienyl, etc.; unsaturated 5- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl [e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl]; unsaturated 5 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl [e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl].

The term also embraces radicals where heterocyclic radicals are fused/condensed with aryl radicals: unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indoliziny, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl [e.g., tetrazolo [1,5-b]pyridazinyl]; unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. benzoxazolyl, benzoxadiazolyl]; unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., benzothiazolyl, benzothiadiazolyl]; and saturated, partially unsaturated and unsaturated condensed heterocyclic group containing 1 to 2 oxygen or sulfur atoms [e.g. benzofuryl, benzothieryl, 2,3-dihydro-benzo[1,4]dioxinyl and dihydrobenzofuryl]. Preferred heterocyclic radicals include five to ten membered fused or unfused radicals. More preferred examples of heteroaryl radicals include quinolyl, isoquinolyl, imidazolyl, pyridyl, thienyl, thiazolyl, oxazolyl, furyl, and pyrazinyl. Other preferred heteroaryl

- 76 -

radicals are 5- or 6-membered heteroaryl, containing one or two heteroatoms selected from sulfur, nitrogen and oxygen, selected from thienyl, furyl, pyrrolyl, indazolyl, pyrazolyl, oxazolyl, triazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, piperidiny and pyrazinyl.

Particular examples of non-nitrogen containing heteroaryl include pyranyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, benzofuryl, benzothienyl, and the like.

Particular examples of partially saturated and saturated heterocyclyl include pyrrolidinyl, imidazolidinyl, piperidiny, pyrrolinyl, pyrazolidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, thiazolidinyl, dihydrothienyl, 2,3-dihydro-benzo[1,4]dioxanyl, indolinyl, isoindolinyl, dihydrobenzothienyl, dihydrobenzofuryl, isochromanyl, chromanyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, 1,2,3,4-tetrahydro-quinolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, 3,4-dihydro-2H-benzo[1,4]oxazinyl, benzo[1,4]dioxanyl, 2,3-dihydro-1H-1λ'-benzo[d]isothiazol-6-yl, dihydropyranyl, dihydrofuryl and dihydrothiazolyl, and the like.

The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals -SO₂-.

The terms "sulfamyl," "aminosulfonyl" and "sulfonamidyl," denotes a sulfonyl radical substituted with an amine radical, forming a sulfonamide (-SO₂NH₂).

The term "alkylaminosulfonyl" includes "N-alkylaminosulfonyl" where sulfamyl radicals are independently substituted with one or two alkyl radical(s). More preferred alkylaminosulfonyl radicals are "lower alkylaminosulfonyl" radicals having one to six carbon atoms. Even more preferred are lower alkylaminosulfonyl radicals

- 77 -

having one to three carbon atoms. Examples of such lower alkylaminosulfonyl radicals include N-methylaminosulfonyl, and N-ethylaminosulfonyl.

The terms "carboxy" or "carboxyl", whether used alone
5 or with other terms, such as "carboxyalkyl", denotes $\text{-CO}_2\text{H}$.

The term "carbonyl", whether used alone or with other terms, such as "aminocarbonyl", denotes -(C=O)- .

The term "aminocarbonyl" denotes an amide group of the formula -C(=O)NH_2 .

10 The terms "N-alkylaminocarbonyl" and "N,N-dialkylaminocarbonyl" denote aminocarbonyl radicals independently substituted with one or two alkyl radicals, respectively. More preferred are "lower alkylaminocarbonyl" having lower alkyl radicals as described above attached to
15 an aminocarbonyl radical.

The terms "N-arylaminocarbonyl" and "N-alkyl-N-arylaminocarbonyl" denote aminocarbonyl radicals substituted, respectively, with one aryl radical, or one alkyl and one aryl radical.

20 The term "heterocyclalkylenyl" embraces heterocyclic-substituted alkyl radicals. More preferred heterocyclalkylenyl radicals are "5- or 6-membered heteroarylalkylenyl" radicals having alkyl portions of one to six carbon atoms and a 5- or 6-membered heteroaryl
25 radical. Even more preferred are lower heteroarylalkylenyl radicals having alkyl portions of one to three carbon atoms. Examples include such radicals as pyridylmethyl and thienylmethyl.

The term "aralkyl" embraces aryl-substituted alkyl
30 radicals. Preferable aralkyl radicals are "lower aralkyl" radicals having aryl radicals attached to alkyl radicals having one to six carbon atoms. Even more preferred are "phenylalkylenyl" attached to alkyl portions having one to three carbon atoms. Examples of such radicals include

- 78 -

benzyl, diphenylmethyl and phenylethyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy.

5 The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent sulfur atom. Even more preferred are lower alkylthio radicals having one to three carbon atoms. An example of "alkylthio" is methylthio, ($\text{CH}_3\text{S}-$).

10 The term "haloalkylthio" embraces radicals containing a haloalkyl radical, of one to ten carbon atoms, attached to a divalent sulfur atom. Even more preferred are lower haloalkylthio radicals having one to three carbon atoms. An example of "haloalkylthio" is trifluoromethylthio.

15 The term "alkylamino" embraces "N-alkylamino" and "N,N-dialkylamino" where amino groups are independently substituted with one alkyl radical and with two alkyl radicals, respectively. More preferred alkylamino radicals are "lower alkylamino" radicals having one or two alkyl radicals of one to six carbon atoms, attached to a nitrogen atom. Even more preferred are lower alkylamino radicals having one to three carbon atoms. Suitable alkylamino radicals may be mono or dialkylamino such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino and the
20 like.
25

The term "arylamino" denotes amino groups which have been substituted with one or two aryl radicals, such as N-phenylamino. The arylamino radicals may be further substituted on the aryl ring portion of the radical.

30 The term "heteroarylamino" denotes amino groups which have been substituted with one or two heteroaryl radicals, such as N-thienylamino. The "heteroarylamino" radicals may be further substituted on the heteroaryl ring portion of the radical.

- 79 -

The term "aralkylamino" denotes amino groups which have been substituted with one or two aralkyl radicals. More preferred are phenyl-C₁-C₃-alkylamino radicals, such as N-benzylamino. The aralkylamino radicals may be further substituted on the aryl ring portion.

The terms "N-alkyl-N-aryl-amino" and "N-aralkyl-N-alkylamino" denote amino groups which have been independently substituted with one aralkyl and one alkyl radical, or one aryl and one alkyl radical, respectively, to an amino group.

The term "aminoalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more amino radicals. More preferred aminoalkyl radicals are "lower aminoalkyl" radicals having one to six carbon atoms and one or more amino radicals. Examples of such radicals include aminomethyl, aminoethyl, aminopropyl, aminobutyl and aminohexyl. Even more preferred are lower aminoalkyl radicals having one to three carbon atoms.

The term "alkylaminoalkyl" embraces alkyl radicals substituted with alkylamino radicals. More preferred alkylaminoalkyl radicals are "lower alkylaminoalkyl" radicals having alkyl radicals of one to six carbon atoms. Even more preferred are lower alkylaminoalkyl radicals having alkyl radicals of one to three carbon atoms. Suitable alkylaminoalkyl radicals may be mono or dialkyl substituted, such as N-methylaminomethyl, N,N-dimethyl-aminoethyl, N,N-diethylaminomethyl and the like.

The term "alkylaminoalkoxy" embraces alkoxy radicals substituted with alkylamino radicals. More preferred alkylaminoalkoxy radicals are "lower alkylaminoalkoxy" radicals having alkoxy radicals of one to six carbon atoms. Even more preferred are lower alkylaminoalkoxy radicals having alkyl radicals of one to three carbon atoms. Suitable

- 80 -

alkylaminoalkoxy radicals may be mono or dialkyl substituted, such as N-methylaminoethoxy, N,N-dimethylaminoethoxy, N,N-diethylaminoethoxy and the like.

The term "alkylaminoalkoxyalkoxy" embraces alkoxy radicals substituted with alkylaminoalkoxy radicals. More preferred alkylaminoalkoxyalkoxy radicals are "lower alkylaminoalkoxyalkoxy" radicals having alkoxy radicals of one to six carbon atoms. Even more preferred are lower alkylaminoalkoxyalkoxy radicals having alkyl radicals of one to three carbon atoms. Suitable alkylaminoalkoxyalkoxy radicals may be mono or dialkyl substituted, such as N-methylaminomethoxyethoxy, N-methylaminoethoxyethoxy, N,N-dimethylaminoethoxyethoxy, N,N-diethylaminomethoxymethoxy and the like.

The term "carboxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more carboxy radicals. More preferred carboxyalkyl radicals are "lower carboxyalkyl" radicals having one to six carbon atoms and one carboxy radical. Examples of such radicals include carboxymethyl, carboxypropyl, and the like. Even more preferred are lower carboxyalkyl radicals having one to three CH₂ groups.

The term "halosulfonyl" embraces sulfonyl radicals substituted with a halogen radical. Examples of such halosulfonyl radicals include chlorosulfonyl and fluorosulfonyl.

The term "arylthio" embraces aryl radicals of six to ten carbon atoms, attached to a divalent sulfur atom. An example of "arylthio" is phenylthio.

The term "aralkylthio" embraces aralkyl radicals as described above, attached to a divalent sulfur atom. More preferred are phenyl-C₁-C₃-alkylthio radicals. An example of "aralkylthio" is benzylthio.

- 81 -

The term "aryloxy" embraces optionally substituted aryl radicals, as defined above, attached to an oxygen atom. Examples of such radicals include phenoxy.

5 The term "aralkoxy" embraces oxy-containing aralkyl radicals attached through an oxygen atom to other radicals. More preferred aralkoxy radicals are "lower aralkoxy" radicals having optionally substituted phenyl radicals attached to lower alkoxy radical as described above.

10 The term "heteroaryloxy" embraces optionally substituted heteroaryl radicals, as defined above, attached to an oxygen atom.

The term "heteroarylalkoxy" embraces oxy-containing heteroarylalkyl radicals attached through an oxygen atom to other radicals. More preferred heteroarylalkoxy radicals are 15 "lower heteroarylalkoxy" radicals having optionally substituted heteroaryl radicals attached to lower alkoxy radical as described above.

The term "cycloalkyl" includes saturated carbocyclic groups. Preferred cycloalkyl groups include C₃-C₆ rings. 20 More preferred compounds include, cyclopentyl, cyclopropyl, and cyclohexyl.

The term "cycloalkenyl" includes carbocyclic groups having one or more carbon-carbon double bonds including "cycloalkyldienyl" compounds. Preferred cycloalkenyl groups 25 include C₃-C₆ rings. More preferred compounds include, for example, cyclopentenyl, cyclopentadienyl, cyclohexenyl and cycloheptadienyl.

The term "comprising" is meant to be open ended, including the indicated component but not excluding other 30 elements.

The term "Formulas I-X" includes any sub formulas.

The compounds of the invention are endowed with kinase inhibitory activity, such as KDR inhibitory activity.


The present invention also comprises the use of a 35 compound of the invention, or pharmaceutically acceptable

- 82 -

salt thereof, in the manufacture of a medicament for the treatment either acutely or chronically of an angiogenesis mediated disease state, including those described previously. The compounds of the present invention are
 5 useful in the manufacture of an anti-cancer medicament. The compounds of the present invention are also useful in the manufacture of a medicament to attenuate or prevent disorders through inhibition of KDR.

The present invention comprises a pharmaceutical
 10 composition comprising a therapeutically-effective amount of a compound of Formulas I-X in association with a least one pharmaceutically-acceptable carrier, adjuvant or diluent.

The present invention also comprises a method of
 treating angiogenesis related disorders in a subject having
 15 or susceptible to such disorder, the method comprising treating the subject with a therapeutically-effective amount of a compound of Formula I



20


wherein each of A¹ and A² is independently C, CH or N;

wherein ring A is selected from

- a) 5- or 6-membered partially saturated heterocyclyl,
- b) 5- or 6-membered heteroaryl,
- 25 c) 9-, 10- or 11-membered fused partially saturated heterocyclyl,
- d) 9- or 10-membered fused heteroaryl,
- e) aryl, and

- 83 -

f) 4-, 5- or 6-membered cycloalkenyl;



wherein X is

wherein Z is oxygen or sulfur;

wherein R is selected from

- 5 a) substituted or unsubstituted 4-6 membered heterocyclyl,
- b) substituted aryl, and
- c) substituted or unsubstituted fused 9-14-membered bicyclic or tricyclic heterocyclyl;
- 10 wherein substituted R is substituted with one or more substituents independently selected from halo, $-OR^3$, $-SR^3$, $-SO_2R^3$, $-CO_2R^3$, $-CONR^3R^3$, $-COR^3$, $-NR^3R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$, cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally
- 15 substituted phenyl, nitro, alkylaminoalkoxyalkoxy, cyano, alkylaminoalkoxy, lower alkyl substituted with R^2 , lower alkenyl substituted with R^2 , and lower alkynyl substituted with R^2 ;
- wherein R^1 is selected from
- 20 a) substituted or unsubstituted 6-10 membered aryl,
- b) substituted or unsubstituted 4-6 membered heterocyclyl,
- c) substituted or unsubstituted 9-14 membered bicyclic or tricyclic heterocyclyl,
- 25 d) cycloalkyl, and
- e) cycloalkenyl,
- wherein substituted R^1 is substituted with one or more substituents independently selected from halo, $-OR^3$, $-SR^3$, $-CO_2R^3$, $-CONR^3R^3$, $-COR^3$, $-NR^3R^3$, $-NH(C_1-C_4$
- 30 alkylenyl $R^{14})$, $-SO_2R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$, optionally substituted cycloalkyl,

- 84 -

- optionally substituted 4-6 membered heterocyclyl,
optionally substituted phenyl, halosulfonyl, cyano,
alkylaminoalkoxy, alkylaminoalkoxyalkoxy, nitro,
lower alkyl substituted with R², lower alkenyl
5 substituted with R², and lower alkynyl substituted
with R²;
- wherein R² is one or more substituents independently selected
from H, halo, -OR³, oxo, -SR³, -CO₂R³, -COR³, -CONR³R³, -
NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, cycloalkyl,
10 optionally substituted phenylalkylenyl, optionally
substituted 4-6 membered heterocyclyl, optionally
substituted heteroarylalkylenyl, optionally substituted
phenyl, lower alkyl, cyano, lower hydroxyalkyl, lower
carboxyalkyl, nitro, lower alkenyl, lower alkynyl, lower
15 aminoalkyl, lower alkylaminoalkyl and lower haloalkyl;
wherein R³ is selected from H, lower alkyl, optionally
substituted phenyl, optionally substituted 4-6 membered
heterocyclyl, optionally substituted C₃-C₆-cycloalkyl,
optionally substituted phenylalkyl, optionally
20 substituted 4-6 membered heterocyclylalkyl, optionally
substituted C₃-C₆ cycloalkylalkyl, and lower haloalkyl;
wherein R⁴ is selected from a direct bond, C₂₋₄-alkylenyl, C₂₋₄-
alkenylenyl and C₂₋₄-alkynylenyl, where one of the CH₂
groups may be substituted with an oxygen atom or an -NH-,
25 wherein R⁴ is optionally substituted with hydroxy;
wherein R⁵ is selected from H, lower alkyl, optionally
substituted phenyl and lower aralkyl;
wherein R¹⁴ is selected from H, phenyl, 4-6 membered
heterocyclyl and C₃-C₆ cycloalkyl;
30 and pharmaceutically acceptable derivatives thereof;
provided R is not unsubstituted 2-thienyl, 2-pyridyl or 3-
pyridyl.

- 85 -

COMBINATIONS

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more compounds of the invention or other agents. When administered as a combination, the therapeutic agents can be formulated as separate compositions that are administered at the same time or sequentially at different times, or the therapeutic agents can be given as a single composition.

The phrase "co-therapy" (or "combination-therapy"), in defining use of a compound of the present invention and another pharmaceutical agent, is intended to embrace administration of each agent in a sequential manner in a regimen that will provide beneficial effects of the drug combination, and is intended as well to embrace co-administration of these agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of these active agents or in multiple, separate capsules for each agent.

Specifically, the administration of compounds of the present invention may be in conjunction with additional therapies known to those skilled in the art in the prevention or treatment of neoplasia, such as with radiation therapy or with cytostatic or cytotoxic agents.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the accepted dosage ranges. Compounds of Formula I may also be administered sequentially with known anticancer or cytotoxic agents when a combination formulation is inappropriate. The invention is not limited in the sequence of administration; compounds of the invention may be administered either prior to, simultaneous with or after administration of the known anticancer or cytotoxic agent.

- 86 -

Currently, standard treatment of primary tumors consists of surgical excision followed by either radiation or IV administered chemotherapy. The typical chemotherapy regime consists of either DNA alkylating agents, DNA
5 intercalating agents, CDK inhibitors, or microtubule poisons. The chemotherapy doses used are just below the maximal tolerated dose and therefore dose limiting toxicities typically include, nausea, vomiting, diarrhea, hair loss, neutropenia and the like.

10 There are large numbers of antineoplastic agents available in commercial use, in clinical evaluation and in pre-clinical development, which would be selected for treatment of neoplasia by combination drug chemotherapy. Such antineoplastic agents fall into several major
15 categories, namely, antibiotic-type agents, alkylating agents, antimetabolite agents, hormonal agents, immunological agents, interferon-type agents and a category of miscellaneous agents.

A first family of antineoplastic agents which may be
20 used in combination with compounds of the present invention consists of antimetabolite-type/thymidilate synthase inhibitor antineoplastic agents. Suitable antimetabolite antineoplastic agents may be selected from but not limited to the group consisting of 5-FU-fibrinogen, acanthifolic
25 acid, aminothiadiazole, brequinar sodium, carmofur, Ciba-Geigy CGP-30694, cyclopentyl cytosine, cytarabine phosphate stearate, cytarabine conjugates, Lilly DATHF, Merrel Dow DDFC, dezaguanine, dideoxycytidine, dideoxyguanosine, didox, Yoshitomi DMDC, doxifluridine, Wellcome EHNA, Merck & Co.
30 EX-015, fazarabine, floxuridine, fludarabine phosphate, 5-fluorouracil, N-(2'-furanidyl)-5-fluorouracil, Daiichi Seiyaku FO-152, isopropyl pyrrolizine, Lilly LY-188011, Lilly LY-264618, methobenzaprim, methotrexate, Wellcome MZPES, norspermidine, NCI NSC-127716, NCI NSC-264880, NCI

- 87 -

NSC-39661, NCI NSC-612567, Warner-Lambert PALA, pentostatin, piritrexim, plicamycin, Asahi Chemical PL-AC, Takeda TAC-788, thioguanine, tiazofurin, Erbamont TIF, trimetrexate, tyrosine kinase inhibitors, Taiho UFT and uricytin.

- 5 A second family of antineoplastic agents which may be used in combination with compounds of the present invention consists of alkylating-type antineoplastic agents. Suitable alkylating-type antineoplastic agents may be selected from but not limited to the group consisting of Shionogi 254-S,
- 10 aldo-phosphamide analogues, altretamine, anaxirone, Boehringer Mannheim BBR-2207, bestrabucil, budotitane, Wakunaga CA-102, carboplatin, carmustine, Chinoin-139, Chinoin-153, chlorambucil, cisplatin, cyclophosphamide, American Cyanamid CL-286558, Sanofi CY-233, cyplatate,
- 15 Degussa D-19-384, Sumimoto DACHP(Myrr)2, diphenylspiromustine, diplatinum cytostatic, Erba distamycin derivatives, Chugai DWA-2114R, ITI E09, elmustine, Erbamont FCE-24517, estramustine phosphate sodium, fotemustine, Unimed G-6-M, Chinoin GYKI-17230, hepsul-fam, ifosfamide,
- 20 iproplatin, lomustine, mafosfamide, mitolactol, Nippon Kayaku NK-121, NCI NSC-264395, NCI NSC-342215, oxaliplatin, Upjohn PCNU, prednimustine, Proter PTT-119, ranimustine, semustine, SmithKline SK&F-101772, Yakult Honsha SN-22, spiromustine, Tanabe Seiyaku TA-077, tauromustine,
- 25 temozolomide, teroxirone, tetraplatin and trimelamol.

- A third family of antineoplastic agents which may be used in combination with compounds of the present invention consists of antibiotic-type antineoplastic agents. Suitable antibiotic-type antineoplastic agents may be selected from
- 30 but not limited to the group consisting of Taiho 4181-A, aclarubicin, actinomycin D, actinoplanone, Erbamont ADR-456, aeroplysinin derivative, Ajinomoto AN-201-II, Ajinomoto AN-3, Nippon Soda anisomycins, anthracycline, azinomycin-A, bisucaberin, Bristol-Myers BL-6859, Bristol-Myers BMY-25067,

- 88 -

Bristol-Myers BMY-25551, Bristol-Myers BMY-26605, Bristol-Myers BMY-27557, Bristol-Myers BMY-28438, bleomycin sulfate, bryostatin-1, Taiho C-1027, calicheomycin, chromoximycin, dactinomycin, daunorubicin, Kyowa Hakko DC-102, Kyowa Hakko DC-79, Kyowa Hakko DC-88A, Kyowa Hakko DC89-A1, Kyowa Hakko DC92-B, ditrisarubicin B, Shionogi DOB-41, doxorubicin, doxorubicin-fibrinogen, elsamicin-A, epirubicin, erbstatin, esorubicin, esperamicin-A1, esperamicin-Alb, Erbamont FCE-21954, Fujisawa FK-973, fostriecin, Fujisawa FR-900482, glidobactin, gregatin-A, grincamycin, herbimycin, idarubicin, illudins, kazusamycin, kesarirhodins, Kyowa Hakko KM-5539, Kirin Brewery KRN-8602, Kyowa Hakko KT-5432, Kyowa Hakko KT-5594, Kyowa Hakko KT-6149, American Cyanamid LL-D49194, Meiji Seika ME 2303, menogaril, mitomycin, mitoxantrone, SmithKline M-TAG, neoenactin, Nippon Kayaku NK-313, Nippon Kayaku NKT-01, SRI International NSC-357704, oxalysine, oxaunomycin, peplomycin, pilatin, pirarubicin, porothramycin, pyrindanycin A, Tobishi RA-I, rapamycin, rhizoxin, rodorubicin, sibanomicin, siwenmycin, Sumitomo SM-5887, Snow Brand SN-706, Snow Brand SN-07, sorangicin-A, sparsomycin, SS Pharmaceutical SS-21020, SS Pharmaceutical SS-7313B, SS Pharmaceutical SS-9816B, steffimycin B, Taiho 4181-2, talisomycin, Takeda TAN-868A, terpentecin, thrazine, tricrozarin A, Upjohn U-73975, Kyowa Hakko UCN-10028A, Fujisawa WF-3405, Yoshitomi Y-25024 and zorubicin.

A fourth family of antineoplastic agents which may be used in combination with compounds of the present invention consists of a miscellaneous family of antineoplastic agents, including tubulin interacting agents, topoisomerase II inhibitors, topoisomerase I inhibitors and hormonal agents, selected from but not limited to the group consisting of α -carotene, α -difluoromethyl-arginine, acitretin, Biotec AD-5, Kyorin AHC-52, alstonine, amonafide, amphetinile, amsacrine, Angiostat, ankinomycin, anti-neoplaston A10,

- 89 -

antineoplaston A2, antineoplaston A3, antineoplaston A5,
antineoplaston AS2-1, Henkel APD, aphidicolin glycinate,
asparaginase, Avarol, baccharin, batracylin, benfluron,
benzotript, Ipsen-Beaufour BIM-23015, bisantrene, Bristol-
5 Myers BMY-40481, Vestar boron-10, bromofosfamide, Wellcome
BW-502, Wellcome BW-773, caracemide, carmethizole
hydrochloride, Ajinomoto CDAF, chlorsulfaquinoxalone, Chemes
CHX-2053, Chemex CHX-100, Warner-Lambert CI-921, Warner-
Lambert CI-937, Warner-Lambert CI-941, Warner-Lambert CI-
10 958, clanfenur, claviridenone, ICN compound 1259, ICN
compound 4711, Contracan, Yakult Honsha CPT-11, crisnatol,
curaderm, cytochalasin B, cytarabine, cytocytin, Merz D-609,
DABIS maleate, dacarbazine, datelliptinium, didemnin-B,
dihaematoporphyrin ether, dihydrolenperone, dinaline,
15 distamycin, Toyo Pharmar DM-341, Toyo Pharmar DM-75, Daiichi
Seiyaku DN-9693, docetaxel elliprabin, elliptinium acetate,
Tsumura EPMTc, the epothilones, ergotamine, etoposide,
etretinate, fenretinide, Fujisawa FR-57704, gallium nitrate,
genkwadaphnin, Chugai GLA-43, Glaxo GR-63178, grifolan NMF-
20 5N, hexadecylphosphocholine, Green Cross HO-221,
homoharringtonine, hydroxyurea, BTG ICRF-187, ilmofosine,
isoglutamine, isotretinoin, Otsuka JI-36, Ramot K-477,
Otsuak K-76COONa, Kureha Chemical K-AM, MECT Corp KI-8110,
American Cyanamid L-623, leukoregulin, lonidamine, Lundbeck
25 LU-23-112, Lilly LY-186641, NCI (US) MAP, marycin, Merrel
Dow MDL-27048, Medco MEDR-340, merbarone, merocyanine
derivatives, methylanilinoacridine, Molecular Genetics MGI-
136, minactivin, mitonafide, mitotidone mopidamol,
motretinide, Zenyaku Kogyo MST-16, N-(retinoyl) amino acids,
30 Nisshin Flour Milling N-021, N-acylated-dehydroalanines,
nafazatrom, Taisho NCU-190, nocodazole derivative,
Normosang, NCI NSC-145813, NCI NSC-361456, NCI NSC-604782,
NCI NSC-95580, ocreotide, Ono ONO-112, oquizanocine, Akzo
Org-10172, paclitaxel, pancratistatin, pazelliptine, Warner-

- 90 -

Lambert PD-111707, Warner-Lambert PD-115934, Warner-Lambert PD-131141, Pierre Fabre PE-1001, ICRT peptide D, piroxantrone, polyhaematoporphyrin, polypreic acid, Efamol porphyrin, probimane, procarbazine, proglumide, Invitron
5 protease nexin I, Tobishi RA-700, razoxane, Sapporo Breweries RBS, restrictin-P, retelliptine, retinoic acid, Rhone-Poulenc RP-49532, Rhone-Poulenc RP-56976, SmithKline SK&F-104864, Sumitomo SM-108, Kuraray SMANCS, SeaPharm SP-10094, spatol, spirocyclopropane derivatives,
10 spirogermanium, Unimed, SS Pharmaceutical SS-554, strypoldinone, Stypoldione, Suntory SUN 0237, Suntory SUN 2071, superoxide dismutase, Toyama T-506, Toyama T-680, taxol, Teijin TEI-0303, teniposide, thaliblastine, Eastman Kodak TJB-29, tocotrienol, topotecan, Topostin, Teijin TT-
15 82, Kyowa Hakko UCN-01, Kyowa Hakko UCN-1028, ukrain, Eastman Kodak USB-006, vinblastine sulfate, vincristine, vindesine, vinestramide, vinorelbine, vintriptyl, vinzolidine, withanolides and Yamanouchi YM-534.

Alternatively, the present compounds may also be used
20 in co-therapies with other anti-neoplastic agents, such as acemannan, aclarubicin, aldesleukin, alemtuzumab, alitretinoin, altretamine, amifostine, aminolevulinic acid, amrubicin, amsacrine, anagrelide, anastrozole, ANCER, aneastim, ARGLABIN, arsenic trioxide, BAM 002 (Novelos),
25 bexarotene, bicalutamide, broxuridine, capecitabine, celmoleukin, cetorelix, cladribine, clotrimazole, cytarabine ocfosfate, DA 3030 (Dong-A), daclizumab, denileukin diftitox, deslorelin, dexrazoxane, dilazep, docetaxel, docosanol, doxercalciferol, doxifluridine,
30 doxorubicin, bromocriptine, carmustine, cytarabine, fluorouracil, HIT diclofenac, interferon alfa, daunorubicin, doxorubicin, tretinoin, edelfosine, edrecolomab, eflornithine, emitefur, epirubicin, epoetin beta, etoposide phosphate, exemestane, exisulind,

- 91 -

fadrozole, filgrastim, finasteride, fludarabine phosphate, formestane, fotemustine, gallium nitrate, gemcitabine, gemtuzumab zogamicin, gimeracil/oteracil/tegafur combination, glycopine, goserelin, heptaplatin, human

5 chorionic gonadotropin, human fetal alpha fetoprotein, ibandronic acid, idarubicin, (imiquimod, interferon alfa, interferon alfa, natural, interferon alfa-2, interferon alfa-2a, interferon alfa-2b, interferon alfa-N1, interferon alfa-n3, interferon alfacon-1, interferon alpha, natural,

10 interferon beta, interferon beta-1a, interferon beta-1b, interferon gamma, natural interferon gamma-1a, interferon gamma-1b, interleukin-1 beta, iobenguane, irinotecan, irsogladine, lanreotide, LC 9018 (Yakult), leflunomide, lenograstim, lentinan sulfate, letrozole, leukocyte alpha

15 interferon, leuprorelin, levamisole + fluorouracil, liarozole, lobaplatin, lonidamine, lovastatin, masoprocol, melarsoprol, metoclopramide, mifepristone, miltefosine, mirimostim, mismatched double stranded RNA, mitoguanzone, mitolactol, mitoxantrone, molgramostim, nafarelin, naloxone

20 + pentazocine, nartograstim, nedaplatin, nilutamide, noscapine, novel erythropoiesis stimulating protein, NSC 631570 octreotide, oprelvekin, osaterone, oxaliplatin, paclitaxel, pamidronic acid, pegaspargase, peginterferon alfa-2b, pentosan polysulfate sodium, pentostatin,

25 picibanil, pirarubicin, rabbit antithymocyte polyclonal antibody, polyethylene glycol interferon alfa-2a, porfimer sodium, raloxifene, raltitrexed, rasburicase, rhenium Re 186 etidronate, RII retinamide, rituximab, romurtide, samarium (153 Sm) lexidronam, sargramostim, sizofiran,

30 sobuzoxane, sonermin, strontium-89 chloride, suramin, tasonermin, tazarotene, tegafur, temoporfin, temozolomide, teniposide, tetrachlorodecaoxide, thalidomide, thymalfasin, thyrotropin alfa, topotecan, toremifene, tositumomab-iodine 131, trastuzumab, treosulfan, tretinoin, trilostane,

- 92 -

- trimetrexate, triptorelin, tumor necrosis factor alpha, natural, ubenimex, bladder cancer vaccine, Maruyama vaccine, melanoma lysate vaccine, valrubicin, verteporfin, vinorelbine, VIRULIZIN, zinostatin stimalamer, or
- 5 zoledronic acid; abarelix; AE 941 (Aeterna), ambamustine, antisense oligonucleotide, bcl-2 (Genta), APC 8015 (Dendreon), cetuximab, decitabine, dexaminoglutethimide, diaziquone, EL 532 (Elan), EM 800 (Endorecherche), eniluracil, etanidazole, fenretinide, filgrastim SD01
- 10 (Amgen), fulvestrant, galocitabine, gastrin 17 immunogen, HLA-B7 gene therapy (Vical), granulocyte macrophage colony stimulating factor, histamine dihydrochloride, ibritumomab tiuxetan, ilomastat, IM 862 (Cytran), interleukin-2, iproxifene, LDI 200 (Milkhaus), leridistim, lintuzumab, CA
- 15 125 MAb (Biomira), cancer MAb (Japan Pharmaceutical Development), HER-2 and Fc MAb (Medarex), idiotypic 105AD7 MAb (CRC Technology), idiotypic CEA MAb (Trilex), LYM-1-iodine 131 MAb (Techniclone), polymorphic epithelial mucin-yttrium 90 MAb (Antisoma), marimastat, menogaril,
- 20 mitumomab, motexafin gadolinium, MX 6 (Galderma), nelarabine, nolatrexed, P 30 protein, pegvisomant, pemetrexed, porfiromycin, prinomastat, RL 0903 (Shire), rubitecan, satraplatin, sodium phenylacetate, sparfosic acid, SRL 172 (SR Pharma), SU 5416 (SUGEN), TA 077
- 25 (Tanabe), tetrathiomolybdate, thaliblastine, thrombopoietin, tin ethyl etiopurpurin, tirapazamine, cancer vaccine (Biomira), melanoma vaccine (New York University), melanoma vaccine (Sloan Kettering Institute), melanoma oncolysate vaccine (New York Medical College),
- 30 viral melanoma cell lysates vaccine (Royal Newcastle Hospital), or valspodar.

Alternatively, the present compounds may also be used in co-therapies with other anti-neoplastic agents, such as other kinase inhibitors including p38 inhibitors and CDK

- 93 -

inhibitors, TNF inhibitors, metallomatrix proteases inhibitors (MMP), COX-2 inhibitors including celecoxib, rofecoxib, parecoxib, valdecoxib, and etoricoxib, NSAID's, SOD mimics or $\alpha_v\beta_3$ inhibitors.

5 The present invention comprises processes for the preparation of a compound of Formula I-X.

Also included in the family of compounds of Formula I-X are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts
10 commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I-X
15 may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, arylaliphatic,
20 heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, adipic, butyric, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic,
25 4-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, ethanedisulfonic, benzenesulfonic, pantothenic, 2-hydroxyethanesulfonic, toluenesulfonic, sulfanilic, cyclohexylaminosulfonic, camphoric, camphorsulfonic, digluconic,
30 cyclopentanepropionic, dodecylsulfonic, glucoheptanoic, glycerophosphonic, heptanoic, hexanoic, 2-hydroxyethanesulfonic, nicotinic, 2-naphthalenesulfonic, oxalic, palmoic, pectinic, persulfuric, 2-phenylpropionic, picric, pivalic propionic, succinic, tartaric, thiocyanic, mesylic,

- 94 -

undecanoic, stearic, algenic, β -hydroxybutyric, salicylic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I-X include metallic salts, such as salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc, or salts made from organic bases including primary, secondary and tertiary amines, substituted amines including cyclic amines, such as caffeine, arginine, diethylamine, N-ethyl piperidine, histidine, glucamine, isopropylamine, lysine, morpholine, N-ethyl morpholine, piperazine, piperidine, triethylamine, trimethylamine. All of these salts may be prepared by conventional means from the corresponding compound of the invention by reacting, for example, the appropriate acid or base with the compound of Formula I-X.

Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids that may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid. Other examples include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic bases. Preferred salts include hydrochloride, phosphate and edisylate.


Additional examples of such salts can be found in Berge et al., J. Pharm. Sci., 66, 1 (1977).

- 95 -

GENERAL SYNTHETIC PROCEDURES

The compounds of the invention can be synthesized according to the following procedures of Schemes 1-31, wherein the substituents are as defined for Formulas I-X, above, except where further noted.

Scheme 1



Substituted carboxamides 3 can be prepared from the corresponding halo analogs 1 by the process outlined in Scheme 1. Substituted amino acids 2 are prepared from the corresponding chloro compounds 1 such as by reacting with an amine at a suitable temperature, such as about 80°C. The acid 2 is coupled with an amine, preferably in the presence of a coupling agent such as EDC, to form the corresponding amide 3.

The amination process can be carried out as an Ullmann type reaction using a copper catalyst, such as copper[0] or a copper[I] compound such as copper[I]oxide, copper[I]bromide or copper[I]iodide in the presence of a suitable base (such as a metal carbonate, for example K₂CO₃ to neutralize the acid generated in the reaction.

- 96 -

This reaction is reviewed in Houben-Weyl "Methoden der Organischen Chemie", Band 11/1, page 32 -33, 1958, in Organic Reactions, 14, page 19-24, 1965 and by J. Lindley (1984) in Tetrahedron, 40, page 1433-1456. The amount of catalyst is typically in the range of 1 to 20 mole percent. The reaction is carried out in an inert, aprotic solvent such as an ether (for example dimethoxyethane or dioxane) or an amide (for example dimethylformamide or *N*-methylpyrrolidone), under an inert atmosphere in the temperature range of 60-180°C.

An alternative amination process involves using a Group VIII element, where the metal core of the catalyst should be a zero-valent transition metal, such as palladium or nickel, which has the ability to undergo oxidative addition to the aryl-halogen bond. The zero valent state of the metal may be generated in situ from the M[II] state. The catalyst complexes may include chelating ligands, such as alkyl, aryl or heteroaryl derivatives of phosphines or biphosphines, imines or arsines. Preferred catalysts contain palladium or nickel. Examples of such catalysts include palladium[II]chloride, palladium[II]acetate, tetrakis(triphenyl-phosphine)palladium[0] and nickel[II]acetylacetonate. The metal catalyst is typically in the range of 0.1 to 10 mole percent. The chelating ligands may be either monodentate, as in the case for example of trialkylphosphines, such as tributylphosphine, triarylphosphines, such as tri-(*ortho*-tolyl)phosphine, and triheteroaryl phosphines, such as tri-2-furylphosphine; or they may be bidentate such as in the case of 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, 1,2-bis(diphenylphosphino)ethane, 1,1'-bis(diphenylphosphino)ferrocene and 1-(*N,N*-dimethyl-amino)-1'-(dicyclohexylphosphino)biphenyl. The supporting ligand may be complexed to the metal center in the form of a metal


- 97 -

complex prior to being added to the reaction mixture or may be added to the reaction mixture as a separate compound. The supporting ligand is typically present in the range 0.01 to 20 mole percent. It is often necessary to add a suitable
5 base to the reaction mixture, such as a trialkylamine (for example, DIEA or 1,5-diazabicyclo[5,4,0]undec-5-ene), a Group I alkali metal alkoxide (for example potassium tert-butoxide) or carbonate (for example cesium carbonate) or potassium phosphate. The reaction is typically carried out
10 in an inert aprotic solvent such as an ether (for example dimethoxyethane or dioxane) or an amide (for example, DMF or *N*-methylpyrrolidone), under an inert atmosphere in the temperature range of 60-180°C.

The amination is preferably carried out in an inert,
15 aprotic, preferably anhydrous, solvent or solvent mixture, for example in a carboxylic acid amide, for example DMF or dimethylacetamide, a cyclic ether, for example THF or dioxane, or a nitrile, for example CH₃CN, or in a mixture thereof, at an appropriate temperature, for example in a
20 temperature range of from about 40°C to about 180°C, and if necessary under an inert gas atmosphere, for example a nitrogen or argon atmosphere.

- 98 -

Scheme 2



5

Substituted carboxamides 3 can be prepared from the corresponding halo analogs 1A by the process outlined in Scheme 2. The chloro acid 1 (LG is OH) is coupled with an amine, preferably in the presence of a coupling agent such as EDC, to form the corresponding chloro amide 4.

Substituted amino-amides 3 are prepared from the corresponding chloro compounds 4 such as by reacting with an amine at a suitable temperature, such as about 80°C. The amination reaction can be run in the presence of an appropriate catalyst such as a palladium catalyst, in the presence of an aprotic base such as sodium *t*-butoxide or cesium carbonate, or a nickel catalyst, or a copper catalyst.


Alternatively, carboxamides 3 can be prepared from 2-chloro-heterocyclyl acid chloride 1A (LG is Cl) by coupling first with R¹-NH₂, such as in the presence of base, e.g., NaHCO₃, in a suitable solvent, such as CH₂Cl₂, to form the

- 99 -

amide 1B, then coupling with a primary or secondary amine to yield the substituted carboxamide 3.

Additionally, where A is a pi-electron rich heterocycle, the addition of KF, such as 40% KF on alumina
 5 in IpOH, at a temperature over about 100°C, preferably about 160°C, can be used in the formation of 3 from 1B.

Scheme 3




10

3

Substituted carboxamides 3 can be prepared from the corresponding bromo/chloro analogs 5 by the process outlined in Scheme 3. The bromo/chloro acid 5 is coupled with an
 15 amine, preferably in the presence of a coupling agent such as EDC, to form the corresponding bromo substituted amide 6. Suzuki coupling with the bromo amide 6 and suitable boronic acids provides the substituted amide 4. Substituted amino-amides 3 are prepared from the corresponding chloro
 20 compounds 4 as described in Scheme 2.

- 100 -

Scheme 4



5

Substituted pyridines can be prepared by the process outlined in Scheme 4. A solution of sodium hypobromite is freshly prepared and added to a 2-hydroxynicotinic acid 7 and heated, preferably at a temperature at about 50°C.


- 10 Additional sodium hypobromide may be needed to form the bromo compound 8. The 5-bromo-2-hydroxynicotinic acid 8 is reacted with thionyl chloride, preferably at a temperature >RT, more preferably at about 80°C to form the 2-chloro-nicotinic acid analog 9. The acid is coupled with an amine,

- 101 -

preferably in the presence of EDC, HOBT, and DIEA to form the corresponding substituted amide 10. Suzuki coupling with the bromo amide and suitable boronic acids, provides the substituted nicotinamide 11. 2-Amino-nicotinamides 12
5 are prepared from the corresponding chloro compounds 11 such as by reacting with substituted amines at a suitable temperature, such as about 80°C.


Scheme 5

10



Alkylated indazoles can be prepared by the process outlined in Scheme 5. To a solution of 6-nitroindazole 13
15 in a solvent such as THF is added strong base, such as NaH at a temperature below RT, preferably at about 0°C. Alkylhalides, such as where Rⁿ is methyl, are added and reacted at a temperature about RT to give 1-alkyl-6-nitro-1H-indazole 14. The nitro indazole 14 is hydrogenated, such
20 as with an H₂ atmosphere in the presence of a catalyst, such as Pd/C to give the 1-substituted-6-amino-1H-indazole 15.


Scheme 6



- 102 -

Brominated indazoles can be prepared by the process outlined in Scheme 6. NBS is slowly added to an acidic solution, such as a mixture of TFA:H₂SO₄ (5:1) and tert-butyl-4-nitrobenzene 16 at a temperature of about RT to yield the brominated compound 17.

Scheme 7




10

Substituted anilines can be prepared by the process outlined in Scheme 7. A mixture of 1-(substituted)-2-bromo-4-nitrobenzene 18 (where R^x is a substituent selected from those available for substituted R²) and N-methylpiperazine is heated, such as with or without solvent, preferably without solvent, at a temperature above RT, preferably at a temperature above about 100°C, and more preferably at a temperature at about 130°C to give the 1-[5-(substituted)-2-nitrophenyl]-4-methylpiperazine 19. The nitro compound 19 is hydrogenated, such as with an H₂ atmosphere in the presence of a catalyst, such as Pd/C to furnish 4-(substituted)-2-(4-methylpiperazinyl)phenylamine 20.

20

- 103 -

Scheme 8




5

Tricyclic heterocycles can be prepared by the process outlined in Scheme 8. 7-Nitro-2,3,4-trihydroisoquinolin-1-one 21 is heated in POCl₃ at a temperature above RT, preferably at a temperature sufficient for reflux, to form the 1-chloro-7-nitro-3,4-dihydroisoquinoline 22. The 1-chloro-7-nitro-3,4-dihydroisoquinoline 22 is dissolved in a solvent, such as THF, and H₂NNH₂ is added. The reaction is heated with HC(OEt)₃ at a temperature above RT, preferably at a temperature above about 75°C, and more preferably at a temperature at about 115°C to give the nitro-substituted tricyclic. Hydrogenation, such as with an H₂ atmosphere in the presence of a catalyst, such as Pd/C, gives 2-amino-5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinoline 23.

- 104 -

Scheme 9



5


Indazolyl ethers can be prepared by the process outlined in Scheme 9. 6-Nitro-1H-2-hydroindazol-3-one **24** is protected such as with Boc_2O and DMAP in CH_2Cl_2 at a temperature of about RT, to give the protected 6-nitro-2-hydroindazol-3-one. The protected 6-nitro-2-hydroindazol-3-one is reacted with an alcohol (where R^* is an appropriate substituent selected from the possible substituents on R^1) and Ph_3P in a solvent, such as THF, and DEAD, at a temperature of about RT, to give the protected 6-nitro(indazol-3-yl) ether. The nitro intermediate is hydrogenated, such as with an H_2 atmosphere in the presence of a catalyst, such as Pd/C, to give the protected 6-amino(indazol-3-yl) ether **25**. The amine **25** is coupled with 2-chloronicotinic acid in a solvent, such as an alcohol, preferably pentanol, at a temperature above RT, preferably

- 105 -

at a temperature above about 75°C, and more preferably at a temperature at about 130°C to give the coupled and deprotected compound 26.

5

Scheme 10




Indolinyl substituted carboxamides can be prepared from the corresponding nitro indoline 27 by the process outlined in Scheme 10. For example, 3,3-dimethyl-6-nitroindoline 27 is alkylated, such as with N-protected-4-formylpiperidine in the presence of NaHB(OAc)₃ and acid, such as glacial AcOH, and solvent, such as dichloromethane, at a temperature of about RT, to afford the alkylated indane

- 106 -

28. Hydrogenation of the alkylated indane **28**, such as with an H₂ atmosphere in the presence of a catalyst, such as Pd/C, in the presence of a solvent, such as an alcohol, preferably MeOH, to give the amino intermediate **29**.
- 5 Alternatively, other hydrogenation methods can be used, such as Fe powder with NH₄Cl. Coupling of the amine **29**, such as with 2-chloronicotinic acid and DIEA, HOBt and EDC, in a solvent such as CH₂Cl₂ at a temperature of about RT provides the protected carboxamide **30**, which upon deprotection and
- 10 alkylation yields other compounds of the invention, **31** and **32**, respectively. Alternatively, amine **29** is reacted with 2-fluoronicotinoyl chloride to form a 2-fluoronicotinamide, which can be alkylated, such as in Scheme 10.

15

Scheme 11




- 107 -

Substituted anilines can be prepared by the process outlined in Scheme 11 (where R^x is a substituent selected those available for substituted R¹, preferably haloalkyl and alkyl). 1-Methyl-4-piperidinone is added to a solution of
5 strong base such as LiHMDS, in a solvent such as THF, at a temperature below RT, preferably lower than about -50°C, more preferably at about -78°C. Tf₂NPh is reacted with the enolate at a temperature of about RT, to give 1-methyl-4-(1,2,5,6-tetrahydro)pyridyl-(trifluoromethyl)sulfonate. A
10 mixture of the triflate intermediate, bis(pinacolato)diboron, potassium acetate, 1,1'-bis(diphenylphosphino)ferrocene-palladium dichloride, and bis(diphenylphosphino)ferrocene in a solvent such as dioxane and heated at a temperature above RT, preferably at a
15 temperature above about 50°C, and more preferably at a temperature at about 80°C to give 4,4,5,5-tetramethyl-2-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-1,3,2-dioxaborolane **34**. The substituted aniline **35** is formed from the 1,3,2-dioxaborolane **34** such as with treatment with an amine in the
20 presence of 1,1'-bis(diphenylphosphino)ferrocene-palladium dichloride and base, such as K₂CO₃, in a solvent such as DMF at a temperature above RT, preferably at a temperature above about 50°C, and more preferably at a temperature at about 80°C.

25

- 108 -

Scheme 12



5

Substituted anilines can be prepared by the process outlined in Scheme 12. 4-Cyano-4-phenylpiperidine hydrochloride **36** is treated with base, such as KOH , at a temperature above RT, preferably at a temperature above about 100°C , and more preferably at a temperature at about 160°C , to provide the phenyl piperidine **37**. Alkylation of the phenyl piperidine **37**, such as with formaldehyde and NaCNBH_3 in a solvent such as CH_3CN , with sufficient acid to maintain the reaction pH near 7, to provide the alkylated piperidine **38**. Nitration of the phenylpiperidine **38**, such as with H_2SO_4 and fuming HNO_3 at a temperature below RT, and preferably at about 0°C , gives the nitro intermediate **39**. Hydrogenation of the nitro intermediate **39**, such as with an H_2 atmosphere in the presence of a catalyst, such as Pd/C , in the presence of a solvent, such as an alcohol, preferably MeOH , to give the amino intermediate **40**.


10

15

20

- 109 -

Scheme 13




5

Substituted amides can be prepared by the process outlined in Scheme 13. 3-Nitrocinnamic acid 41 is coupled with 1-methylpiperazine in the presence of EDC and a solvent such as CH₂Cl₂, at a temperature of about RT gives the carboxamide 42.

10

Scheme 14



15

1,2,3,6-Tetrahydro-pyridyl substituted anilines are prepared such as by the procedure described in Scheme 14. Nitrobenzenes 43 are brominated, such as with bromine in the presence of acid, H₂SO₄ for example, or with NBS to yield the 3-bromo derivative 44. Suzuki coupling of the bromo-derivative 44 and a substituted pyridylboronic acid, such as


20

- 110 -

at a temperature above RT, preferably above about 50°C, and more preferably at about 80°C, yields the pyridyl derivative 45. Alkylation of the nitrophenyl-pyridine 45, such as by treatment with iodomethane, preferably above about 50°C, and more preferably at about 80°C, yields the pyridinium compound 46, which upon reduction, such as by NaBH₄, yields the tetrahydropyridine 47.

Scheme 15

10




A series of substituted anilines are prepared such as by the procedure described in Scheme 15. A nitrobenzyl bromide 48 is coupled with morpholine, such as at a temperature at about RT, to yield the heterocyclylmethyl nitrobenzene derivative. Reduction of the nitro compound, such as with iron powder, preferably above about 50°C, and

more preferably at about 80°C, yields the heterocyclylmethyl substituted aniline 49.

Protected alkylamine substituted anilines can be prepared from the nitro free amines 50, such as with standard protecting agents and chemistry known in the art, such as BOC chemistry. Reduction of the protected nitro compound, such as with iron powder, preferably above about 50°C, and more preferably at about 80°C, yields the aniline 51.

10 Sulfonamide substituted anilines can be prepared from
nitrobenzenesulfonyl chlorides 52. Coupling of
nitrobenzenesulfonyl chlorides 52 with reactive heterocyclic
compounds, such as substituted piperazines, piperidines, and
the like, in a protic solvent such as EtOH, such as at a
15 temperature about RT, yields the nitrobenzenesulfonamides 52.
Reduction of the nitro benzenesulfonamide, such as with iron
powder, preferably above about 50°C, and more preferably at
about 80°C, yields the aniline 53.

Scheme 16



A series of perhaloalkyl-substituted anilines **56**,
 25 where R^y represents perhaloalkyl radicals, are prepared such
 as by the procedure described in Scheme 16. 1-Nitro-4-
 (perfluoroethyl)benzene can be synthesized by the method
 described in the reference [John N. Freskos, Synthetic
 Communications, 18(9), 965-972 (1988)]. Alternatively, 1-
 30 Nitro-4-(perfluoroalkyl)benzene can be synthesized from the
 nitro compound, where X^a is a leaving group, such as bromo


- 112 -

or iodo, by the method described by W. A. Gregory, et al. [J. Med. Chem., 1990, 33, 2569-2578].

Reduction of the nitrobenzenes 55, such as with iron powder, at a temperature above about 50°C, and preferably at about 80°C, yields the aniline 56. Hydrogenation, such as with H₂ atmosphere in the presence of a catalyst, such as 10% Pd/C, is also possible.

Scheme 17

10



Additional series of substituted anilines are prepared such as by the procedures described in Scheme 17 (where R^x is a substituent selected those available for substituted R¹, preferably haloalkyl and alkyl). 2-Alkoxy substituted anilines 59 are prepared from the corresponding phenol compounds 57 such as by the Mitsunobu reaction, including treatment with a N,N-dialkylethanolamine and PPh₃ and DEAD to give the corresponding nitro compound 58, followed by hydrogenation, such as with H₂ to give the aniline 59.


- 113 -

Alternatively, piperazinyl substituted anilines 62 can be prepared by the treatment of an aniline 60 with an N-substituted-bis(2-chloroethyl)amine, base, such as K_2CO_3 and NaI, at a temperature above about $50^\circ C$, preferably above about $100^\circ C$, and more preferably at about $170^\circ C$, to give the piperazinylbenzene compound 61. Nitration, such as with H_2SO_4 and HNO_3 , at a temperature above $0^\circ C$, and preferably at about RT, followed by hydrogenation, such as with H_2 atmosphere gives the substituted aniline 62.

Alternatively, piperazinyl substituted anilines 65 can be prepared by the treatment of a fluoro-nitro-substituted aryl compounds 63. The fluoro-nitro-substituted aryl compounds 63 and 1-substituted piperazines are heated, preferably neat, at a temperature above about $50^\circ C$, and preferably at about $90^\circ C$, to yield the piperazinyl-nitroaryl compounds 64. Hydrogenation, such as with H_2 atmosphere in the presence of a catalyst, such as 10% Pd/C, gives the substituted aniline 65.

20

Scheme 18




- 114 -

Substituted indolines are prepared such as by the procedures described in Scheme 18. Substituted amino-indolines **68** are prepared from the nitroindoline **66** and a ketone in the presence of $\text{NaHB}(\text{OAc})_3$ to form the 1-substituted indoline **67**. The nitroindoline **67** is hydrogenated, such as with H_2 in the presence of a catalyst, such as Pd/C , to yield the amino-indoline **68**.

Alternatively, substituted amino-indolines **71** are prepared from the nitroindoline **66**. Nitroindoline **66**, is reacted with an acid chloride to form an amide. Further treatment with a primary or secondary amine, preferably a secondary amine, such as in the presence of NaI , at a temperature above about 50°C , and preferably at about 70°C yields the nitroindoline **69**. The nitro compound **69** is hydrogenated, such as with H_2 in the presence of a catalyst, such as Pd/C , to yield the amino-indoline) **70**. The carbonyl is reduced, such as with $\text{BH}_3\text{-THF}$ yields 1-aminoalkyl-indolines **71**.

20

Scheme 19




- 115 -

Substituted indolines are prepared such as by the procedures described in Scheme 19. Substituted acetamides 73 are prepared from the coupling of halo-5-nitroanilines 72 (where LG is bromo or chloro, preferably chloro) and an acylating agent, such as acetyl chloride or acetic anhydride, under standard coupling chemistry, such as with DIEA, and DMAP, at a temperature of about RT, in a suitable solvent, such as CH₂Cl₂, DMF and/or DMAC. The N-(2-methylprop-2-enyl)acetamide 74 is prepared from the acetamide 73, such as by the treatment of base, such as NaH in a suitable solvent such as NMP or anhydrous DMF and a 3-halo-2-methylpropene such as 3-bromo-2-methylpropene or 3-chloro-2-methylpropene, at a temperature between about 0°C and RT, and preferably at about RT; or with CsCO₃ at a temperature above RT, preferably above about 50°C and more preferably above about 60°C. Cyclization of the N-(2-methylprop-2-enyl)acetamide 74, such as by the Heck-type reaction (treatment with Pd(OAc)₂ in the presence of base, for example tetraethyl-ammonium chloride, sodium formate, and NaOAc) at a temperature above about 50°C, and preferably at about 80°C, yields the protected (3,3-dimethyl-6-nitro-2,3-dihydro-indol-1-yl)ethanone 75. Deprotection, such as with strong acid such as HCl or AcOH at a temperature above about 50°C, and preferably at about 70-80°C, yields the 3,3-dimethyl-6-nitro-2,3-dihydro-indol-1-yl 76. Alternatively, the protected dihydro-6-nitro indoline 75 can be reduced, such as with Fe, or with 10% Pd/C in the presence of an excess of NH₄CO₂H, or with H₂ in the presence of a catalyst to form the protected dihydro-6-amino indoline 75a.

30

- 116 -

Scheme 20



5

Substituted anilines are prepared such as by the procedures described in Scheme 20. Nitrophenyl esters 78 are formed from the acid 77, such as by treatment with MeOH and acid. Alkylation of the ester 78, such as by treatment


10 with base, followed by alkyl halide, yields the branched alkyl compounds 79. Reduction of the ester 79, such as with BH_3 , yields the alcohol 80. The aldehyde 81 is prepared from the alcohol 80, such as by treatment with TPAP in the presence of N-methylmorpholine-N-oxide. Subsequent

15 treatment with methoxymethyltriphenylphosphonium chloride and KHMDS yields 81. Coupling of the aldehyde 81 and morpholine, such as with NaBH(OAc)_3 , yields the tertiary amine 82. Reduction of the nitro compound, such as with acid, for example AcOH, and zinc yields the aniline 83.

20

- 117 -

Scheme 21




5

Substituted aniline compounds are prepared such as by the procedure described in Scheme 21 (where R^x is a substituent selected those available for substituted R^1 , preferably haloalkyl and alkyl). Alkynyl-aniline 90, prepared similar to that described in Scheme 22, is hydrogenated such as with H_2 in the presence of a catalyst, such as $\text{Pd}(\text{OH})_2$, to yield the substituted alkyl 91.

10

Scheme 22

15




20

Substituted bromophenyl compounds are prepared such as by the procedure described in Scheme 22. Bromine is added to a optionally substituted nitrobenzene 92, silver(II)sulfate and acid, such as H_2SO_4 , to provide the bromo derivative 93.

- 118 -

Scheme 23




5

Substituted anilines are prepared such as by the procedure described in Scheme 23 (where R^t and R^v are alkyl, or together with the nitrogen atom form a 4-6 membered heterocyclic ring). Acryloyl chloride **94** is reacted with an amine, preferably a secondary amine, such as at a temperature between about 0°C and about RT, to form the amide **95**. A bromo-nitrobenzene **93** is reacted with the amide **95**, such as in the presence of base, for example TEA, together with Pd(OAc)_2 and $\text{Pd(PPh}_3)_4$, at a temperature above about 50°C, and preferably at about 120°C, such as in a sealed container, to form the substituted alkene **96**. Hydrogenation of the alkene **96**, such as with H_2 -in the presence of a catalyst, for example Pd/C catalyst yields the substituted aniline **97**. Reduction of the amide **97**, such as with LiAlH_4 , at a temperature above about 50°C, and preferably at about 80°C yields the aniline **98**.

20


- 119 -

Scheme 24



5 Substituted indoles are prepared such as by the
 procedure described in Scheme 24. A nitroindole 99 is
 coupled with a halo compound, in the presence of base, for
 example K_2CO_3 . Heating at a temperature above about $50^\circ C$,
 and preferably at about reflux yields the substituted-nitro-
 10 1H-indole 100. Hydrogenation similar to conditions
 described above yield the amino derivative 101.

Scheme 25



15


Substituted aminomethyl compounds are prepared such as
 by the procedure described in Scheme 25. NaH is added to an
 alcohol and heated at a temperature above about RT, and
 20 preferably at about $50^\circ C$, to form the sodium alkoxide, which
 is added to a halo compound 102, such as a 2-chloro-4-
 cyanopyridine and heated at a temperature above about $50^\circ C$,

- 120 -

and preferably at about 70°C to form the ether 103.
Hydrogenation yields the aminomethyl derivative 104.


Scheme 26

5



Substituted anilines are prepared such as by the
procedure described in Scheme 26. Treatment with the
haloalkyl alcohol 105 with an alcohol, such as in the
presence of DEAD and PPh₃ yields the ether 107 or 106.

Scheme 27




Functionalized pyridines are prepared such as by the
procedure described in Scheme 27. 2-Fluoropyridine 108 is
treated with base, such as LDA, at a temperature below about
0°C, and preferably at about -78°C, and quenched with a
stream of dry CO₂ to form the nicotinic acid 109.
Alternatively, solid CO₂ (dry ice) can be used, preferably
dried with N₂ prior to use. The acid 109 is converted to the
acid halide 110, such as by treatment with thionyl chloride

- 121 -

and heating at a temperature above about 50°C, and preferably at about reflux.


Scheme 28

5



Chloro-substituted pyridines 113 are prepared such as by the procedure described in Scheme 28. 2-Chloronicotinic acid 112 is activated with ethyl chloroformate, in the presence of base, such as TEA, at a temperature of about RT. Reaction with an amine produces amide 113. Alternatively, the amine can be coupled with the acid chloride 111, such as with polymer-supported DIPEA. Excess acid chloride is removed by treating the reaction mixture with polymer-supported trisamine resin, to form amide 113.

Scheme 29




Amino-substituted indoles 116 are prepared such as by the procedure described in Scheme 291. Nitroindoline 114 is

- 122 -

reacted with N-methyl-4-piperidone in the presence of NaOMe at a temperature above about 50°C, and preferably at about reflux, to form the 3-substituted indole **115**. Hydrogenation as previously discussed yields the amino indole **116**.

5

Scheme 30




10 Substituted carboxamides **118** can be prepared from the corresponding phenols **117** of the invention, by the process outlined in Scheme 30. A carboxamide **117** is coupled with an alcohol, such as 4-hydroxy-N-methylpiperidine, in the presence of DEAD and triphenylphosphine, in a solvent such as THF, at a temperature of about RT, provides the ether **118**.

15

- 123 -

Scheme 31



5

2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluoren-6-ylamine may be prepared by the method found in Scheme 31.

Nitrobenzylpyridines **119** are alkylated, such as with MeI, in the presence of TBAI and base to form the pyridinium compound **120**. The pyridinium compounds **120** are halogenated, such as brominated with NBS, to form the brominated pyridinium compounds **121** which are reduced such as with NaBH_4 , dehalogenated and reduced to form the hexahydro-fluorenes **124**.

15

The starting compounds defined in Schemes 1-31 may also be present with functional groups in protected form if necessary and/or in the form of salts, provided a salt-forming group is present and the reaction in salt form is possible. If so desired, one compound of formula I-X can be converted into another compound of formula I-X or a N-oxide

20

- 124 -

thereof; a compound of formula I-X can be converted into a salt; a salt of a compound of formula I-X can be converted into the free compound or another salt; and/or a mixture of isomeric compounds of formula I-X can be separated into the
5 individual isomers.

N-Oxides can be obtained in a known matter by reacting a compound of formula I-X with hydrogen peroxide or a peracid, e.g. 3-chloroperoxy-benzoic acid, in an inert solvent, e.g. dichloromethane, at a temperature between
10 about -10-35°C, such as about 0°C - RT.

If one or more other functional groups, for example carboxy, hydroxy, amino, or mercapto, are or need to be protected in a compound of formula I-X or in the preparation of compounds of formula I-X, because they should not take
15 part in the reaction, these are such groups as are usually used in the synthesis of peptide compounds, and also of cephalosporins and penicillins, as well as nucleic acid derivatives and sugars.

The protecting groups may already be present in
20 precursors and should protect the functional groups concerned against unwanted secondary reactions, such as acylations, etherifications, esterifications, oxidations, solvolysis, and similar reactions. It is a characteristic of protecting groups that they lend themselves readily, i.e.
25 without undesired secondary reactions, to removal, typically by solvolysis, reduction, photolysis or also by enzyme activity, for example under conditions analogous to physiological conditions, and that they are not present in the end-products. The specialist knows, or can easily
30 establish, which protecting groups are suitable with the reactions mentioned above and hereinafter.

The protection of such functional groups by such protecting groups, the protecting groups themselves, and their removal reactions are described for example in

- 125 -

standard reference works, such as J. F. W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London and New York 1973, in T. W. Greene, "Protective Groups in Organic Synthesis", Wiley, New York 1981, in "The Peptides"; Volume 3 (editors: E. Gross and J. Meienhofer), Academic Press, London and New York 1981, in "Methoden der organischen Chemie" (Methods of organic chemistry), Houben Weyl, 4th edition, Volume 15/1, Georg Thieme Verlag, Stuttgart 1974, in H.-D. Jakubke and H. Jescheit, "Aminosäuren, Peptide, Proteine" (Amino acids, peptides, proteins), Verlag Chemie, Weinheim, Deerfield Beach, and Basel 1982, and in Jochen Lehmann, "Chemie der Kohlenhydrate: Monosaccharide und Derivate" (Chemistry of carbohydrates: monosaccharides and derivatives), Georg Thieme Verlag, Stuttgart 1974.

In the additional process steps, carried out as desired, functional groups of the starting compounds which should not take part in the reaction may be present in unprotected form or may be protected for example by one or more of the protecting groups mentioned above under "protecting groups". The protecting groups are then wholly or partly removed according to one of the methods described there.

Salts of a compound of formula I-X with a salt-forming group may be prepared in a manner known *per se*. Acid addition salts of compounds of formula I-X may thus be obtained by treatment with an acid or with a suitable anion exchange reagent. A salt with two acid molecules (for example a dihalogenide of a compound of formula I) may also be converted into a salt with one acid molecule per compound (for example a monohalogenide); this may be done by heating to a melt, or for example by heating as a solid under a high vacuum at elevated temperature, for example from 130 to

- 126 -

170°C, one molecule of the acid being expelled per molecule of a compound of formula I-X.

Salts can usually be converted to free compounds, e.g. by treating with suitable basic agents, for example with
5 alkali metal carbonates, alkali metal hydrogen carbonates, or alkali metal hydroxides, typically potassium carbonate or sodium hydroxide.

A compound of formula I-X, wherein Z is oxygen, can be converted into the respective compound wherein Z is sulfur, for example, by using an appropriate sulfur compound, e. g.
10 using reaction with Lawesson's reagent (2,4-bis-(4-methoxyphenyl)2,4-dithioxo-1,2,3,4-dithiaphosphetan) in a halogenated hydrocarbon, such as CH_2Cl_2 , or an aprotic solvent, such as toluene or xylene, at temperatures from
15 about 30°C to reflux.

All process steps described here can be carried out under known reaction conditions, preferably under those specifically mentioned, in the absence of or usually in the presence of solvents or diluents, preferably such as are
20 inert to the reagents used and able to dissolve these, in the absence or presence of catalysts, condensing agents or neutralizing agents, for example ion exchangers, typically cation exchangers, for example in the H^+ form, depending on the type of reaction and/or reactants at reduced, normal, or
25 elevated temperature, for example in the range from about -100°C to about 190°C, preferably from about -80°C to about 150°C, for example at about -80 to about 60°C, at room temperature, at about -20 to about 40°C or at the boiling point of the solvent used, under atmospheric pressure or in
30 a closed vessel, where appropriate under pressure, and/or in an inert atmosphere, for example under argon or nitrogen.

Salts may be present in all starting compounds and transients, if these contain salt-forming groups. Salts may

- 127 -

also be present during the reaction of such compounds, provided the reaction is not thereby disturbed.

In certain cases, typically in hydrogenation processes, it is possible to achieve stereoselective reactions, allowing for example easier recovery of individual isomers.

The solvents from which those can be selected which are suitable for the reaction in question include for example water, esters, typically lower alkyl-lower alkanoates, e.g., ethyl acetate, ethers, typically aliphatic ethers, e.g., diethylether, or cyclic ethers, e.g., THF, liquid aromatic hydrocarbons, typically benzene or toluene, alcohols, typically MeOH, EtOH or 1-propanol, IPOH, nitriles, typically CH_3CN , halogenated hydrocarbons, typically CH_2Cl_2 , acid amides, typically DMF, bases, typically heterocyclic nitrogen bases, e.g. pyridine, carboxylic acids, typically lower alkanecarboxylic acids, e.g., AcOH, carboxylic acid anhydrides, typically lower alkane acid anhydrides, e.g., acetic anhydride, cyclic, linear, or branched hydrocarbons, typically cyclohexane, hexane, or isopentane, or mixtures of these solvents, e.g., aqueous solutions, unless otherwise stated in the description of the process. Such solvent mixtures may also be used in processing, for example in chromatography.

The invention relates also to those forms of the process in which one starts from a compound obtainable at any stage as a transient and carries out the missing steps, or breaks off the process at any stage, or forms a starting material under the reaction conditions, or uses said starting material in the form of a reactive derivative or salt, or produces a compound obtainable by means of the process according to the invention and processes the said compound *in situ*. In the preferred embodiment, one starts

- 128 -

from those starting materials which lead to the compounds described above as preferred.

The compounds of formula I-X, including their salts, are also obtainable in the form of hydrates, or their
5 crystals can include for example the solvent used for crystallization (present as solvates).

New starting materials and/or intermediates, as well as processes for the preparation thereof, are likewise the subject of this invention. In the preferred embodiment, such
10 starting materials are used and reaction conditions so selected as to enable the preferred compounds to be obtained.

Starting materials of the invention, are known, are commercially available, or can be synthesized in analogy to
15 or according to methods that are known in the art.

For example, amine 1 can be prepared by reduction of the corresponding nitro. The reduction preferably takes place in the presence of a suitable reducing agent, such as tin(II) chloride or hydrogen in the presence of an
20 appropriate catalyst, such as Raney nickel (then preferably the hydrogen is used under pressure, e.g. between 2 and 20 bar) or PtO_2 , in an appropriate solvent, e.g. an alcohol, such as MeOH. The reaction temperature is preferably between about 0°C and about 80°C, especially about 15°C to about
25 30°C.

It would also be possible to reduce the nitro compound after forming the amide compound under reaction conditions analogous to those for the reduction of nitro compounds described above. This would eliminate the need to protect
30 the free amino group as described in Scheme 1.

In the preparation of starting materials, existing functional groups which do not participate in the reaction should, if necessary, be protected. Preferred protecting

- 129 -

groups, their introduction and their removal are described above or in the examples.


All remaining starting materials are known, capable of being prepared according to known processes, or commercially obtainable; in particular, they can be prepared using processes as described in the examples.

Compounds of the present invention can possess, in general, one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or non-racemic mixtures thereof. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, e.g., by formation of diastereoisomeric salts, by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric, and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts. A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting compounds of the invention with an optically pure acid in an activated form or an optically pure isocyanate. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active compounds of the invention can likewise be obtained by using optically active starting materials. These isomers may be in the form of a free acid, a free base, an ester or a salt.

- 130 -

The compounds of this invention may contain one or more asymmetric centers and thus occur as racemates and racemic mixtures, scalemic mixtures, single enantiomers, individual diastereomers and diastereomeric mixtures. All such isomeric forms of these compounds are expressly included in the present invention.

The compounds of this invention may also be represented in multiple tautomeric forms, for example, as illustrated below:



The invention expressly includes all tautomeric forms of the compounds described herein.

The compounds may also occur in cis- or trans- or E- or Z- double bond isomeric forms. All such isomeric forms of such compounds are expressly included in the present invention. All crystal forms of the compounds described herein are expressly included in the present invention.

Substituents on ring moieties (e.g., phenyl, thienyl, etc.) may be attached to specific atoms, whereby they are intended to be fixed to that atom, or they may be drawn unattached to a specific atom, whereby they are intended to be attached at any available atom that is not already substituted by an atom other than H (hydrogen).

The compounds of this invention may contain heterocyclic ring systems attached to another ring system. Such heterocyclic ring systems may be attached through a carbon atom or a heteroatom in the ring system.

Alternatively, a compound of any of the formulas delineated herein may be synthesized according to any of the processes delineated herein. In the processes delineated

- 131 -

herein, the steps may be performed in an alternate order and may be preceded, or followed, by additional protection/deprotection steps as necessary. The processes may further comprise use of appropriate reaction conditions, including inert solvents, additional reagents, such as bases (e.g., LDA, DIEA, pyridine, K_2CO_3 , and the like), catalysts, and salt forms of the above. The intermediates may be isolated or carried on *in situ*, with or without purification. Purification methods are known in the art and include, for example, crystallization, chromatography (liquid and gas phase, simulated moving bed ("SMB")), extraction, distillation, trituration, reverse phase HPLC and the like. Reaction conditions such as temperature, duration, pressure, and atmosphere (inert gas, ambient) are known in the art and may be adjusted as appropriate for the reaction.

As can be appreciated by the skilled artisan, the above synthetic schemes are not intended to comprise a comprehensive list of all means by which the compounds described and claimed in this application may be synthesized. Further methods will be evident to those of ordinary skill in the art. Additionally, the various synthetic steps described above may be performed in an alternate sequence or order to give the desired compounds. Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing the inhibitor compounds described herein are known in the art and include, for example, those such as described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, 3rd. Ed., John Wiley and Sons (1999); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); A. Katritzky and A. Pozharski,

- 132 -

Handbook of Heterocyclic Chemistry, 2nd Ed. (2001); M. Bodanszky, A. Bodanszky: *The practice of Peptide Synthesis* Springer-Verlag, Berlin Heidelberg 1984; J. Seyden-Penne: *Reductions by the Alumino- and Borohydrides in Organic*
5 *Synthesis*, 2nd Ed., Wiley-VCH, 1997; and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995).

The compounds of this invention may be modified by appending appropriate functionalities to enhance selective
10 biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological compartment (e.g., blood, lymphatic system, central nervous system), increase oral availability, increase solubility to allow administration by injection,
15 alter metabolism and alter rate of excretion.

The following examples contain detailed descriptions of the methods of preparation of compounds of Formulas I-X. These detailed descriptions fall within the scope, and serve to exemplify, the above described General Synthetic
20 Procedures which form part of the invention. These detailed descriptions are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention.

Unless otherwise noted, all materials were obtained
25 from commercial suppliers and used without further purification. Anhydrous solvents such as DMF, THF, CH₂Cl₂ and toluene were obtained from the Aldrich Chemical Company. All reactions involving air- or moisture-sensitive compounds were performed under a nitrogen atmosphere. Flash
30 chromatography was performed using Aldrich Chemical Company silica gel (200-400 mesh, 60A) or Biotage pre-packed column. Thin-layer chromatography (TLC) was performed with Analtech gel TLC plates (250 μ). Preparative TLC was performed with Analtech silica gel plates (1000-2000 μ). Preparative HPLC

- 133 -

was conducted on a Beckman or Waters HPLC system with 0.1% TFA/H₂O and 0.1% TFA/CH₃CN as mobile phase. The flow rate was at 20 ml/min. and gradient method was used. ¹H NMR spectra were determined with super conducting FT NMR spectrometers operating at 400 MHz or a Varian 300 MHz instrument. Chemical shifts are expressed in ppm downfield from internal standard tetramethylsilane. All compounds showed NMR spectra consistent with their assigned structures. Mass spectra (MS) were determined on a Perkin Elmer - SCIEX API 165 electrospray mass spectrometer(positive and, or negative) or an HP 1100 MSD LC-MS with eletrospray ionization and quadrupole detection. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated.

15

- 134 -

The following abbreviations are used:

	AIBN -	2,2'-azobisisobutyronitrile
5	Ar -	argon
	AgSO ₄ -	silver sulfate
	ATP -	adenosine triphosphate
	BH ₃ -	borane
	Boc -	tert-butyloxycarbonyl
10	Boc ₂ O -	Boc anhydride
	BOP-Cl -	bis(2-oxo-3-oxazolidinyl)phosphinic chloride
	Br ₂ -	bromine
	BSA -	bovine serum albumin
15	t-BuOH -	tert-butanol
	CAN -	ammonium cerium(IV) nitrate
	CH ₃ CN, AcCN -	acetonitrile
	CH ₂ Cl ₂ -	dichloromethane
	CH ₃ I, MeI -	iodomethane, methyl iodide
20	CCl ₄ -	carbon tetrachloride
	CCl ₃ -	chloroform
	CO ₂ -	carbon dioxide
	Cs ₂ CO ₃ -	cesium carbonate
	DIEA -	diisopropylethylamine
25	CuI -	copper iodide
	DCE -	1,2-dichloroethane
	DEAD -	diethyl azodicarboxylate
	DIEA -	diisopropylethylamine
	dppf -	1,1-diphenylphosphinoferrocene
30	DMAP -	4-(dimethylamino)pyridine
	DMAC -	N,N-dimethylacetamide
	DMF -	dimethylformamide
	DMSO -	dimethylsulfoxide
	DTT -	dithiothreitol
35	EDC, EDAC-	1-(3-dimethylaminopropyl)-3-

- 135 -

		ethylcarbodiimide hydrochloride
	EGTA -	ethylene glycol-bis(β -aminoethyl ether) - N,N,N',N'-tetraacetic acid
	EtOAc -	ethyl acetate
5	EtOH -	ethanol
	Et ₂ O -	diethyl ether
	Fe -	iron
	g -	gram
	h -	hour
10	HATU -	O-(7-azabenzotriazol-1-yl)-N,N,N',N'- tetramethyluronium hexafluorophosphate
	H ₂ -	hydrogen
	H ₂ O -	water
	HCl -	hydrochloric acid
15	H ₂ SO ₄ -	sulfuric acid
	H ₂ NNH ₂ -	hydrazine
	HC(OEt) ₃ -	triethylorthoformate
	HCHO, H ₂ CO -	formaldehyde
	HCO ₂ Na -	sodium formate
20	HOAc, AcOH -	acetic acid
	HOAt -	1-hydroxy-7-azabenzotriazole
	HOBt -	hydroxybenzotriazole
	IpOH -	isopropanol
	K ₂ CO ₃ -	potassium carbonate
25	KHMDS -	potassium hexamethylsilazane
	KNO ₃ -	potassium nitrate
	KOAc -	potassium acetate
	KOH -	potassium hydroxide
	LAH, LiAlH ₄ -	lithium aluminum hydride
30	LDA -	lithium diisopropylamide
	LiCl -	lithium chloride
	LiHMDS -	lithium hexamethyldisilazide
	MeOH -	methanol
	MgCl ₂ -	magnesium chloride

- 136 -

	MgSO ₄ -	magnesium sulfate
	mg -	milligram
	ml -	milliliter
	MnCl ₂ -	manganese chloride
5	NBS -	N-bromosuccinimide
	NMO -	4-methylmorpholine, N-oxide
	NMP -	N-methylpyrrolidone
	Na ₂ SO ₄ -	sodium sulfate
	Na ₂ S ₂ O ₅ -	sodium metabisulfite
10	NaHCO ₃ -	sodium bicarbonate
	Na ₂ CO ₃ -	sodium carbonate
	NaCl -	sodium chloride
	NaH -	sodium hydride
	NaI -	sodium iodide
15	NaOH -	sodium hydroxide
	NaOMe -	sodium methoxide
	NaCNBH ₃ -	sodium cyanoborohydride
	NaBH ₄ -	sodium borohydride
	NaNO ₂ -	sodium nitrate
20	NaBH(OAc) ₃ -	sodium triacetoxyborohydride
	NH ₄ Cl -	ammonium chloride
	N ₂ -	nitrogen
	Pd/C -	palladium on carbon
	PdCl ₂ (PPh ₃) ₂ -	palladium chloride bis(triphenylphosphine)
25	PdCl ₂ (dppf) -	1,1-bis(diphenylphosphino)ferrocene palladium chloride
	Pd(PPh ₃) ₄ -	palladium tetrakis triphenylphosphine
	Pd(OH) ₂ -	palladium hydroxide
	Pd(OAc) ₂ -	palladium acetate
30	PMB -	para methoxybenzyl
	POCl ₃ -	phosphorus oxychloride
	PPh ₃ -	triphenylphosphine
	PtO ₂ -	platinum oxide
	RT -	room temperature

- 137 -

	SiO ₂ -	silica
	SOCl ₂ -	thionyl chloride
	TBAI -	tetrabutylammonium iodide
	TEA -	triethylamine
5	Tf ₂ NPh -	N-phenyltrifluoromethanesulfonimide
	TFA -	trifluoroacetic acid
	THF -	tetrahydrofuran
	TPAP -	tetrapropylammonium perruthenate
	Tris-HCl -	Tris(hydroxymethyl)aminomethane
10		hydrochloride salt
	Zn -	zinc

Preparation I - 3-nitro-5-trifluoromethyl-phenol

1-Methoxy-3-nitro-5-trifluoromethyl-benzene (10g, Aldrich)
15 and pyridine-HCl (41.8g, Aldrich) were mixed together and heated neat at 210°C in an open flask. After 2.5 h the mixture was cooled to RT and partitioned between 1N HCl and EtOAc. The EtOAc fraction was washed with 1N HCl (4x), brine (1x), dried with Na₂SO₄, filtered and concentrated in vacuo
20 to form 3-nitro-5-trifluoromethyl-phenol as an off-white solid.

Preparation II - 1-Boc-4-(3-nitro-5-trifluoromethyl-phenoxy)-piperidine

25 3-Nitro-5-trifluoromethyl-phenol (8.81g) was dissolved in THF (76 ml). 1-Boc-4-hydroxy-piperidine (8.81 g, Aldrich) and Ph₃P (11.15 g) were added and the solution was cooled to -20°C. A solution of DEAD (6.8 ml, Aldrich) in THF (36 ml) was added dropwise, maintaining the temperature between -20
30 and -10°C. The reaction was warmed to RT and stirred overnight. The reaction was concentrated in vacuo and triturated with hexane. The yellow solid was removed by filtration and washed with Et₂O (25 ml), and hexane. The white filtrate was washed with 1N NaOH (2x), brine (1x) and

- 138 -

the hexane layer was dried over Na_2SO_4 , filtered and concentrated in vacuo. The crude material was purified with flash chromatography (SiO_2 , 5-10% EtOAc/hexane) to obtain 1-Boc-4-(3-nitro-5-trifluoromethyl-phenoxy)-piperidine.

5

The following compounds were prepared similarly to the procedure outlined above:

- 10 a) (S)-1-Boc-[2-(5-nitro-2-trifluoromethylphenoxy)methyl]-pyrrolidine
- b) (R)-1-Boc-[2-(5-nitro-2-trifluoromethylphenoxy)methyl]-pyrrolidine.
- c) (R) 1-Boc-2-(3-Nitro-5-trifluoromethyl-phenoxy)methyl)-pyrrolidine
- 15 d) 4-(2-tert-Butyl-5-nitro-phenoxy)methyl)-1-methyl-piperidine.
- e) (S) 1-Boc-2-(3-Nitro-5-trifluoromethyl-phenoxy)methyl)-pyrrolidine
- f) 1-Boc-3-(5-nitro-2-pentafluoroethyl-phenoxy)methyl)-azetidine.
- 20 g) N-Boc-[2-(5-nitro-2-pentafluoroethyl-phenoxy)-ethyl]amine.
- h) (R) 3-(2-tert-Butyl-5-nitro-phenoxy)methyl)-1-Boc-pyrrolidine.
- 25 i) 3-(2-tert-Butyl-5-nitro-phenoxy)methyl)-1-Boc-azetidine.
- j) (S)-1-Boc-[2-(5-nitro-2-tert-butylphenoxy)methyl]-pyrrolidine
- k) (S) 3-(2-tert-Butyl-5-nitro-phenoxy)methyl)-1-Boc-pyrrolidine.
- 30 l) (R)-1-Boc-[2-(5-nitro-2-tert-butylphenoxy)methyl]-pyrrolidine

- 139 -

Preparation III - 1-Boc-4-(3-amino-5-trifluoromethyl-phenoxy)-piperidine

1-Boc-4-(3-nitro-5-trifluoromethyl-phenoxy)-piperidine (470
5 mg) was dissolved in MeOH (12 ml) and Pd/C (10 mg) was
added. After sparging briefly with H₂, the mixture was
stirred under H₂ for 6 H. The catalyst was removed by
filtration and the MeOH solution was concentrated in
vacuo to yield 1-Boc-4-(3-amino-5-trifluoromethyl-
10 phenoxy)-piperidine as an off-white foam.

The following compounds were prepared similarly to the
procedure outlined above:

- 15 a) 1-Boc-2-(3-Amino-5-trifluoromethyl-phenoxy-methyl)-
pyrrolidine.
b) 2-(3-Amino-5-trifluoromethyl-phenoxy-methyl)-1-methyl-
pyrrolidine.
c) [2-(1-Methylpiperidin-4-yloxy)-pyridin-4-yl]methylaniline.
20 ESI (M+H)=222.
d) [2-(2-Morpholin-4-yl-ethoxy)-pyridin-4-yl]methylaniline.
e) [2-(2-Morpholin-4-yl-propoxy)-pyridin-4-yl]methylaniline.
f) [2-(1-Methyl-pyrrolidin-2-ylmethoxy)-pyridin-4-
yl]methylaniline. ESI MS: (M+H)=222.
25 g) (4-Aminomethyl-pyridin-2-yl)-(3-morpholin-4-yl-propyl)-
amine. ESI MS: (M+H)=251.
h) 4-tert-Butyl-3-(1-methyl-piperidin-4-ylmethoxy)-
phenylaniline.
i) 4-tert-Butyl-3-(2-piperidin-1-yl-ethoxy)-phenylaniline.
30 j) 3-(1-Methyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-
phenylaniline.
k) 3-(1-Isopropyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-
phenylaniline.
l) (S) 3-Oxiranylmethoxy-4-pentafluoroethyl-phenylaniline.

- 140 -

- m) 3-(2-Pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenylamine.
- n) 3-(2-Piperidin-1-yl-ethoxy)-4-trifluoromethyl-phenylamine.
- 5 o) (S) 3-(1-Boc-pyrrolidin-2-ylmethoxy)-4-pentafluoroethyl-phenylamine.
- p) (R) 3-(1-Boc-pyrrolidin-2-ylmethoxy)-4-pentafluoroethyl-phenylamine.
- 10 q) (R) 3-(1-Methyl-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenylamine.
- r) (S) 3-(1-Methyl-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenylamine
- s) (R) 3-Oxiranylmethoxy-4-pentafluoroethyl-phenylamine.
- t) (R) 2-(5-Amino-2-pentafluoroethyl-phenoxy)-1-pyrrolidin-15 1-yl-ethanol.
- u) 3-(1-Boc-azetidin-3-ylmethoxy)-4-pentafluoroethyl-phenylamine.
- v) 3-(2-(Boc-amino)ethoxy)-4-pentafluoroethyl-phenylamine.
- w) 6-Amino-2,2-dimethyl-4H-benzo[1,4]oxazin-3-one. M+H 20 193.2. Calc'd 192.1.
- x) 2,2,4-Trimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-ylamine.
- y) 1-(6-Amino-2,2-dimethyl-2,3-dihydro-benzo[1,4]oxazin-4-yl)-ethanone. M+H 221.4. Calc'd 220.3.
- 25 z) [2-(1-Benzhydryl-azetidin-3-yloxy)-pyridin-4-yl]-methylamine.
- aa) [2-(1-Methyl-piperidin-4-ylmethoxy)-pyridin-4-yl]-methylamine. M+H 236.3. Calc'd 235.2.
- ab) 3-(4-Boc-piperazin-1-ylmethyl)-5-trifluoromethyl-30 phenylamine. M+H 360.3.
- ac) 2-Boc-4,4-dimethyl-1,2,3,4-tetrahydro-isoquinolin-7-ylamine.
- ad) 3-Morpholin-4-ylmethyl-4-pentafluoroethyl-phenylamine.

- 141 -

- ae) 3-(4-Methyl-piperazin-1-ylmethyl)-4-pentafluoroethyl-phenylamine. M+H 410.3. Calc'd 409.4.
- af) 7-Amino-2-(4-methoxy-benzyl)-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one. M+H 311.1.
- 5 ag) 7-Amino-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one.
- ah) (3-Amino-5-trifluoromethyl-phenyl)-(4-Boc-piperazin-1-yl)-methanone. M+H 374.3; Calc'd 373.
- ai) 3-(4-Boc-Piperazin-1-ylmethyl)-5-trifluoromethyl-phenylamine.
- 10 aj) 1-(7-Amino-4,4-dimethyl-3,4-dihydro-1H-isoquinolin-2-yl)-ethanone. M+H 219.2.
- ak) {2-[2-(1-Methylpiperidin-4-yl)ethoxy]-pyridin-4-yl}-methylamine.
- al) {2-[2-(1-Pyrrolidinyl)ethoxy]-pyridin-4-yl}-methylamine.
- 15 am) {2-[2-(1-Methylpyrrolin-2-yl)ethoxy]-pyridin-4-yl}-methylamine.
- an) (2-Chloro-pyrimidin-4-yl)-methylamine.
- ao) 3-(1-Boc-azetidin-3-ylmethoxy)-5-trifluoromethyl-phenylamine.
- 20 ap) 4-tert-Butyl-3-(1-Boc-pyrrolidin-3-ylmethoxy)-phenylamine. M+H 385.
- aq) 4-tert-Butyl-3-(1-Boc-azetidin-3-ylmethoxy)-phenylamine. M+Na 357.
- ar) (S) 4-tert-Butyl-3-(1-Boc-pyrrolidin-2-ylmethoxy)-phenylamine. M+Na 371.
- 25 as) 3-tert-Butyl-4-(4-Boc-piperazin-1-yl)-phenylamine
- at) 3-(1-Methyl-piperidin-4-yl)-5-trifluoromethyl-phenylamine.
- au) 3,3-Dimethyl-2,3-dihydro-benzofuran-6-ylamine.
- 30 av) 3,9,9-Trimethyl-2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluoren-6-ylamine.
- aw) 4-[1-Methyl-1-(1-methyl-piperidin-4-yl)-ethyl]-phenylamine was prepared using EtOH as the solvent.

- 142 -

- ax) 4-tert-Butyl-3-(4-pyrrolidin-1-yl-but-1-enyl)-phenylamine.
- ay) (R) 3-(1-Boc-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenylamine.
- 5 az) (S) 3-(1-Boc-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenylamine.

Preparation IV - 1-Boc-4-{3-[(2-fluoro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy}-piperidine

- 10 1-Boc-4-(3-amino-5-trifluoromethyl-phenoxy)-piperidine (4.37 g) was dissolved in CH₂Cl₂ (100 ml) and NaHCO₃ (2.4 g, Baker) was added. 2-Fluoropyridine-3-carbonyl chloride (2.12 g) was added and the reaction was stirred at RT for 2.5 h. The reaction was filtered and concentrated in vacuo to yield a
- 15 yellow foam. (30%) EtOAc/Hexane was added and 1-Boc-4-{3-[(2-fluoro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy}-piperidine precipitated as an off white solid.

The following compounds were prepared similarly to the

20 procedure outlined above:

- a) 2-Fluoro-N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide.
- b) N-[4-tert-Butyl-3-(2-piperidin-1-yl-ethoxy)-phenyl]-2-fluoro-nicotinamide.
- 25 c) N-[3,3-Dimethyl-1-(1-methyl-piperidin-4-ylmethyl)-2,3-dihydro-1H-indol-6-yl]-2-fluoro-nicotinamide.
- d) N-[1-(2-Dimethylamino-acetyl)-3,3-dimethyl-2,3-dihydro-1H-indol-6-yl]-2-fluoro-nicotinamide
- 30 e) N-[3,3-Dimethyl-1-(2-(Boc-amino)acetyl)-2,3-dihydro-1H-indol-6-yl]-2-fluoro-nicotinamide.
- f) N-(4-Acetyl-2,2-dimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-2-fluoro-nicotinamide. M+H 344.5. Calc'd 343.4.

- 143 -

- g) 2-Fluoro-N-(2,2,4-trimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-nicotinamide. M+H 316.2. Calc'd 315.1.
- h) N-(2,2-Dimethyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-2-fluoro-nicotinamide. M+H 316.1. Calc'd 315.10.
- 5 i) 2-Fluoro-N-[3-(4-methyl-piperazin-1-ylmethyl)-5-trifluoromethyl-phenyl]-nicotinamide. M+H 481. Calc'd 480.
- j) 2-Fluoro-N-(2-Boc-4,4-dimethyl-1,2,3,4-tetrahydro-isoquinolin-7-yl)-nicotinamide. M+H 400.
- 10 k) 2-Fluoro-N-[3-(4-methyl-piperazin-1-ylmethyl)-4-pentafluoroethyl-phenyl]-nicotinamide. M+H 447.0. Calc'd 446.
- l) 2-Fluoro-N-(3-morpholin-4-ylmethyl-4-pentafluoroethyl-phenyl)-nicotinamide.
- 15 m) 2-Fluoro-N-[4-iodophenyl]-nicotinamide.
- n) 2-Fluoro-N-(4,4-dimethyl-1-oxo-1,2,3,4-tetrahydro-isoquinolin-7-yl)-nicotinamide. M+H 314.0, Calc'd 311.
- o) 2-Fluoro-N-[3-(4-Boc-piperazine-1-carbonyl)-5-trifluoromethyl-phenyl]-nicotinamide. M+H 495.
- 20 p) 2-Fluoro-N-[3-(4-Boc-piperazin-1-ylmethyl)-5-trifluoromethyl-phenyl]-nicotinamide. M+H 483.3; Calc'd 482.
- q) N-(2-Acetyl-4,4-dimethyl-1,2,3,4-tetrahydro-isoquinolin-7-yl)-2-fluoro-nicotinamide. M+H 430.0.
- 25 r) N-[3,3-Dimethyl-1-(1-methyl-piperidin-4-yl)-2,3-dihydro-1H-indol-6-yl]-2-fluoro-nicotinamide. M+H 383.2; Calc'd 382.5.
- s) N-(4-tert-Butylphenyl)-2-fluoronicotinamide.
- 30 t) N-(4-Trifluoromethylphenyl)-2-fluoronicotinamide.
- u) 2-Fluoro-N-[3-(1-Boc-azetidin-3-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide. M-H 468.2; Calc'd 469.16.

- 144 -

- v) 2-Fluoro-N-[3-(1-Boc-azetidin-3-ylmethoxy)-4-tert-butyl-phenyl]-nicotinamide.
- w) (S) N-[4-tert-Butyl-3-(1-Boc-pyrrolidin-2-ylmethoxy)-phenyl]-2-fluoro-nicotinamide. M+Na 494.
- 5 x) N-[3-(1-Methyl-piperidin-4-yl)-5-trifluoromethyl-phenyl]-2-fluoro-nicotinamide was prepared with K_2CO_3 . instead of $NaHCO_3$.
- y) N-(3-Bromo-5-trifluoromethyl-phenyl)-2-fluoro-nicotinamide.
- 10 z) 2-Fluoro-N-(3,9,9-trimethyl-2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluoren-6-yl)-nicotinamide.
- aa) 2-Fluoro-N-{4-[1-methyl-1-(1-methyl-piperidin-4-yl)-ethyl]-phenyl}-nicotinamide
- ab) N-[3,3-Dimethyl-1-(1-Boc-piperidin-4-ylmethyl)-2,3-
- 15 dihydro-1H-indol-6-yl]-2-fluoro-nicotinamide.

Preparation V - 1-Boc-4-{3-[(2-chloro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy}-piperidine

- 1-Boc-4-{3-[(2-chloro-pyridine-3-carbonyl)-amino]-5-
- 20 trifluoromethyl-phenoxy}-piperidine was prepared from 1-Boc-4-(3-amino-5-trifluoromethyl-phenoxy)-piperidine and 2-chloropyridine-3-carbonyl chloride by a procedure similar to that described in the preparation of 1-Boc-4-{3-[(2-fluoro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy}-
- 25 piperidine.

The following compounds were prepared similarly to the procedure outlined above:

- 30 a) N-(4-tert-Butyl-3-nitro-phenyl)-2-chloro-nicotinamide.
- b) 2-Chloro-N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide.
- c) 2-Chloro-N-[3-(3-morpholin-4-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide.

- 145 -

- d) 2-Chloro-N-[3-(1-methylpiperidin-4-yl)-5-trifluoromethyl-phenyl]-nicotinamide.
- e) 2-Chloro-N-[3-(1-methyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide.
- 5 f) 2-Chloro-N-[3-(1-isopropyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide.
- g) (S) 2-Chloro-N-[4-(oxiranylmethoxy)-3-pentafluoroethyl-phenyl]-nicotinamide.
- h) 2-Chloro-N-[3-(2-pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenyl]-nicotinamide.
- 10 i) 2-Chloro-N-[3-(2-piperidin-1-yl-ethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide.
- j) (R) 2-Chloro-N-[3-(1-Boc-pyrrolidin-2-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide.
- 15 k) (S) 2-Chloro-N-[3-(1-Boc-pyrrolidin-2-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide.
- l) (R) 2-Chloro-N-[3-(1-methyl-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide.
- m) (S) 2-Chloro-N-[3-(1-methyl-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide.
- 20 n) (R) 2-Chloro-N-[4-(oxiranylmethoxy)-3-pentafluoroethyl-phenyl]-nicotinamide.
- o) (R) Acetic acid 2-(5-[(2-chloro-pyridine-3-carbonyl)-amino]-2-pentafluoroethyl-phenoxy)-1-pyrrolidin-1-yl-ethyl ester.
- 25 p) 2-Chloro-N-[3-(4-methyl-piperazin-1-ylmethyl)-5-trifluoromethyl-phenyl]-nicotinamide.
- q) 2-Chloro-N-[2-(4-methoxy-benzyl)-4,4-dimethyl-1-oxo-1,2,3,4-tetrahydro-isoquinolin-7-yl]-nicotinamide. M+H 450.2. Calc'd 449.
- 30 r) 2-Chloro-N-(4,4-dimethyl-1-oxo-1,2,3,4-tetrahydro-isoquinolin-7-yl)-nicotinamide. M+H 330.1, Calc'd 329.
- s) 2-Chloro-N-[3-(4-Boc-piperazin-1-ylmethyl)-5-trifluoromethyl-phenyl]-nicotinamide.

- 146 -

- t) 2-{3-[(2-Chloro-pyridine-3-carbonyl)-amino]-phenyl}-2-methyl-propionic acid methyl ester. M+H 405
- u) N-{4-tert-Butyl-3-[2-(1-Boc-piperidin-4-yl)-ethyl]-phenyl}-2-chloro-nicotinamide. M+Na 524. Calc'd 501.1.
- 5 v) N-[3,3-Dimethyl-1,1-dioxo-2,3-dihydro-1H-benzo[d]isothiazol-6-yl]-2-chloro-nicotinamide.
- w) N-[1,1,4,4-Tetramethyl-1,2,3,4-tetrahydro-naphth-6-yl]-2-chloro-nicotinamide.
- x) 2-Chloro-N-[3,3-dimethyl-2,3-dihydro-benzofuran-6-yl]-2-chloro-nicotinamide.
- 10 y) 2-Chloro-N-[3-(1-Boc-piperidin-4-yloxy)-5-trifluoromethyl-phenyl]-nicotinamide.
- z) 2-Chloro-N-[3-(1-methyl-piperidin-4-ylmethyl)-5-trifluoromethyl-phenyl]-nicotinamide.
- 15 aa) 2-Chloro-N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide.
- ab) N-[4-tert-Butyl-3-(4-pyrrolidin-1-yl-but-1-enyl)-phenyl]-2-chloro-nicotinamide.
- ac) (R) 2-Chloro-N-[3-(1-Boc-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide.
- 20 ad) (S) 2-Chloro-N-[3-(1-Boc-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide.

Preparation VI - 1-Boc-2-{3-[(2-fluoro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy-methyl}-pyrrolidine

25 1-Boc-2-{3-[(2-Fluoro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy-methyl}-pyrrolidine was prepared from 1-Boc-2-(3-amino-5-trifluoromethyl-phenoxy-methyl)-pyrrolidine by a procedure similar to that described in the

30 preparation of 1-Boc-4-{3-[(2-fluoro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy}-piperidine.

- 147 -

Preparation VII - 2-(3-nitro-5-trifluoromethyl-phenoxymethyl)-pyrrolidine

1-Boc-2-(3-nitro-5-trifluoromethyl-phenoxymethyl)-pyrrolidine (2.35 g) was dissolved in CH_2Cl_2 (60 ml) and TFA (20 ml) was added. After stirring for 1 h at RT, the mixture was concentrated in vacuo to yield 2-(3-nitro-5-trifluoromethyl-phenoxymethyl)-pyrrolidine as an oil that solidified upon standing. The material was used as is without further purification.

10

The following compounds were prepared similarly to the procedure outlined above:

- a) (4-Aminomethyl-pyrimidin-2-yl)-(3-morpholin-4-yl-propyl)-amine.
- b) (4-Aminomethyl-pyrimidin-2-yl)-[2-(1-methyl-pyrrolidin-2-yl)-ethyl]-amine.

15

Preparation VIII - 1-methyl-2-(3-nitro-5-trifluoromethyl-phenoxymethyl)-pyrrolidine

20

2-(3-Nitro-5-trifluoromethyl-phenoxymethyl)-pyrrolidine (6 mmol) was dissolved in CH_3CN (20 ml) and formaldehyde (2.4 ml, 37% aqueous) was added. NaBH_3CN (607 mg) was added, an exotherm was observed. The pH is monitored every 15 min and adjusted to ~7 with AcOH. After 45 min, the mixture was concentrated in vacuo and the residue is dissolved in EtOAc, washed with 6N NaOH, 1N NaOH, and 2N HCl (3x). The acid washings were combined, adjusted to ~pH 10 with solid Na_2CO_3 and extracted with EtOAc (2x). The EtOAc fractions were combined, dried with Na_2SO_4 , and purified with flash chromatography (SiO_2 , 95:5:0.5 CH_2Cl_2 :MeOH: NH_4OH) to afford 1-methyl-2-(3-nitro-5-trifluoromethyl-phenoxymethyl)-pyrrolidine.

30

- 148 -

The following compounds were prepared similarly to the procedure outlined above:

- a) 2-(1-Methylpiperidin-4-yl)-ethanol.
- 5 b) 2-{3-[(2-Fluoro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy-methyl}-1-methylpyrrolidine.

Preparation IX - 4-tert-butyl-3-nitro-phenylamine

A mixture of 1,3-dinitro-4-tert-butylbenzene (10.0 g) in H₂O
10 (56 ml) was heated to reflux. A mixture of Na₂S (21.42 g) and sulfur (2.85 g) in H₂O (34 ml) was added over 1 h via an addition funnel. The reaction maintained at reflux for 1.5 h then cooled to RT and extracted with EtOAc. The organic
15 extracts were combined and washed with H₂O, brine, dried over MgSO₄ and concentrated in vacuo to afford 4-tert-butyl-3-nitro-phenylamine which was used as is without further purification.

Preparation X - N-(3-bromo-5-trifluoromethyl-phenyl)-acetamide

3-Bromo-5-(trifluoromethyl)phenylamine (5 g, Alfa-Aesar) was dissolved in AcOH (140 ml) and Ac₂O (5.9 ml, Aldrich) was added. The reaction was stirred at RT overnight. The
20 mixture was added slowly to H₂O (~700 ml) forming a white precipitate. The solid was isolated by filtration, washed with H₂O and dried under vacuum to yield N-(3-bromo-5-trifluoromethyl-phenyl)-acetamide.

Preparation XI - N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide

30 Allylpiperidine (1.96 g, Lancaster) was degassed under vacuum, dissolved in 0.5 M 9-BBN in THF (31.2 ml, Aldrich), and heated to reflux for 1 h, then cooled to RT. PD(dppf)Cl₂/CH₂Cl₂ was added to a degassed mixture of N-(3-

- 149 -

bromo-5-trifluoromethyl-phenyl)-acetamide, K_2CO_3 (9.8 g) DMF (32.1 ml and H_2O (3 ml). The allyl piperidine solution was added heated to $60^\circ C$ for 3 h. After cooling to RT and reheating at $60^\circ C$ for 6 h, the mixture was cooled to RT and
5 poured into H_2O . The mixture was extracted with EtOAc (2x), and the EtOAc portion was washed with 2 N HCl (2x) and brine. The aqueous phases were combined and the pH was adjusted to ~11 with NaOH (15%) forming a cloudy suspension. The cloudy suspension was extracted with EtOAc (2x) and the
10 EtOAc portion was dried with Na_2SO_4 , filtered and concentrated in vacuo. The crude material was purified by flash chromatography (SiO_2 , 95:5:0.5 CH_2Cl_2 :MeOH: NH_4OH) to afford N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide as a brown oil that solidified under
15 vacuum.

The following compounds were prepared similarly to the procedure outlined above:

- 20 a) N-(3-Morpholin-4-ylpropyl-5-trifluoromethyl-phenyl)-acetamide from 4-allyl-morpholine.
b) N-(3-(1-methylpiperidin-4-ylmethyl-5-trifluoromethyl-phenyl)-acetamide from 1-Methyl-4-methylene-piperidine.

25 **Preparation XII - 3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenylamine**

N-[3-(3-Piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide (1.33 g) was dissolved in EtOH (40 ml) and 12 N HCl (40 ml) was added. After stirring overnight at $70^\circ C$ and
30 RT, the mixture was concentrated in vacuo, affording 3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenylamine as a brown oil.

- 150 -

The following compounds were prepared similarly to the procedure outlined above:

- a) 3,3-Dimethyl-6-nitro-2,3-dihydro-1H-indole. M+H 193.1;
5 Calc'd 192.2.
b) 3-(1-Methyl-piperidin-4-ylmethyl)-5-trifluoromethyl-phenylamine.
c) 3-Morpholin-4-ylmethyl-5-trifluoromethyl-phenylamine.
- 10 **Preparation XIII - 3,3-Dimethyl-6-nitro-1-piperidin-4-ylmethyl-2,3-dihydro-1H-indole**
3,3-Dimethyl-1-(1-Boc-piperidin-4-ylmethyl)-6-nitro-2,3-dihydro-1H-indole was dissolved in HCl/EtOAc and stirred for 2 h. The mixture was concentrated in vacuo and partitioned
15 between 1,2-dichloroethane and 1N NaOH. The organic layer was removed, washed with brine, dried (Na₂SO₄) and filtered. The material was used without further purification.
- Preparation XIV - N-[3-(3-morpholin-4-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide**
20 N-[3-(3-Morpholin-4-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide was prepared from allyl morpholine and N-(3-bromo-5-trifluoromethyl-phenyl)-acetamide similar to that described in the preparation of N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide.
25
- Preparation XV - 3-(3-morpholin-4-yl-propyl)-5-trifluoromethyl-phenylamine**
3-(3-Morpholin-4-yl-propyl)-5-trifluoromethyl-phenylamine
30 was prepared from N-[3-(3-morpholin-4-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide similar to that described in the preparation of 3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenylamine.

- 151 -

Preparation XVI - 1-methyl-4-methylene-piperidine

Ph₃PCH₃I (50 g, Aldrich) was suspended in Et₂O (20 ml) and butyllithium (77.3 ml, 1.6 M in hexanes, Aldrich) was added dropwise. The reaction was stirred for 2 h at RT then 1-methylpiperidone (12.3 ml, Aldrich) was added slowly. The mixture was stirred at RT overnight. The solid was removed by filtration, the volume was reduced to ~400 ml and additional solid was removed by filtration. The Et₂O was washed with H₂O (2x) and 2N HCl (4x). The pH of the acid washings was adjusted to ~11 with 6 N NaOH, then they were extracted with CH₂Cl₂ (4x). The CH₂Cl₂ washings were dried over Na₂SO₄ and concentrated cold in vacuo to provide 1-methyl-4-methylene-piperidine which was used as is.

15 Preparation XVII - N-[3-(1-methylpiperidin-4-yl)-5-trifluoromethyl-phenyl]-acetamide

N-[3-(1-Methylpiperidin-4-yl)-5-trifluoromethyl-phenyl]-acetamide was prepared from 1-methyl-4-methylene-piperidine and N-(3-bromo-5-trifluoromethyl-phenyl)-acetamide similar to that described in the preparation of N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-acetamide.

Preparation XVIII - 3-(1-methylpiperidin-4-yl)-5-trifluoromethyl-phenylamine

25 3-(1-Methylpiperidin-4-yl)-5-trifluoromethyl-phenylamine was prepared from N-[3-(1-methylpiperidin-4-yl)-5-trifluoromethyl-phenyl]-acetamide similar to the procedure described in the preparation of 3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenylamine.

30

Preparation XIX - 2-(1-methylpiperidin-4-yloxy)-4-pyridylcarbonitrile

4-Hydroxy-1-methylpiperidine (25.4 g) was dissolved in THF (50 ml) in a 100 mL r.b. flask. NaH/mineral oil mixture

- 152 -

(9.58 g) was slowly added to the flask and stirred for 20 min. 2-Chloro-4-cyanopyridine was added to the mixture and stirred at RT until completion. Diluted mixture with EtOAc and added H₂O to quench mixture, then transferred contents to a sep. funnel. The organic phase was collected while the aqueous phase was washed two times with EtOAc. The combined organics were dried over Na₂SO₄, filtered, then concentrated *in vacuo*. Then redissolved mixture in CH₂Cl₂, 10% HCl (300 ml) was added and the mixture was transferred to sep. funnel. The org. was extracted, while EtOAc along with 300 mL 5N NaOH was added to the sep. funnel. The organic phases were collected, dried over Na₂SO₄, filtered and concentrated *in vacuo* affording 2-(1-methylpiperidin-4-yloxy)-4-pyridylcarbonitrile as a brown solid. ESI (M+H) = 218.

15

The following compounds were prepared similarly to the procedure outlined above:

- a) 2-(1-methylpiperidin-4-ylmethoxy)-4-pyridylcarbonitrile.
20 M+H 232.1. Calc'd 231.1.
- b) 2-(1-Benzhydryl-azetidin-3-yloxy)-4-pyridylcarbonitrile.
M+H 342.2. Calc'd 341.2.
- c) 2-(1-methylpiperidin-4-ylethoxy)-4-pyridylcarbonitrile.
- d) 2-(1-pyrrolidinylethoxy)-4-pyridylcarbonitrile.
- 25 e) 2-(1-methylpyrrolin-2-ylethoxy)-4-pyridylcarbonitrile.
- f) 2-[2-(1-Boc-azetidin-3-yl)-ethoxy]-4-pyridylcarbonitrile.

Preparation XX - [2-(1-methylpiperidin-4-yloxy)-pyridin-4-yl]methylaniline bis hydrochloride

- 30 [2-(1-Methylpiperidin-4-yloxy)-pyridin-4-yl]methylaniline was diluted with Et₂O (50 ml) and 1M HCl/Et₂O (47 ml) was added. The vessel was swirled until precipitate formed.

- 153 -

Preparation XXI - 2-(2-morpholin-4-yl-ethoxy)-4-pyridylcarbonitrile

2-(2-Morpholin-4-yl-ethoxy)-4-pyridylcarbonitrile was prepared from 2-chloro-4-cyanopyridine and 2-morpholin-4-yl-ethanol by a procedure similar to that described in the preparation of 2-(1-methylpiperidin-4-yloxy)-4-pyridylcarbonitrile. The hydrochloride salt was prepared similar to that described for [2-(1-methylpiperidin-4-yloxy)-pyridin-4-yl]methylaniline bis hydrochloride.

10

Preparation XXII - 2-morpholin-4-yl-propanol

LAH powder (1.6 g) was added to a flask while under N₂ atmosphere, immediately followed by THF (50 ml). The mixture was chilled to 0°C, methyl 2-morpholin-4-yl-propionate (5 g) was added dropwise to the reaction mixture and stirred at 0°C. After 1 h, the mixture was worked up by adding H₂O (44 mL), 2N NaOH (44 mL), then H₂O (44 mL, 3x). After 30 min of stirring, the mixture was filtered through Celite® and the organic portion was concentrated in vacuo providing 2-morpholin-4-yl-propanol as a colorless oil.

15
20

The following compounds were prepared similarly to the procedure outlined above:

- 25 a) (1-Methyl-piperidin-4-yl)-methanol. M+H 130.2. Calc'd 129.1.

Preparation XXIII - 2-(2-morpholin-4-yl-propoxy)-4-pyridylcarbonitrile

30 2-(2-Morpholin-4-yl-propoxy)-4-pyridylcarbonitrile was prepared from 2-chloro-4-cyanopyridine and 2-morpholin-4-yl-propanol by a procedure similar to that described in the preparation of 2-(1-methylpiperidin-4-yloxy)-4-pyridylcarbonitrile.

- 154 -

Preparation XXIV - 2-(1-Methyl-pyrrolidin-2-ylmethoxy)-4-pyridylcarbonitrile

2-(1-Methyl-pyrrolidin-2-ylmethoxy)-4-pyridylcarbonitrile
5 was prepared from 2-chloro-4-cyanopyridine and 1-methyl-pyrrolidin-2-ylmethanol by a procedure similar to that described in the preparation of 2-(1-methylpiperidin-4-yloxy)-4-pyridylcarbonitrile. ESI MS: (M+H)=218.

10 **Preparation XXV - 2-(3-morpholin-4-yl-propylamino)-4-pyridylcarbonitrile**

To a flask charged with 2-chloro-4-cyanopyridine (2.0 g),
was added the aminopropyl morpholine (2.11 ml). The mixture
was heated to 79°C for 5 h and stirred. After 5 h the
15 reaction was incomplete. The mixture was then heated at
60°C overnight. The crude compound was purified on silica
gel (1-5% MeOH/CH₂Cl₂ gradient). ESI MS: (M+H)=247, (M-
H)=245.

20 **Preparation XXVI - 5-Nitro-2-pentafluoroethylphenol**

Combined 2-methoxy-4-nitro-1-pentafluoroethylbenzene (9.35
g) and pyridine hydrochloride in a round bottom flask and
heated at 210°C for 1 h then cooled to RT. The mixture was
diluted with EtOAc and 2N HCl (>500 ml) until all residue
25 dissolved. The organic layer was removed, washed with 2N
HCl (2x) and concentrated *in vacuo*. The residue was
dissolved in hexanes and Et₂O, washed with 2N HCl, then
brine. Dried organic layer over Na₂SO₄, filtered,
concentrated *in vacuo* and dried under high vacuum to provide
30 5-nitro-2-pentafluoromethylphenol.

Preparation XXVII - 2-tert-Butyl-5-nitro-aniline

To H₂SO₄ (98%, 389 mL) in a 500 mL 3-neck flask was added 2-
tert-butyl aniline (40.6 mL). The reaction was cooled to -

- 155 -

10°C and KNO₃ in 3.89 g aliquots was added every 6 min for a total of 10 aliquots. Tried to maintain temperature at -5°C to -10°C. After final addition of KNO₃, stirred the reaction for five min then it was poured onto ice (50 g).

- 5 The black mix was diluted with H₂O and extracted with EtOAc. The aqueous layer was basified with solid NaOH slowly then extracted with EtOAc (2x). The combined organic layers were washed with 6N NaOH and then with a mix of 6N NaOH and brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*
- 10 to obtain crude 2-tert-butyl-5-nitro-aniline as a dark red-black oil which solidified when standing at RT. The crude material was triturated with about 130 mL hexanes. After decanting the hexanes, the material was dried to obtain a dark-red black solid.

15

Preparation XXVIII - 2-tert-Butyl-5-nitrophenol

- In a 250 ml round bottom flask, 20 mL concentrated H₂SO₄ was added to 2-tert-butyl-5-nitro-aniline (7.15 g) by adding 5 mL aliquots of acid and sonicating with occasional heating
- 20 until all of the starting aniline went into solution. H₂O (84 ml) was added with stirring, then the reaction was cooled to 0°C forming a yellow-orange suspension. A solution of NaNO₂ (2.792 g) in H₂O (11.2 mL) was added dropwise to the suspension and stirred for 5 min. Excess
- 25 NaNO₂ was neutralized with urea, then the cloudy solution was transferred to 500 ml 3-necked round bottom flask then added 17 mL of 1:2 H₂SO₄:H₂O solution, and heated at reflux. Two additional 5 mL aliquots of 1:2 H₂SO₄:H₂O solution, a 7 mL aliquot of 1:2 H₂SO₄:H₂O solution and another 10 mL of 1:2
- 30 H₂SO₄: H₂O were added while heating at reflux. The mixture was cooled to RT forming a black layer floating on top of the aqueous layer. The black layer was diluted with EtOAc (300 mL) and separated. The organic layer was washed with H₂O then brine, dried over Na₂SO₄ and concentrated *in vacuo*.

- 156 -

Crude oil was purified on silica gel column with 8% EtOAc/Hexanes. Upon drying under vacuum, the 2-tert-butyl-5-nitrophenol was isolated as a brown solid.

5 **Preparation XXIX - 1-methylpiperidine-4-carboxylic acid ethyl ester**

Piperidine-4-carboxylic acid ethyl ester (78 g) was dissolved in MeOH (1.2 L) at RT then formaldehyde (37%, 90 ml) and acetic acid (42 ml) were added and stirred for 2 h.
10 The mixture was cooled to 0°C, NaCNBH₃ (70 g) was added, and the mix was stirred for 20 min at 0°C, then overnight at RT. The mixture was cooled to 0°C then quenched with 6N NaOH. The mixture was concentrated *in vacuo* to an aqueous layer, which was extracted with EtOAc (4x), brine-washed, dried
15 over Na₂SO₄, and concentrated *in vacuo* to provide 1-methylpiperidine-4-carboxylic acid ethyl ester.

The following compounds were prepared similarly to the procedure outlined above:

20

a) (1-Methyl-piperidin-4-yl)-methanol. M+H 130.2. Calc'd 129.1.

25 **Preparation XXX - N-[4-tert-Butyl-3-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-2-chloro-nicotinamide**

N-[4-tert-Butyl-3-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-2-chloro-nicotinamide was prepared from 4-tert-butyl-3-(1-methyl-piperidin-4-ylmethoxy)-phenylamine by a procedure similar to that described in the preparation of 1-Boc-4-(3-
30 [(2-chloro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy)-piperidine.

- 157 -

Preparation XXXI - 1-[2-(2-tert-Butyl-5-nitro-phenoxy)-ethyl]-piperidine

To 2-tert-butyl-5-nitrophenol (1.01 g) and K_2CO_3 (1.72 g) was added acetone (35 ml) and H_2O (10.5 mL), then 1-(2-chloroethyl)piperidine HCl (1.909 g) and TBAI (153 mg). The mixture was stirred at reflux overnight. Additional K_2CO_3 (850 mg) and 1-(2-chloroethyl)-piperidine HCl (950 mg) were added and the mixture was heated at reflux for 6 h. The mixture was concentrated in vacuo to an aqueous layer which was acidified with 2N HCl and extracted with EtOAc. The aqueous layer was basified with 6N NaOH and washed with CH_2Cl_2 (3x). The combined organic layers were washed with brine/1N NaOH and dried over Na_2SO_4 . Washed the EtOAc layer with 2N NaOH/brine and dried over Na_2SO_4 . The crude material was purified by silica gel column chromatography with 15% EtOAc/Hexanes to yield 1-[2-(2-tert-butyl-5-nitro-phenoxy)-ethyl]-piperidine as a light tan solid.

(M+1)=307.3.

20 Preparation XXXII - 1-Boc-Piperidine-4-carboxylic acid ethyl ester

To a stirred solution of piperidine-4-carboxylic acid ethyl ester (23.5 g) in EtOAc (118 ml) at 0°C was added dropwise Boc_2O in EtOAc (60 ml). The reaction was warmed to RT and stirred overnight. Washed reaction with H_2O , 0.1N HCl, H_2O , $NaHCO_3$ and brine. The organic layer was dried over Na_2SO_4 , filtered and concentrated in vacuo. The liquid was dried under vacuum to provide 1-Boc-piperidine-4-carboxylic acid ethyl ester.

30

The following compounds were prepared similarly to the procedure outlined above:

a) N-Boc-(2-chloropyrimidin-4-yl)-methylaniline.

- 158 -

- b) 1-(2-tert-Butyl-4-nitrophenyl)-4-Boc-piperazine.
- c) 1-Boc-azetidine-3-carboxylic acid
- d) 1-Boc-4-Hydroxymethyl-piperidine using TEA.

5 **Preparation XXXIII - 1-Boc-4-hydroxymethyl-piperidine**

1-Boc-4-Hydroxymethyl-piperidine was prepared from 1-Boc-piperidine-4-carboxylic acid ethyl ester by a procedure similar to that described in the preparation of 2-morpholin-4-yl-propanol.

10

Preparation XXXIV - 1-Boc-4-Methylsulfonyloxymethyl-piperidine

Dissolved 1-Boc-4-hydroxymethyl-piperidine in anhydrous CH_2Cl_2 (50 ml) and TEA (4.5 ml) and cooled to 0°C . Mesyl chloride (840 μl) was added and the mixture was stirred for 15 min then at RT for 45 min. The mixture was washed with brine/1N HCl and then brine, dried over Na_2SO_4 , concentrated in vacuo and dried under high vacuum to provide 1-Boc-4-methylsulfonyloxymethyl-piperidine as a yellow orange thick oil.

20

The following compounds were prepared similarly to the procedure outlined above:

- 25 a) 1-Boc-3-methylsulfonyloxymethyl-azetidine.

Preparation XXXV - 1-Boc-4-(3-nitro-6-pentafluoroethyl-phenoxy)methyl-piperidine

To a slurry of 60% NaH suspension in DMF (30 mL) at RT added a solution of 5-nitro-2-pentafluoroethyl-phenol (3.6 g) in 5 mL DMF. The dark red mixture was stirred at RT for 10 min then added a solution of 1-Boc-4-methylsulfonyloxymethyl-piperidine (3.1 g) in 5 mL DMF. The reaction was stirred at 60°C and 95°C . After 1h, added 2.94 g K_2CO_3 and stirred

30

- 159 -

overnight at 105°C . After cooling to RT, the reaction was diluted with hexanes and 1N NaOH. Separated layers, and washed organic layer with 1N NaOH and with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. Purification
5 with silica gel column chromatography with 8% EtOAc/Hexanes yielded 1-Boc-4-(3-nitro-6-pentafluoroethyl-phenoxyethyl)-piperidine as a light yellow thick oil.

10 **Preparation XXXVI - 4-(3-nitro-6-pentafluoroethyl-phenoxyethyl)-piperidine**

4-(3-Nitro-6-pentafluoroethyl-phenoxyethyl)-piperidine was prepared from 1-Boc-4-(3-nitro-6-pentafluoroethyl-phenoxyethyl)-piperidine by a procedure similar to that described in the preparation of 2-(3-nitro-5-
15 trifluoromethyl-phenoxyethyl)-pyrrolidine.

Preparation XXXVII - 1-methyl-4-(3-nitro-6-pentafluoroethyl-phenoxyethyl)-piperidine

4-(3-Nitro-6-pentafluoroethyl-phenoxyethyl)-piperidine
20 (316.5 mg) was dissolved in 2.7 mL acetonitrile, then added 37% formaldehyde/H₂O (360 ul) and then NaBH₃CN (90 mg). Upon addition of NaCNBH₃ the reaction exothermed slightly. The reaction was stirred at RT and pH was maintained at ~7 by addition of drops of glacial acetic acid. After about 1
25 h, the mixture was concentrated in vacuo, treated with 8 mL 2N KOH and extracted two times with 10 mL Et₂O. The organic layers were washed with 0.5N KOH and then the combined organic layers were extracted two times with 1N HCl. The aqueous layer was basified with solid KOH and extracted two
30 times with Et₂O. This organic layer was then washed with brine/1N NaOH, dried over Na₂SO₄, filtered, concentrated in vacuo and dried under high vacuum to give pure compound.

- 160 -

Preparation XXXVIII - 1-Isopropyl-4-(5-nitro-2-pentafluoroethyl-phenoxyethyl)-piperidine

Dissolved 4-(5-nitro-2-pentafluoroethyl-phenoxyethyl)-piperidine (646 mg) in 1,2-dichloroethane (6.4 ml), then
5 added acetone (136 ul), NaBH(OAc)₃ (541 mg) and finally acetic acid (105 ul). Stirred the cloudy yellow solution under N₂ at RT overnight. Added another 130 uL acetone and stirred at RT over weekend. Quenched the reaction with 30 mL N NaOH/H₂O and stirred 10 min. Extracted with Et₂O and
10 the organic layer was brine-washed, dried over Na₂SO₄, filtered and concentrated in vacuo. Dried under high vacuum for several h to obtain 1-isopropyl-4-(5-nitro-2-pentafluoroethyl-phenoxyethyl)-piperidine as a yellow orange solid.

15

The following compounds were prepared similarly to the procedure outlined above:

- a) 3,3-Dimethyl-1-(1-methyl-piperidin-4-yl)-6-nitro-2,3-dihydro-1H-indole was prepared using 1-methyl-piperidin-
20 4-one. M+H 290; Calc'd 289.4.
- b) 3,3-Dimethyl-1-(1-Boc-piperidin-4-ylmethyl)-6-nitro-2,3-dihydro-1H-indole using 1-Boc-4-formyl-piperidine.

Preparation XXXIX - 3,3-Dimethyl-1-(1-methyl-piperidin-4-ylmethyl)-6-nitro-2,3-dihydro-1H-indole

25 3,3-Dimethyl-1-piperidin-4-ylmethyl-6-nitro-2,3-dihydro-1H-indole was treated with an excess of formaldehyde and NaBH(OAc)₃ and stirred overnight at RT. The reaction was quenched with MeOH and concentrated in vacuo. The residue
30 was partitioned between EtOAc and 1N NaOH. The organic layer was removed, washed with brine, dried (Na₂SO₄), filtered and concentrated to provide the compound.

- 161 -

Preparation XL - (S) 2-(5-Nitro-2-pentafluoroethyl-phenoxymethyl)-oxirane

Combined 5-nitro-2-pentafluoromethylphenol (2.69 g), DMF (25 ml) K_2CO_3 (3.03 g) and (S)-toluene-4-sulfonic acid oxiranylmethyl ester (2.27 g) and stirred the mixture at 90°C. After about 4 hours, the mix was cooled, diluted with EtOAc, washed with H_2O , 1N NaOH (2x), 1N HCl and then with brine. Dried over Na_2SO_4 , filtered and concentrated *in vacuo*. Purified the crude on silica gel column with 5% EtOAc/hexane and drying under high vacuum provided the (S)-2-(5-nitro-2-pentafluoroethyl-phenoxymethyl)-oxirane.

The following compounds were prepared similarly to the procedure outlined above:

a) (R)-2-(5-Nitro-2-pentafluoroethyl-phenoxymethyl)-oxirane.

Preparation XLI - (S) 2-Chloro-N-[3-(2-hydroxy-3-pyrrolidin-1-yl-propoxy)-4-pentafluoroethyl-phenyl]-nicotinamide

(S) 2-Chloro-N-[4-(2-oxiranylmethoxy)-3-pentafluoroethyl-phenyl]-nicotinamide (1.11 g) in a sealed tube and added pyrrolidine (285 μ l). Stirred after sealing tube at 60°C. After 12 h, the mix was concentrated *in vacuo* and purified on a silica gel column (5:95:0.5 MeOH: CH_2Cl_2 : NH_4OH - 8:92:1, MeOH: CH_2Cl_2 : NH_4OH). Concentrated *in vacuo* and dried under high vacuum to obtain pure compound.

The following compounds were prepared similarly to the procedure outlined above:

a) (R) 1-(5-Nitro-2-pentafluoroethyl-phenoxy)-3-pyrrolidin-1-yl-propan-2-ol.

- 162 -

Preparation XLII - 5-nitro-2-trifluoromethylanisole

Cooled 140 mL pyridine in a large sealable vessel to -40°C. Bubbled in trifluoromethyl iodide from a gas cylinder which had been kept in freezer overnight. After adding ICF₃ for 5 20 min, added 2-iodo-5-nitroanisole (24.63 g) and copper powder (67.25 g). Sealed vessel and stirred vigorously for 22 h at 140°C. After cooling to -50°C, carefully unsealed reaction vessel and poured onto ice and Et₂O. Repeatedly washed with Et₂O and H₂O. Allowed the ice - Et₂O mixture to 10 warm to RT. Separated layers, washed organic layer with 1N HCl (3x), then brine, dried over Na₂SO₄, filtered and concentrated in vacuo. Eluted material through silica gel plug (4.5:1 Hex:CH₂Cl₂) to provide 5-nitro-2-trifluoromethylanisole.

15

Preparation XLIII - 1-[2-(5-nitro-2-trifluoromethylphenoxy)ethyl]pyrrolidine

1-[2-(5-Nitro-2-trifluoromethylphenoxy)ethyl]-pyrrolidine was prepared from 5-nitro-2-trifluoromethyl-phenol and 1-(2-chloroethyl)pyrrolidine by a procedure similar to that 20 described for 1-[2-(2-tert-butyl-5-nitro-phenoxy)-ethyl]-piperidine.

Preparation XLIV - 1-[2-(5-Nitro-2-pentafluoroethyl-phenoxy)-ethyl]-piperidine

1-[2-(5-Nitro-2-pentafluoroethyl-phenoxy)-ethyl]-piperidine was prepared from 5-nitro-2-pentafluoroethylphenol and 1-(2-chloroethyl)piperidine by a procedure similar to that described in the preparation of 1-[2-(2-tert-butyl-5-nitro- 30 phenoxy)-ethyl]-piperidine.

Preparation XLV - 3-(1-Boc-pyrrolidin-2-ylmethoxy)-4-pentafluoroethyl-phenylamine

- 163 -

3-(2-Pyrrolidin-1-yl-methoxy)-4-trifluoromethyl-phenylamine was prepared from 1-[2-(5-nitro-2-trifluoromethylphenoxy)methyl]-pyrrolidine by a procedure similar to that described in the preparation of 1-Boc-4-(3-amino-5-trifluoromethyl-phenoxy)-piperidine.

Preparation XLVI - 2-Chloro-N-[3-(2-pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenyl]-nicotinamide

2-Chloro-N-[3-(2-pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenyl]-nicotinamide was prepared from 3-(2-pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenylamine and 2-chloropyridine-3-carbonyl chloride by a procedure similar to that described in the preparation of 1-Boc-4-{3-[(2-chloro-pyridine-3-carbonyl)-amino]-5-trifluoromethyl-phenoxy}-piperidine.

Preparation XLVII - (R) Acetic acid 2-(5-nitro-2-pentafluoroethyl-phenoxy)-1-pyrrolidin-1-ylmethyl-ethyl ester

Dissolved 1-(5-nitro-2-pentafluoroethyl-phenoxy)-3-pyrrolidin-1-yl-propan-2-ol (3.5 g) in CH_2Cl_2 (15 ml), added TEA (2.55 ml) and cooled to 0°C. Acetyl chloride (781.3 μl) was added dropwise, forming a suspension. The mixture was warmed to RT and stirred for 1.5 h. Additional acetyl chloride (200 μl) was added and the mix was stirred for another h. The mixture was diluted with CH_2Cl_2 and washed with sat. NaHCO_3 . The organic layer was removed, washed with brine and back extracted with CH_2Cl_2 . Dried the combined organic layers over Na_2SO_4 , filtered and concentrated in vacuo. The residue was purified over silica gel column (5:94.5:0.5 MeOH: CH_2Cl_2 : NH_4OH) to provide acetic acid 2-(5-nitro-2-pentafluoroethyl-phenoxy)-1-pyrrolidin-1-ylmethyl-ethyl ester as a yellow brown oil.

- 164 -

The following compounds were prepared similarly to the procedure outlined above:

- 5 a) (R) Acetic acid 2-(5-amino-2-pentafluoroethyl-phenoxy)-1-pyrrolidin-1-yl-methyl-ethyl ester.
- b) 1-(2,2-Dimethyl-6-nitro-2,3-dihydro-benzo[1,4]oxazin-4-yl)-ethanone. M-NO₂ 206.4; Calc'd 250.1.

10 **Preparation XLVIII - (R) 2-Chloro-N-[3-(2-hydroxy-2-pyrrolidin-1-yl-propoxy)-4-pentafluoroethyl-phenyl]-nicotinamide**

(R) Acetic acid 2-{5-[(2-chloro-pyridine-3-carbonyl)-amino]-2-pentafluoroethyl-phenoxy}-1-pyrrolidin-1-yl-ethyl ester (408 mg) was dissolved in MeOH (15 ml) and NH₄OH (6 ml) was added and the mixture was stirred at RT for 6 h. The reaction was concentrated in vacuo and dried under high vacuum. The residue was purified over silica gel column (8:92:0.6 MeOH: CH₂Cl₂:NH₄OH). The purified fractions were concentrated in vacuo and dried again to provide (R)-2-chloro-N-[3-(2-hydroxy-2-pyrrolidin-1-yl-ethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide as a white foam.

25 **Preparation XLIX - 2-Dimethylamino-1-(3,3-dimethyl-6-nitro-2,3-dihydro-indol-1-yl)-ethanone**

3,3-Dimethyl-6-nitro-2,3-dihydro-1H-indole (5 g) was dissolved in DMF (100 ml) and HOAt (3.89 g) dimethylamino-acetic acid (5.83 g) and EDC (3.89 g) were added. The reaction was stirred overnight. The mixture was diluted with CH₂Cl₂ (1L) and washed with sat'd NaHCO₃ (3x200 ml). The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by flash chromatography (SiO₂, EtOAc to 5%MeOH/EtOAc) to afford the title compound.

- 165 -

The following compounds were prepared similarly to the procedure outlined above:

- 5 a) 1-(3,3-Dimethyl-6-nitro-2,3-dihydro-indol-1-yl)-2-(N-Boc-amino)-ethanone.

Preparation L - 1-(6-Amino-3,3-dimethyl-2,3-dihydro-indol-1-yl)-2-(N-Boc-amino)-ethanone

- 10 1-(3,3-Dimethyl-6-nitro-2,3-dihydro-indol-1-yl)-2-(N-Boc-amino)-ethanone (3.9 g) was dissolved in EtOH (30 ml) and Fe powder (3.1 g) NH_4Cl (299 mg) and H_2O (5 ml) were added. The reaction was stirred at 80°C overnight. The reaction was filtered through Celite® and evaporated off the MeOH. The residue was partitioned between CH_2Cl_2 and sat'd NaHCO_3 . The organic layer was removed, washed with brine, dried over Na_2SO_4 , filtered and concentrated in vacuo. The residue was purified by flash chromatography (SiO_2 , 25% EtOAc/hexane). The purified fractions were concentrated in vacuo to afford the compound as a white powder.

20

The following compounds were prepared similarly to the procedure outlined above:

- 25 a) 1-(6-Amino-3,3-dimethyl-2,3-dihydro-indol-1-yl)-2-dimethylamino-ethanone.
- b) 3,3-Dimethyl-1-(1-methyl-piperidin-4-ylmethyl)-2,3-dihydro-1H-indol-6-ylamine.
- c) 3-(4-Methyl-piperazin-1-ylmethyl)-4-pentafluoroethyl-phenylamine. M+H 324.2. Calc'd 323.
- 30 d) 3,3-Dimethyl-1-(1-methyl-piperidin-4-yl)-2,3-dihydro-1H-indol-6-ylamine. M+H 259.6; Calc'd 259.3.
- e) 3,3-Dimethyl-1,1-dioxo-2,3-dihydro-1H-116-benzo[d]isothiazol-6-ylamine
- f) 1,1,4,4-Tetramethyl-1,2,3,4-tetrahydro-naphth-6-ylamine.

- 166 -

g) 3,3-Dimethyl-1-(1-Boc-piperidin-4-ylmethyl)-2,3-dihydro-1H-indol-6-ylamine.

Preparation LI - 2-Boc-4,4-dimethyl-7-nitro-1,2,3,4-

5 tetrahydro-isoquinoline

4,4-Dimethyl-7-nitro-1,2,3,4-tetrahydro-isoquinoline (150 mg) was dissolved with CH₂Cl₂ (3 ml) DIEA (100 ul) DMAP (208 mg and Boc₂O (204 mg) and the mixture was stirred for 6 h at RT. The reaction was diluted with CH₂Cl₂, washed with sat'd
10 NaHCO₃ and dried over MgSO₄, filtered and concentrated to provide the compound which was used without further purification.

The following compounds were prepared similarly to the
15 procedure outlined above substituting Ac₂O:

a) 1-(4,4-Dimethyl-7-nitro-3,4-dihydro-1H-isoquinolin-2-yl)-ethanone. M+H 249.3.

20 Preparation LII - 2-Bromo-N-(4-methoxy-benzyl)-5-nitro-benzamide

PMB-amine (5.35 ml) in CH₂Cl₂ (130 ml) was slowly added to 2-bromo-5-nitro-benzoyl chloride (10.55 g) and NaHCO₃ (9.6 g) and the mixture was stirred at RT for 1 h. The mixture
25 was diluted with CH₂Cl₂ (1 L), filtered, washed with dilute HCl, dried, filtered again, concentrated and dried under vacuum to provide the compound as a white solid. M+H 367. Calc'd 366.

30 Preparation LIII - 2-Bromo-N-(4-methoxy-benzyl)-N-(2-methyl-allyl)-5-nitro-benzamide

To a suspension of NaH (1.22 g) in DMF (130 ml) was added 2-bromo-N-(4-methoxy-benzyl)-5-nitro-benzamide (6.2 g) in DMF (60 ml) at -78C. The mixture was warmed to 0°C, 3-bromo-2-

- 167 -

5 methyl-propene (4.57 g) was added and the mixture was stirred for 2 h at 0°C. The reaction was poured into ice water, extracted with EtOAc (2x400 ml), dried over MgSO₄, filtered and concentrated to a DMF solution which was used without further purification.

Preparation LIV - of 2-(4-Methoxy-benzyl)-4,4-dimethyl-7-nitro-3,4-dihydro-2H-isoquinolin-1-one

10 2-Bromo-N-(4-methoxy-benzyl)-N-(2-methyl-allyl)-5-nitro-benzamide (23.4 mmol) was dissolved in DMF (150 ml) and Et₃NCl (4.25 g), HCO₂Na (1.75 g) and NaOAc (4.99 g) were added. N₂ was bubbled through the solution for 10 min, then Pd(OAc)₂ (490 mg) was added and the mixture was stirred overnight at 70°C. The mixture was extracted with EtOAc, 15 washed with sat'd NH₄Cl, dried over MgSO₄, filtered and concentrated until the compound precipitated as a white solid.

20 The following compounds were prepared similarly to the procedure outlined above:

- a) 3,3-Dimethyl-6-nitro-2,3-dihydro-benzofuran was prepared from 1-bromo-2-(2-methyl-allyloxy)-4-nitro-benzene.
- b) 3,9,9-Trimethyl-6-nitro-4,9-dihydro-3H-3-aza-fluorene was 25 prepared from 4-[1-(2-bromo-4-nitro-phenyl)-1-methyl-ethyl]-1-methyl-1,2,3,6-tetrahydro-pyridine.

Preparation LV - 4,4-Dimethyl-7-nitro-3,4-dihydro-2H-isoquinolin-1-one

30 2-(4-Methoxy-benzyl)-4,4-dimethyl-7-nitro-3,4-dihydro-2H-isoquinolin-1-one (2.0 g) was dissolved in CH₃CN (100 ml) and H₂O (50 ml) and cooled to 0°C. CAN (9.64 g) was added and the reaction was stirred at 0°C for 30 min, then warmed to RT and stirred for 6 h. The mixture was extracted with

- 168 -

CH₂Cl₂ (2x300 ml) washed with sat'd NH₄Cl, dried over MgSO₄, filtered and concentrated. The crude material was recrystallized in CH₂Cl₂/EtOAc (1:1) to give 4,4-dimethyl-7-nitro-3,4-dihydro-2H-isoquinolin-1-one as a white solid.

5

Preparation LVI - 4,4-Dimethyl-7-nitro-1,2,3,4-tetrahydro-isoquinoline

4,4-Dimethyl-7-nitro-3,4-dihydro-2H-isoquinolin-1-one (230 mg) was dissolved in THF (10 ml) and BH₃Me₂S (400 ul) was added and the reaction was stirred overnight at RT. The reaction was quenched with MeOH (10 ml) and NaOH (200 mg) and heating at reflux for 20 min. The mixture was extracted with EtOAc, washed with sat'd NH₄Cl, extracted with 10% HCl (20 ml). The acidic solution was treated with 5N NaOH (15 ml), extracted with EtOAc (30 ml) dried, filtered and evaporated to give the compound as a yellow solid. M+H 207.2, Calc'd 206.

10
15
20

The following compounds were prepared similarly to the procedure outlined above:

a) 4-Boc-2,2-dimethyl-6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine.

25 Preparation LVII - 2-Bromomethyl-4-nitro-1-pentafluoroethyl-benzene

2-Methyl-4-nitro-1-pentafluoroethyl-benzene (2.55 g) was dissolved in CCl₄ (30 ml) and AIBN (164 mg) and NBS (1.96 g) were added. The reaction was heated to reflux and stirred for 24 h. The mix was diluted with CH₂Cl₂, washed with sat'd NaHCO₃, dried over MgSO₄ and concentrated to give the compound as an oil which was used without further purification.

30

- 169 -

Preparation LVIII - 1-Methyl-4-(5-nitro-2-pentafluoroethyl-benzyl)-piperazine

2-Bromomethyl-4-nitro-1-pentafluoroethyl-benzene (2.6 g) was added to N-methylpiperazine (5 ml) and stirred at RT for 3 h. The mixture was filtered and the filtrate was treated with 1-chlorobutane, extracted with 2N HCl (100 ml). The acidic solution was treated with 5N NaOH (6 ml) then extracted with EtOAc. The organic layer was removed, dried over MgSO₄ and concentrated to give the compound as an oil.

10

The following compounds were prepared similarly to the procedure outlined above:

a) 4-(5-Nitro-2-pentafluoroethyl-benzyl)-morpholine.

15

Preparation LIX - 1-Boc-4-(5-nitro-2-pentafluoroethyl-benzyl)-piperazine.

2-Bromomethyl-4-nitro-1-pentafluoroethyl-benzene (2.5 g) was dissolved in CH₂Cl₂ and added to N-Boc-piperazine (2.5 g) and NaHCO₃ (1 g) and stirred at RT overnight. The mixture was diluted with CH₂Cl₂ (100 ml), washed with sat'd NH₄Cl, dried over MgSO₄, filtered and concentrated. The residue was purified by silica gel chromatography (hexane, CH₂Cl₂:hexane 2:8) to give the compound as an yellow solid.

25

Preparation LX - (4-Boc-piperazin-1-yl)-(3-nitro-5-trifluoromethyl-phenyl)-methanone

A mixture of 3-nitro-5-trifluoromethyl-benzoic acid (4.13 g), 4-Boc-piperazine (2.97 g), EDC (3.88 g), HOBt (2.74 g), DIEA (3.33 ml) in CH₂Cl₂ (120 ml) was stirred at RT for 3 h. The mixture was diluted with CH₂Cl₂ (100 ml), washed with sat'd NH₄Cl, dried over MgSO₄, filtered and concentrated. The residue was purified by silica gel chromatography

30

- 170 -

(hexane, CH₂Cl₂:hexane 1:2) to give the compound as a white solid.

Preparation LXI - 1-Boc-4-(3-nitro-5-trifluoromethyl-benzyl)-piperazine

(4-Boc-piperazin-1-yl)-(3-nitro-5-trifluoromethyl-phenyl)-methanone (403 mg) was dissolved in THF (6 ml) and BH₃Me₂S (300 µl) was added and the reaction was stirred for 3 h at 60°C and 2 h at RT. The reaction was quenched with MeOH (5 ml) and NaOH (100 mg) and stirred at RT for 1 h. The mixture was concentrated and dissolved in CH₂Cl₂, washed with sat'd NH₄Cl/NaHCO₃, dried (MgSO₄), filtered and evaporated to give the compound as an oil. M+H 390.3.

Preparation LXII - 2-Ethyl-4-aminomethyl pyridine

To a solution of 2-ethyl-4-thiopyridylamide (10 g) in MeOH (250 ml) was added Raney 2800 Nickel (5 g, Aldrich) in one portion. The mixture was stirred at RT for 2 days then at 60°C for 16 h. The mixture was filtered, concentrated to provide the desired compound.

Preparation LXIII - N-Boc-[2-(4-morpholin-4-yl-butyl)-pyrimidin-4-ylmethyl]-amine

N-Boc-(2-chloropyrimidine)-methylamine (663 mg) and 4-(aminopropyl)morpholine (786 mg) were dissolved in MeOH and concentrated *in vacuo*. The residue was heated at 100°C for 15 min, forming a solid which was dissolved in CH₂Cl₂/MeOH then concentrated again and heated 15 min more. Concentrated *in vacuo* and dried under high vacuum. Triturated with a small amount of IpOH and allowed to settle over a weekend. Filtered, rinsing with a small amount of IpOH to provide the compound as a white solid.

- 171 -

The following compounds were prepared similarly to the procedure outlined above:

- a) (4-Bocaminomethyl-pyrimidin-2-yl)-[2-(1-methyl-pyrrolidin-2-yl)-ethyl]-amine. M+H 336.5; Calc'd 335.45.

Preparation LXIV - 2-fluoronicotinic acid

In a flame dried 3-necked round bottom flask equipped with a dropping funnel and thermometer, under N₂, THF (250 ml) was added via cannula. LDA (2M in cyclohexane, 54 ml) was added via cannula as the flask was cooled to -78°C. At -78°C, 2-fluoropyridine (8.87 ml) was added dropwise over 10 min. The reaction was stirred for 3 h. Condensation was blown off (with N₂) a few cubes of solid CO₂ and they were added to the mixture. The mixture was warmed to RT once the solution turned yellow, and it was stirred overnight. The reaction was cooled to 0°C and the pH was adjusted to ~2.5 with 5N HCl. The mixture was concentrated in vacuo and extracted with EtOAc. The EtOAc layer was washed with brine, dried over MgSO₄, filtered and concentrated to dryness. The resulting solid was slurried in EtOAc (100 ml), filtered, washed with cold EtOAc and dried at 50°C for 1 h to afford 2-fluoronicotinic acid. M+H 142.1; Calc'd 141.0.

Preparation LXV - 4-cyano-2-methoxypyridine

Under a stream of N₂ and with cooling, Na metal (2.7 g) was added to MeOH (36 ml) with a considerable exotherm. After the Na is dissolved, a solution of 2-chloro-4-cyanopyridine (15 g) in dioxane:MeOH (1:1, 110 ml) was added via dropping funnel over a 10 min period. The reaction was heated to reflux for 3.5 h then cooled at ~10°C overnight. Solid was filtered off and the solid was washed with MeOH. The filtrate was concentrated to ~60 ml and H₂O (60 ml) was added to redissolve a precipitate. Upon further

- 172 -

concentration, a precipitate formed which was washed with H_2O . Further concentration produced additional solids. The solids were combined and dried in vacuo overnight at $35^\circ C$ to provide 4-cyano-2-methoxypyridine which was used as is.

5

Preparation LXVI - (2-methoxypyridin-4-yl)methylamine

4-Cyano-2-methoxypyridine (1.7 g) was dissolved in MeOH (50 ml) and conc. HCl (4.96 ml) was added. Pd/C (10%) was added and H_2 was added and let stand overnight. The solids were
10 filtered through Celite® and the cake was washed with MeOH (~250 ml). Concentration in vacuo produced an oil which was dissolved in MeOH (~20 ml). Et_2O (200 ml) was added and stirred for 1 h. The resulting precipitate was filtered and washed with Et_2O to afford (2-methoxypyridin-4-
15 yl)methylamine (hydrochloride salt) as an off-white solid.

Preparation LXVII - 2-(4-Amino-phenyl)-2-methyl-propionic acid methyl ester

2-Methyl-2-(4-nitro-phenyl)-propionic acid methyl ester (2.1
20 g) was dissolved in THF (70 ml) and acetic acid (5 ml) and Zn (10 g) were added. The mixture was stirred for 1 h and filtered through Celite®. The filtrate was rinsed with EtOAc and the organics were evaporated to a residue which was purified on silica gel chromatography (40%EtOAc/hexanes) to
25 provide the desired compound as a yellow oil. M+H 194.

Preparation LXVIII - 1-(2-tert-Butyl-phenyl)-4-methyl-piperazine

2-tert-Butyl-phenylamine and bis-(2-chloro-ethyl)-
30 methylamine were mixed together with K_2CO_3 (25 g), NaI (10 g) and diglyme (250 mL) and heated at $170^\circ C$ for 8 h. Cooled and filtered solid and evaporated solvent. Diluted with EtOAc, washed with $NaHCO_3$ solution, extracted twice more

- 173 -

with EtOAc, washed with brine, dried over Na_2SO_4 and evaporated to give the compound as a dark solid.

The following compounds were prepared similarly to the
5 procedure outlined above:

a) 1-Bromo-2-(2-methyl-allyloxy)-4-nitro-benzene was prepared from methallyl bromide.

10 **Preparation LXIX 3-(1-Methyl-1,2,3,6-tetrahydro-pyridin-4-yl)-5-trifluoromethyl-phenylamine**
3-(5,5-Dimethyl-[1,3,2]dioxaborinan-2-yl)-5-trifluoromethyl-phenylamine (8.8g, 0.032mol) was added to trifluoro-
methanesulfonic acid 1-methyl-1,2,3,6-tetrahydro-pyridin-4-
15 yl ester (7.91g, 0.032mol) and 2N Na_2CO_3 aqueous solution (25mL) was bubbled through N_2 for 5 min. $\text{Pd}(\text{PPh}_3)_4$ (3.7g, 3.2mmol) was added and the reaction was heated to 80°C for 16 h. The reaction was cooled to RT and diluted with Et_2O (100 mL). The mixture was filtered through Celite® and the
20 filtrate was washed with NaHCO_3 aqueous solution (25 ml) followed by brine (25 mL). The organic phase was dried over Na_2SO_4 and concentrated in vacuo. The desired product was isolated by passing through silica gel column chromatography (EtOAc, then (2M NH_3) in MeOH/EtOAc) to provide a yellow
25 oil.

Preparation LXX - 3,3-Dimethyl-6-nitro-2,3-dihydro-benzo[d]isothiazole 1,1-dioxide

3,3-Dimethyl-2,3-dihydro-benzo[d]isothiazole 1,1-dioxide was
30 added to KNO_3 in H_2SO_4 cooled to 0°C and stirred for 15 min. The reaction was warmed to RT and stirred overnight. The mix was poured into ice and extracted with EtOAc (3x), washed with H_2O and brine, dried and evaporated to give the product which was used without further purification.

- 174 -

The following compounds were prepared similarly to the procedure outlined above:

- 5 a) 1,1,4,4-Tetramethyl-6-nitro-1,2,3,4-tetrahydro-naphthalene

Preparation LXXI - 3-(1-Methyl-1,2,3,4-tetrahydro-pyridin-4-yl)-5-trifluoromethyl-phenylamine

- 10 3-(5,5-Dimethyl-[1,3,2]dioxaborinan-2-yl)-5-trifluoromethyl-phenylamine (1.2 g) was added to trifluoro-methanesulfonic acid 1-methyl-1,2,3,6-tetrahydro-pyridin-4-yl ester (1.0 g), LiCl (500 mg, Aldrich), PPh₃ (300 mg, Aldrich) and 2M Na₂CO₃ aqueous solution (6 ml) and was bubbled with N₂ for 5 min.
- 15 Pd(PPh₃)₄ (300 mg, Aldrich) was added and the reaction was heated to 80°C for 16 h. The reaction was cooled to RT and diluted with Et₂O (100 mL). The mixture was filtered through Celite® and the filtrate was washed with NaHCO₃ aqueous solution (25 ml) followed by brine (25 mL). The
- 20 organic phase was dried over Na₂SO₄ and concentrated in vacuo. The desired compound was isolated by silica gel column chromatography (EtOAc 10% (2M NH₃) in MeOH/EtOAc) to provide yellow oil. M+H 257.2; Calc'd 256.1.

- 25 **Preparation LXXII - Trifluoromethylsulfonic acid 1-methyl-1,2,3,6-tetrahydro-pyridin-4-yl ester**

- In a three-necked round bottom flask equipped with a thermometer and an additional funnel was placed anhydrous THF (200 mL) and 2M LDA (82.8 mL). The solution was cooled
- 30 to -78°C and a solution of 1-methyl-piperidin-4-one (20 mL) in anhydrous THF (70 mL) was added drop-wise. The reaction was warmed to -10°C over 30 min and cooled down again to -78°C. Tf₃NPh (54.32 g) in 200 mL of anhydrous THF was added through the additional funnel over 30 min and anhydrous THF

- 175 -

(30 mL) was added to rinse the funnel. The reaction was warmed to RT and the reaction solution was concentrated in vacuo. The residue was dissolved in Et₂O purified on neutral Al₂O₃ column chromatography (Et₂O as elutant). The product was obtained as orange oil. (20 g)

Preparation LXXIII - 3-(5,5-Dimethyl-[1,3,2]dioxaborinan-2-yl)-5-trifluoromethyl-phenylamine

N₂ was bubbled through a solution of 3-bromo-5-trifluoromethyl-phenylamine (2.38 g), 5,5,5',5'-tetramethyl-[2,2']bi[[1,3,2]dioxaborinanyl] (2.24 g, Frontier Scientific) and KOAc (2.92 g), dppf (165 mg, Aldrich) in anhydrous dioxane (50 ml) for 2 min. PdCl₂ (dppf) (243 mg, Aldrich) was added and the reaction was heated to 80°C for 4 h. After cooling to RT, the mix was diluted with 50 mL of Et₂O, filtered through Celite®, and the filtrate was concentrated in vacuo. The residue was dissolved in Et₂O (100 mL), washed with sat. NaHCO₃ aqueous solution (50 mL) followed by brine (50 mL). The organic phase was dried over Na₂SO₄ and concentrated in vacuo. The residue was dissolved in 3:2 Et₂O/Hex (100 mL), filtered through Celite® and the filtrate was concentrated in vacuo to afford a dark brown semi-solid.

25 Preparation LXXIV - 1-Boc-3-Hydroxymethyl-azetidine

A solution of 1-Boc-azetidine-3-carboxylic acid (1.6 g) and Et₃N (2 ml) in anhydrous THF (60 ml) was cooled to 0°C. Isopropyl chloroformate (1.3 g) was added via a syringe slowly; forming a white precipitate almost immediately. The reaction was stirred for 1 h at 0°C and the precipitate was filtered out. The filtrate was cooled to 0°C again and aqueous NaBH₄ solution (900 mg, 5 ml) was added via pipette and stirred for 1 h. The reaction was quenched with NaHCO₃ solution (50 mL) and the product was extracted with EtOAc

- 176 -

(200 mL). The organic phase was washed with brine (50 mL), dried over Na_2SO_4 and concentrated in vacuo. The residue was dissolved in EtOAc and passed through a short silica gel pad. Concentrating the filtrate in vacuo provided the
5 compound as a light yellow oil.

Preparation LXXV - 1-Boc-3-(3-nitro-5-trifluoromethyl-phenoxymethyl)-azetidine

A mixture of 1-Boc-3-methylsulfonyloxymethyl-azetidine (1.47
10 g), 3-nitro-5-trifluoromethyl-phenol (1.15 g) and K_2CO_3 (1.15 g) in DMF (20 mL) at 80°C was stirred overnight. The reaction was cooled to RT and diluted with 25 mL of sat. NaHCO_3 and 50 mL of EtOAc. The organic phase was separated and washed with brine (25 mL), dried over Na_2SO_4 and
15 concentrated in vacuo. The crude compound was purified by column chromatography (50% EtOAc/hex).

Preparation LXXVI - 2,2-Dimethyl-6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine

20 2,2-Dimethyl-6-nitro-4H-benzo[1,4]oxazin-3-one was added to BH_3 -THF complex (Aldrich) in THF with ice cooling. The mixture was heated to reflux for 2 h then carefully diluted with 12 mL of MeOH and heated to reflux for an additional 1 h. Concentrated HCl (12 mL) was added and heated to reflux
25 for 1 h. The mixture was concentrated and the resulting solid was suspended in a dilute aqueous solution of NaOH (1 M) and extracted with EtOAc (100 mL x 4). The organic layers were washed with H_2O and dried over MgSO_4 . Evaporation of solvent gave a yellow solid.

30

Preparation LXXVII - 2,2,4-Trimethyl-6-nitro-4H-benzo[1,4]oxazin-3-one

2,2-Dimethyl-6-nitro-4H-benzo[1,4]oxazin-3-one (1.1 g) was mixed with MeI (850 mg, Aldrich), K_2CO_3 (1.38 g, Aldrich)

- 177 -

and DMF (30 ml, Aldrich) at 40°C for 48 h. The DMF was removed in vacuo and the residue was diluted with EtOAc (80 ml). The organic phase was washed with H₂O (50 ml), aqueous Na₂SO₃ (50 ml) and brine (50 ml). The resulting solution
5 was dried (MgSO₄) and concentrated to provide the compound which was used as is.

Preparation LXXVIII - 2-Bromo-N-(2-hydroxy-5-nitro-phenyl)-2-methyl-propionamide

10 2-Amino-4-nitro-phenol (3.08 g, Aldrich) was stirred with THF (30 ml, Aldrich) in an ice bath. 2-Bromo-2-methyl-propionyl bromide (2.47 ml, Aldrich) and Et₃N (2.0 g, Aldrich) was slowly added via syringe. The mixture was stirred for 45 min then poured into ice. The aqueous phase
15 was extracted by EtOAc (50 mL x 4). The organic layer was dried and concentrated. The desired product was crystallized from EtOAc. (*Chem. Pharm. Bull* 1996, 44(1) 103-114).

20 **Preparation LXXIX - 2,2-Dimethyl-6-nitro-4H-benzo[1,4]oxazin-3-one**

2-Bromo-N-(2-hydroxy-5-nitro-phenyl)-2-methyl-propionamide was mixed with K₂CO₃ in 20 mL of DMF and stirred overnight at 50°C. The reaction mixture was poured into ice water.
25 The precipitate was collected by filtration and washed with H₂O. The crude compound was recrystallized from EtOH.

Preparation LXXX -4-[1-(2-Bromo-4-nitro-phenyl)-1-methyl-ethyl]-1-methyl-pyridinium iodide

30 1-Methyl-4-[1-methyl-1-(4-nitro-phenyl)-ethyl]-pyridinium (8 g) was dissolved in glacial HOAc (10 ml) then diluted with H₂SO₄ (50 ml), then NBS (3.8 g) was added. After 1 h, additional NBS (1.2 g) was added, 30 min later another 0.5 g of NBS, then 15 min later 200 mg more NBS. After 1 h, the

- 178 -

mixture was neutralized with NH_4OH (conc.) with ice bath cooling. The neutralized mixture was then concentrated and used as is.

- 5 **Preparation LXXXI - 4-[1-(2-Bromo-4-nitro-phenyl)-1-methyl-ethyl]-1-methyl-1,2,3,6-tetrahydro-pyridine**
4-[1-(2-Bromo-4-nitro-phenyl)-1-methyl-ethyl]-1-methyl-pyridiniumiodide was mixed with MeOH (400 ml) and CH_2Cl_2 (200 ml), then treated with NaBH_4 (2.5 g) in portions.
- 10 After stirring at RT for 2 h, the mixture was extracted with CH_2Cl_2 (300 mL x 3). The CH_2Cl_2 layer was washed with brine, dried over Na_2SO_4 and concentrated in vacuo, to provide the desired product.
- 15 **Preparation LXXXII - 1-Methyl-4-[1-methyl-1-(4-nitro-phenyl)-ethyl]-pyridinium iodide**
4-(4-Nitro-benzyl)-pyridine (4.3 g) was mixed with MeI (4 ml, 9.12 g)/NaOH (5N, 30 ml), Bu_4NI (150 mg) and CH_2Cl_2 (50 ml) and stirred at RT overnight. Additional MeI (2 mL) was
- 20 added along with 50 mL of NaOH (5N). 6 h later, more MeI (2 mL) was added. The mixture was stirred at RT over the weekend. The mixture was cooled on ice bath and the base was neutralized by conc. HCl (aq) addition dropwise to pH 7. The compound was used as is.
- 25 **Preparation LXXXIII - 1-Methyl-4-(4-nitro-benzyl)-1,2,3,6-tetrahydro-pyridine**
4-(4-Nitrobenzyl)pyridine (64 g) and TBAI (6 g) were dissolved in CH_2Cl_2 (500 mL) and the solution was suspended
- 30 with NaOH (aq. 5N, 450 mL) in a 3L 3-necked round bottom flask. With vigorous stirring, iodomethane (213 g) was added and stirred vigorously at RT for 60 h (or until blue color disappears). The reaction was quenched with dimethylamine (100 mL) and MeOH (300 mL) and stirred for 2 h. NaBH_4 (19

- 179 -

g) was added to the mixture in small portions. The reaction mixture was stirred for 30 min at RT, then partitioned between $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ (500 mL/500 mL). The organic layer was collected and the aqueous layer was washed with CH_2Cl_2 (300 mL x 3). The combined organic layers was washed with brine then concentrated in vacuo. The residue was purified on a silica wash-column (7% TEA in EtOAc). The desired fractions were combined and concentrated under vacuum to give the desired compound as a dark gray solid. (MS: $M+1=261$).

Preparation LXXXIV - 1-Boc-4-formylpiperidine

4A Molecular sieves were heated to 100°C and a vacuum was applied. They were cooled to RT and purged with N_2 . CH_2Cl_2 (420 ml) and CH_3CN (40 ml), NMO (40 g) and 1-Boc-4-hydroxymethylpiperidine (50 g) were added and the mix was stirred for 5 min then cooled to 15°C . TPAP (4.1 g) is added and an exotherm was observed. The reaction was maintained at RT with external cooling. The reaction was stirred at RT for 3 h, filtered, concentrated, diluted with 50% EtOAc/hexanes and purified on a silica gel plug (50% EtOAc/hexanes). The eluant fractions were concentrated to afford a yellow oil.

25 Preparation LXXXV 2-Chloro-4-cyanopyridine

2-Chloro-4-cyanopyridine was prepared similar to the method described by Daves et al., J. Het. Chem., 1, 130-32 (1964).

30 Preparation LXXXVI 4-(2-tert-Butyl-5-nitro-phenyl)-but-3-en-1-ol

A mix of 1-(tert-butyl)-2-bromo-4-nitrobenzene (3.652 g), TEA (5.92 ml), 3-buten-1-ol (5.48 ml), $\text{Pd}(\text{OAc})_2$ (32 mg), $\text{Pd}(\text{PPh}_3)_4$ (327 mg) and toluene (40 ml) was degassed with nitrogen and heated in a sealed vessel for 16 h at 120°C .

- 180 -

The next day, the reaction mixture was cooled to RT, filtered, and concentrated in vacuo. The crude was eluted on a silica gel column with 15% to 22% EtOAc/hexanes gradient system to yield a yellow-brown oil.

5

Preparation LXXXVII 4-(2-tert-Butyl-5-nitro-phenyl)-but-3-enal

4-(2-tert-Butyl-5-nitro-phenyl)-but-3-en-1-ol (1.024 g) was dissolved in 10 ml of CH₂Cl₂ and added dropwise over 5 min to a -78°C mix of oxalyl chloride (0.645 ml), DMSO (0.583 ml), and 10 ml CH₂Cl₂. The reaction was stirred at -78°C for 1 h, then treated with a solution of TEA (1.52 ml) in 7 ml CH₂Cl₂ and stirred at -78°C for an additional 25 min, then warmed to -30°C for 35 min. The reaction was treated with 50 ml of saturated aqueous NH₄Cl, diluted with H₂O and extracted with EtOAc. The organic layer was brine-washed, dried over Na₂SO₄, filtered, and concentrated in vacuo to yield a yellow oil which was used as is in Preparation LXXXVIII.

20

Preparation LXXXVIII 1-[4-(2-tert-Butyl-5-nitro-phenyl)-but-3-enyl]-pyrrolidine

4-(2-tert-Butyl-5-nitro-phenyl)-but-3-enal (895 mg) was dissolved in 40 ml THF, and to the solution was added pyrrolidine (0.317 ml). To the deep orange solution was added NaBH(OAc)₃ (1.151 g) and glacial AcOH (0.207 ml). The reaction was stirred at RT overnight, then treated with saturated aqueous NaHCO₃ and diluted with Et₂O and some 1N NaOH. The layers were separated, and the organic layer was extracted with aqueous 2N HCl. The acidic aqueous layer was basified to pH>12 with 6 N NaOH, extracted with Et₂O, brine-washed, dried over Na₂SO₄, filtered, and concentrated in vacuo to provide 1-[4-(2-tert-butyl-5-nitro-phenyl)-but-3-enyl]-pyrrolidine as a orange-brown oil.

30

- 181 -

**Preparation LXXXVIX N-Boc-(2-chloropyrimidin-4-yl)-
methylamine**


To 2-chloropyrimidine-4-carbonitrile [2.5 g, prepared by the
5 procedure of Daves et. al. [*J. Het. Chem.* **1964**, 1, 130-132)]
in EtOH (250 ml) under N₂ was added Boc₂O (7.3 g). After
the mixture was briefly placed under high vacuum and flushed
with N₂, 10% Pd/C (219 mg) was added. H₂ was bubbled through
the mixture (using balloon pressure with a needle outlet) as
10 it stirred 4.2 h at RT. After filtration through Celite®,
addition of 1.0 g additional Boc₂O, and concentration, the
residue was purified by silica gel chromatography (5:1 →
4:1 hexanes/EtOAc) to obtain N-Boc-(2-chloropyrimidin-4-yl)-
methylamine.

15

**Preparation XC Methanesulfonic acid 1-Boc-azetidin-3-
ylmethyl ester**

To a solution of (1-Boc-azetidin-3-yl)-methanol (1.06 g, 5.7
mmol), TEA (1.18 mL, 8.52mmol) in CH₂Cl₂ at 0°C was added
20 MeSO₂Cl (0.53 mL, 6.82 mmol) via a syringe. The reaction
was warmed to RT over 2 h and stirring was continued at RT
for 2 h. The white solid formed was removed by filtration
and the filtrate was washed with 25 mL of H₂O. The organic
phase was dried over Na₂SO₄, and concentrated *in vacuo* to
25 afford yellow oil.

182

Example 1

N-(4-Chlorophenyl)[2-(6-quinolylamino)(3-pyridyl)]carboxamide

Step A - Preparation of 2-(6-quinolylamino)pyridine-3-carboxylic acid

A mixture of 2-chloronicotinic acid (1.5 g) and 6-aminoquinoline (1.4 g) was heated at 140°C neat for 2 h. The reaction was cooled to give a brown solid used in the next step without further purification. MS (ES⁺): 266 (M+H)⁺; (ES⁻): 264 (M-H)⁻.


Step B - Preparation of N-(4-chlorophenyl)[2-(6-quinolylamino)(3-pyridyl)]carboxamide

To a mixture of 2-(6-quinolylamino)pyridine-3-carboxylic acid (250 mg, from Step A) and 4-chloroaniline (120 mg) and DIEA (220 ul) in DMF (15 ml) was added EDC (220 mg) and HOBt (130 mg). The reaction was stirred at 50°C for 3 h. The solution was evaporated under reduced pressure and mixed with CH₂Cl₂ (50 ml). The resulting solution was washed with 1N HCl, saturated NaHCO₃, H₂O and brine. The organic layer was dried over Na₂SO₄, evaporated under reduced pressure and triturated with CH₂Cl₂. The precipitate was filtered and washed with EtOH to give the titled compound.

183

MS (ES⁺): 375 (M+H)⁺; (ES⁻): 373 (M-H)⁻. Calc'd for C₂₁H₁₅ClN₄O - 374.8.


Example 2



N-(4-Chlorophenyl)[2-(5-isoquinolyamino)(3-pyridyl)]carboxamide

The title compound was analogously synthesized by the method described in Example 1. MS (ES⁺): 375 (M+H)⁺. Calc'd for C₂₁H₁₅ClN₄O - 374.8.

Example 3



N-(4-Chlorophenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


Step A - Preparation of (2-chloro(3-pyridyl))-N-(4-chlorophenyl)carboxamide

To a mixture of 2-chloronicotinic acid (4 g) and 4-chloroaniline (3.2 g) and DIEA (6 ml) in CH_2Cl_2 (200 ml) was added EDC (6 g) and HOBT (3.3 g). The reaction was stirred at RT overnight and washed with 2 N NaOH (100 ml), H_2O (150 ml) and brine (100 ml). The organic layer was dried over Na_2SO_4 and evaporated to give (2-chloro(3-pyridyl))-N-(4-chlorophenyl)carboxamide.

Step B - Preparation of N-(4-chlorophenyl)[2-(1H-indazol-5-ylamino)(3-pyridyl)]carboxamide

A mixture of (2-chloro(3-pyridyl))-N-(4-chlorophenyl)carboxamide (200 mg, from Step A) and 6-aminoindazole (150 mg) was heated at 150°C neat for 2 h. The reaction was cooled and washed with MeOH. The resulting solid was filtered to give the titled product. MS (ES⁺): 364 (M+H)⁺; (ES⁻): 362 (M-H). Calc'd for $\text{C}_{19}\text{H}_{14}\text{ClN}_5\text{O}$ - 363.8.


Example 4



N-(4-Chlorophenyl)[2-(1H-indazol-5-ylamino)(3-pyridyl)]carboxamide

185

The title compound was analogously synthesized by the method described in Example 3. MS (ES⁺): 364 (M+H)⁺; (ES⁻): 362 (M-H)⁻. Calc'd for C₁₉H₁₄ClN₅O - 363.8.

Example 5

2-(1H-Indazol-6-ylamino)-N-(4-isopropylphenyl)nicotinamide

Step A: Preparation of (2-chloro(3-pyridyl))-N-(4-isopropylphenyl)carboxamide

To a mixture of 2-chloronicotinic acid (6.3 g) and 4-isopropylaniline (5.26 ml) and DIEA (10 ml) in CH₂Cl₂ (200 ml) was added EDC (10 g) and HOBt (5.4 g). The reaction was stirred at RT overnight and washed with 2N NaOH (100 ml), H₂O (250 ml) and brine (100 ml). The organic layer was dried over Na₂SO₄ and evaporated to give (2-chloro(3-pyridyl))-N-(4-isopropylphenyl)carboxamide.

Step B: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylethyl)phenyl]carboxamide hydrochloride


A mixture of (2-chloro(3-pyridyl))-N-(4-isopropylphenyl)carboxamide (1.5 g, Step A) and 6-aminoindazole (880 mg) was heated at 160°C in NMP for 3 h. The reaction was cooled and diluted with CH₂Cl₂ and washed with water twice, followed by brine. The organic layer was

186

dried with Na_2SO_4 and evaporated under reduced pressure. The residue was purified by column chromatography with 35% EtOAc/Hexane and further mixed with MeOH and 1 N HCl/Et₂O (3 ml). The solution was evaporated to furnish the titled compound. MS (ES⁺): 372 (M+H)⁺; (ES⁻): 371 (M-H). Calc'd. for $\text{C}_{22}\text{H}_{21}\text{N}_5\text{O}$ - 371.2.

The following compounds (Examples 6-23) were analogously synthesized by the method described in Example 5. Detailed intermediate preparations are described.


Example 6



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(methylethyl)phenyl]carboxamide

MS (ES⁺): 372 (M+H)⁺. Calc'd. for $\text{C}_{22}\text{H}_{21}\text{N}_5\text{O}$ - 371.2.

Example 7




187

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylpropyl)phenyl]carboxamide

MS (ES⁺): 386 (M+H)⁺; (ES⁻): 384 (M-H). Calc'd. for C₂₃H₂₃N₅O - 385.2.


Example 8



N-[4-(tert-Butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

MS (ES⁺): 386 (M+H)⁺; (ES⁻): 384 (M-H). Calc'd. for C₂₃H₂₃N₅O - 385.2.


Example 9



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(trifluoromethyl)phenyl]carboxamide


MS (ES⁺): 398 (M+H)⁺; (ES⁻): 396 (M-H). Calc'd. for C₂₀H₁₄F₃N₅O - 397.1.

188

Example 10

N-[3-(tert-Butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


MS (ES+): 386 (M+H)⁺; (ES-): 384 (M-H). Calc'd. for
C₂₃H₂₃N₅O - 385.2.

Example 11

[2-(Benzotriazol-6-ylamino)(3-pyridyl)]-N-[4-(tert-butyl)phenyl]carboxamide


MS (ES+): 387 (M+H)⁺; (ES-): 385 (M-H). Calc'd. for
C₂₂H₂₂N₆O - 386.2.

189

Example 12

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(3-phenylpyrazol-5-yl)carboxamide


MS: 396 (M+1). Calc'd. for $C_{22}H_{17}N_7O$ - 395.4.

Example 13

N-3-aminosulfonyl(4-chloro-phenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


MS (ES+): 443 (M+H); (ES-): 441 (M-H). Calc'd. for $C_{19}H_{15}ClN_6O_3S$ - 442.1.

190

Example 14

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-methyl-2-oxo-1,2-dihydroquinol-7-yl)carboxamide


MS (ES⁺): 411 (M+H); (ES⁻): 409 (M-H). Calc'd. for
C₂₃H₁₈N₆O₂ - 410.1.

Example 15

N-[4-(Methylethyl)phenyl]{2-[(4-methyl-2-oxo(7-hydroquinolyl))amino](3-pyridyl)}carboxamide


MS (ES⁺): 413 (M+H); (ES⁻): 411 (M-H). Calc'd. for
C₂₅H₂₄N₄O₂ - 412.2.

191

Example 16

N-[5-(tert-Butyl)isoxazol-3-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


MS (ES+): 377 (M+H); (ES-): 375 (M-H). Calc'd. for $C_{20}H_{20}N_6O_2$ - 412.2.

Example 17

N-[5-(tert-Butyl)-1-methylpyrazol-3-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


MS (ES+): 390 (M+H); (ES-): 388 (M-H). Calc'd. for $C_{21}H_{23}N_7O$ - 389.2.

192

Example 18

N-[4-(tert-Butyl)(1,3-thiazol-2-yl)][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


MS (ES⁺): 393 (M+H); (ES⁻): 391 (M-H). Calc'd. for C₂₀H₂₀N₆OS - 392.5.

Example 19

N-[5-(tert-Butyl)(1,3,4-thiadiazol-2-yl)][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

MS (ES⁺): 394 (M+H); (ES⁻): 392 (M-H). Calc'd. for C₁₉H₁₉N₇OS - 393.5.

193

Example 20

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(1,1,2,2,3,3,4,4,4-nonafluorobutyl)phenyl]carboxamide


Step A: Preparation of 4-(1,1,2,2,3,3,4,4,4-nonafluorobutyl)phenylamine

The title compound was synthesized analogously by the method described in Gregory, W.A. et al., J. Med. Chem, 1990, 33(9) to give 4-nitro-(1,1,2,2,3,3,4,4,4-nonafluorobutyl)benzene. The mixture of the above intermediate (1.0 mmol), iron powder (5.0 mmol) and NH_4Cl (0.7 mmol) in EtOH (3 mL) and H_2O (3 mL) was stirred for 4 hours at 80°C . Filtration and concentration to give crude title product, which was used in next step without further purification.

Step B: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[4-(1,1,2,2,3,3,4,4,4-nonafluorobutyl)phenyl]carboxamide

The title compound was prepared from 4-(1,1,2,2,3,3,4,4,4-nonafluorobutyl)phenylamine (Step A) similar to the method described in Example 5. MS (ES⁺): 548 (M+H); (ES⁻): 546 (M-H). Calc'd. for $\text{C}_{22}\text{H}_{15}\text{F}_9\text{N}_4\text{O}$ - 522.4.

194

Example 21


{2-[(1-Methyl(1H-indazol-6-yl))amino](3-pyridyl))-N-[4-(methylethyl)phenyl]carboxamide

Step A: Preparation of 1-methyl-6-amino-1H-indazole

To a solution of 6-nitroindazole (8 g) in THF (200 ml) was added NaH (2.5 g) at 0°C. The reaction was stirred for 30 min and MeI (3.7 ml) was added and stirred at RT overnight. The reaction was quenched with water and extracted with EtOAc twice, then was further purified by column chromatography to give 1-methyl-6-nitro-1H-indazole which was hydrogenated with an H₂ atmosphere in the presence of Pd/C (400 mg) to give 1-methyl-6-amino-1H-indazole.


Step B Preparation of (2-[(1-methyl(1H-indazol-6-yl))amino](3-pyridyl))-N-[4-(methylethyl)phenyl]-carboxamide

The title compound was prepared from 1-methyl-6-amino-1H-indazole (Step A) similar to the method described in Example 5. MS (ES⁺): 386 (M+H)⁺; (ES⁻): 384 (M-H). Calc'd. for C₂₃H₂₃N₅O - 385.2.

Example 22

N-[4-(tert-Butyl)phenyl]{2-[(7-bromo(1H-indazol-6-yl))amino](3-pyridyl)}carboxamide

A mixture of N-[4-(tert-butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide (620 mg) (Example 8) and NBS (330 mg) in benzene (50 ml) was heated at 80°C for 3 h. The solvent was evaporated under reduced pressure and the residue was purified by column chromatography to give the title compound. MS (ES⁺): 464 (M+H)⁺; (ES⁻): 462 (M-H).
Calc'd. for C₂₃H₂₂BrN₅O - 463.1.

Example 23

2-(1H-Indazol-6-ylamino)-N-[4-tert-butyl-3-(1,2,3,6-tetrahydropyridin-4-yl)phenyl]nicotinamide

Step A: Preparation of 2-bromo-1-tert-butyl-4-nitrobenzene

NBS (125.0 g, 697.5 mmol, 1.5 eq) was slowly added to a solution of TFA:H₂SO₄ (5:1, 750 mL) and tert-butyl-4-nitrobenzene (100.0 g, 558.0 mmol) at RT. The solution was stirred for 24 h and poured over 5 kg of ice. The resulting suspension was filtered and washed with a 1:1 MeOH:H₂O solution (200 mL) and dried in a vacuum oven. MS (ES⁺): 258.1, 260.1 (M+H)⁺. Calc'd for C₁₀H₁₂BrNO₂: 257.0.

Step B: Preparation of 4-(2-tert-butyl-5-nitrophenyl)pyridine

To a solution of 2-bromo-1-tert-butyl-4-nitrobenzene (8.6 g, 33.3 mmol, Step A) and toluene (70 mL) in a 150 mL round bottom flask, 4-pyridylboronic acid (4.5 g, 36.6 mmol, 1.1 eq), Pd(PPh₃)₄ (3.8 g, 3.3 mmol, 0.1 eq) and K₂CO₃ (13.8 g, 99.9 mmol, 3 eq) were added. The solution was stirred for 24 h at 80°C before cooling to RT. The solution was filtered through a pad of Celite® and purified by silica flash chromatography (30% EtOAc/Hexanes). This afforded the desired product as a yellow solid. MS (ES⁺): 257.2 (M+H)⁺; (ES⁻): 255.2 (M-H)⁻. Calc'd for C₁₅H₁₆N₂O₂: 256.1.

Step C: Preparation of 4-(2-tert-butyl-5-nitrophenyl)-1-methylpyridinium

4-(2-tert-Butyl-5-nitrophenyl)pyridine (2.0 g, 7.8 mmol, Step B) was added to a round-bottom flask and dissolved in EtOH (10 mL). CH₃I (30 mL) was added to the flask which was placed in a 80°C sand bath and heated to reflux. After 6 h, the solution was cooled to RT and the excess CH₃I and EtOH were stripped-off under reduced

197


pressure resulting in the desired compound as a light brown solid. MS (ES+): 271.2 (M+H)⁺; (ES-): 269.2 (M-H)⁻. Calc'd for C₁₆H₁₉N₂O₂⁺: 271.1.

Step D: Preparation of 4-tert-butyl-3-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)aniline

4-(2-tert-Butyl-5-nitrophenyl)-1-methylpyridinium (2.1 g, 7.8 mmol, Step C) was added to a 100 mL round-bottom flask and dissolved in a 10% H₂O/EtOH mixture. To the flask iron dust (1.31 g, 23.4 mmol, 3 eq) and NH₄Cl (460 mg, 8.6 mmol, 1.1 eq) were added. The flask was placed in a 100°C sand bath and heated to reflux. After 2 h, the solution was cooled to RT and filtered through a pad of Celite®. The resulting solution was stripped down to a yellow solid and redissolved in MeOH (20 mL, anhydrous). The solution was cooled to 0°C by placing it in an ice bath and slowly adding NaBH₄ (450 mg, 11.7 mmol, 1.5 eq). After addition of the NaBH₄, the solution was cooled to RT and stirred for 30 min. The solvent was stripped-off under vacuum and the solid was redissolved in CH₂Cl₂ and filtered. The solution was stripped-off under vacuum to afford of an amorphous clear yellow solid. MS (ES+): 245.2 (M+H)⁺. Calc'd for C₁₆H₂₄N₂: 244.2.

Step E Preparation of 2-(1H-indazol-6-ylamino)-N-[4-tert-butyl-3-(1,2,3,6-tetrahydropyridin-4-yl)phenyl]nicotinamide

The title compound was prepared from 4-tert-butyl-3-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)aniline (Step D) similar to the method described in Example 5. MS: (ES+) 480.3 (M+H). Calc'd for C₂₉H₃₂N₆O - 480.6.

Example 24

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-
 {4-[2,2,2-trifluoro-1-hydroxy-1-
 (trifluoromethyl)ethyl]phenyl}carboxamide

Step A: Preparation of 2-(1H-indazol-6-ylamino)pyridine-3-
 carboxylic acid

A mixture of 6-aminoindazole (6.7 g) and 2-chloronicotinic acid (8 g) was heated neat at 150°C for 2 h. The reaction was cooled and washed with acetone to give 2-(1H-indazol-6-ylamino)pyridine-3-carboxylic acid.

Step B: Preparation of [2-(1H-indazol-6-ylamino)(3-
 pyridyl)]-N-{4-[2,2,2-trifluoro-1-hydroxy-1-
 (trifluoromethyl)ethyl]phenyl}carboxamide


2-(1H-Indazol-6-ylamino)pyridine-3-carboxylic acid (500 mg, Step A) was reacted with 4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenylamine (340 mg) and EDC (500 mg) and HOBt (270 mg) in DMF at RT overnight. The reaction was diluted with CH₂Cl₂ and washed with H₂O followed by brine. The organic layer was dried with Na₂SO₄ and evaporated under reduced pressure. The residue was purified by column chromatography to give the title compound. MS

199

(ES⁺):496 (M+H)⁺; (ES⁻):394 (M-H). Calc'd. for C₂₂H₁₅F₆N₅O₂ - 495.1.

The following compounds (Examples 25-42) were synthesized similar to the method described in Example 24. Detailed intermediate preparations are described.


Example 25



N-[5-(tert-Butyl)-2-methoxyphenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

MS (ES⁺):416 (M+H)⁺; (ES⁻):414 (M-H). Calc'd. for C₂₄H₂₅N₅O₂ - 415.2.


Example 26



200


[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-{6-[4-(trifluoromethyl)piperidyl](3-pyridyl)}carboxamide

MS (ES⁺): 482 (M+H)⁺; (ES⁻): 480 (M-H). Calc'd. for C₂₄H₂₂F₃N₇O - 481.2.

Example 27

2-(1H-Indazol-6-ylamino)-N-(1-oxo-1,2,3,4-tetrahydroisoquinolin-7-yl)-nicotinamide


MS (ES⁺): 399 (M+H)⁺; (ES⁻): 397 (M-H). Calc'd. C₂₂H₁₈N₆O₂ - 398.1.

Example 28

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylethoxy)phenyl]carboxamide

201

MS (ES⁺): 388 (M+H)⁺; (ES⁻): 386 (M-H). Calc'd. for C₂₂H₂₁N₅O₂
- 387.2.

Example 29

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl)carboxamide


Step A: Preparation of [1-(4-amino-phenyl)-ethyl]carbamic acid tert-butyl ester

A mixture of 1-(S)-1-(4-nitrophenyl)ethylamine hydrochloride (2 g) and Boc₂O (2.6 g) and NaHCO₃ (3 g) in MeOH/H₂O (1:1, 200 ml) was stirred at RT overnight. The reaction was extracted with EtOAc twice then washed with water followed by brine. The organic layer was dried with Na₂SO₄ and evaporated under reduced pressure to give the protected nitrophenyl ethylamine. Boc-1-(S)-1-(4-nitrophenyl)ethylamine (1 g) was hydrogenated at H₂ atmosphere in the presence of Pd/C (200 mg) to give Boc protected aniline (0.8 g). The intermediate was deprotected with 4N HCl/dioxane to give the title compound as the HCl salt.

Step B: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-(4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl)carboxamide


202

The title compound was prepared from [1-(4-amino-phenyl)ethyl]carbamic acid tert-butyl ester (Step A) similar to the method described in Example 24. MS (ES⁺): 373 (M+H)⁺; (ES⁻): 371 (M-H). Calc'd. for C₂₁H₂₀N₆O - 372.2.

Example 30

N-(4-((1S)-1-[(Methylethyl)amino]ethyl)phenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

A mixture of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-(4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)-ethyl]phenyl)carboxamide (100 mg, Example 29), NaBH(OAc)₃ (2 eq), acetone (5 ml) and AcOH (0.2 ml) in CH₂Cl₂ was stirred at RT for 4 h. The solvent was evaporated and the residue was purified by prep-HPLC to give the title compound as TFA salt. MS (ES⁺): 415 (M+H)⁺; (ES⁻): 413 (M-H). Calc'd. for C₂₄H₂₆N₆O - 414.2.

Example 31

N-[4-(tert-Butyl)-3-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

Step A: Preparation of 1-[2-(tert-butyl)-5-aminophenyl]-4-methylpiperazine


A mixture of 2-t-butylaniline (5.4 g) and methylchloroethylamine hydrochloride (7 g) and K_2CO_3 (5 g) in NaI (2 g) in diglyme (150 ml) was heated at 170°C for 8 h. The reaction was filtered and the filtrate was evaporated under high vacuum. The residue was mixed with EtOAc (200 ml) and H_2O (200 ml) and extracted with EtOAc twice. The combined organic layer was washed with brine and dried over Na_2SO_4 and evaporated to give crude 1-[2-(tert-butylphenyl)]-4-methylpiperazine. The crude 1-[2-(tert-butylphenyl)]-4-methylpiperazine (260 mg) was stirred with H_2SO_4 (3 ml) at 0°C and HNO_3 (1.2 ml, 70%) was slowly added to the reaction. The reaction was warmed to RT, stirred for 30 min, poured on ice and basified with K_2CO_3 slowly. The solution was extracted with EtOAc three times, washed with H_2O , followed by brine, dried over Na_2SO_4 and evaporated under reduced pressure. The residue was purified by column chromatography to give 1-[2-(tert-butyl)-5-nitrophenyl]-4-methylpiperazine

(260 mg), which was hydrogenated under H₂ atmosphere to give 1-[2-(tert-butyl)-5-aminophenyl]-4-methylpiperazine.

Step B: Preparation of N-[4-(tert-butyl)-3-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was prepared from 1-[2-(tert-butyl)-5-aminophenyl]-4-methylpiperazine (Step A) similar to the method described in Example 24. MS (ES⁺): 484 (M+H)⁺; (ES⁻): 482 (M-H). Calc'd. C₂₈H₃₃N₇O - 483.3.

Example 32



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(4-methylpiperazinyl)phenyl]carboxamide

Step A: Preparation of 1-(5-aminophenyl)-4-methylpiperazine


The intermediate was analogously synthesized from 3-nitroaniline by the method described in Example 31, Step A.

Step B: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[3-(4-methylpiperazinyl)phenyl]carboxamide

The title compound was prepared from 1-(5-amino-phenyl)-4-methylpiperazine (Step A) similar to the method

205

described in Example 24. MS (ES⁺): 428 (M+H)⁺; (ES⁻): 426 (M-H). Calc'd. C₂₄H₂₅N₇O - 427.2.

Example 33

N-[4-(tert-Butyl)-2-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


Step A: Preparation of 4-(tert-butyl)-2-(4-methylpiperazinyl)phenylamine

A mixture of 1-(tert-butyl)-2-bromo-4-nitrobenzene (3 g) and N-methylpiperazine (8 g) was heated neat at 130°C for 4 h. The residue was purified by column chromatography to give 1-[4-bromo-5-(tert-butyl)-2-nitrophenyl]-4-methylpiperazine, which was hydrogenated to furnish 4-(tert-butyl)-2-(4-methylpiperazinyl)-phenylamine.

Step B: Preparation of N-[4-(tert-butyl)-2-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was prepared from 4-(tert-butyl)-2-(4-methylpiperazinyl)phenylamine (Step A) similar to the method described in Example 24. MS (ES⁺): 484 (M+H)⁺; (ES⁻): 482 (M-H). Calc'd. C₂₈H₃₃N₇O - 483.3.

206

Example 34

N-{2-[2-(Dimethylamino)ethoxy]-5-(tert-butyl)phenyl}[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


Step A: Preparation of {2-[4-(tert-butyl)-2-aminophenoxy]ethyl}dimethylamine

DEAD (2.6 ml) was added to a mixture of 2-nitro-4-tert-butylphenol (2 g) and N,N-dimethylethanolamine (1.3 g) and Ph_3P (4 g) in THF (50 ml). The reaction was stirred at RT for 1 h, diluted with EtOAc (50 ml) and washed with 1 N HCl twice. The aqueous layer was basified with NaHCO_3 , extracted with EtOAc twice and washed with H_2O and brine. The organic layer was dried over Na_2SO_4 and evaporated to give {2-[4-(tert-butyl)-2-nitrophenoxy]ethyl}-dimethylamine. It was hydrogenated under H_2 atmosphere to give {2-[4-(tert-butyl)-2-aminophenoxy]ethyl}-dimethylamine.

Step B: Preparation of N-{2-[2-(dimethylamino)ethoxy]-5-(tert-butyl)phenyl}[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was prepared from {2-[4-(tert-butyl)-2-aminophenoxy]ethyl}dimethylamine (Step A) similar to the method described in Example 24. MS (ES⁺): 473(M+H)⁺; (ES⁻): 471(M-H). Calc'd. for $\text{C}_{27}\text{H}_{32}\text{N}_6\text{O}_2$ -472.3.

207

Example 35


N-(3-[2-(Dimethylamino)ethoxy]phenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

Step A: Preparation of [2-(2-aminophenoxy)ethyl]-
dimethylamine

The intermediate was synthesized from 2-nitrophenol similar to the method described in Example 34, Step A.

Step B: Preparation of N-(3-[2-(
(Dimethylamino)ethoxy]phenyl)[2-(1H-indazol-6-ylamino)(3-
pyridyl)]carboxamide

The title compound was prepared from [2-(2-aminophenoxy)ethyl]dimethylamine (Step A) similar to the method described in Example 24. MS (ES⁺): 417(M+H)⁺; (ES⁻): 415(M-H). Calc'd. for C₂₃H₂₄N₆O₂ - 416.2.


Example 36

208

N-(3-Hydroxy-4-methoxyphenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

MS (ES⁺):376 (M+H)⁺; (ES⁻):374 (M-H). Calc'd. for C₂₀H₁₇N₅O₃
- 375.1.


Example 37



N-{3-[2-(Dimethylamino)ethoxy]-4-methoxyphenyl}[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was prepared from N-(3-hydroxy-4-methoxyphenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide (Example 36) by a method similar to that described in Example 34. MS (ES⁺):447 (M+H)⁺; (ES⁻): 445 (M-H). Calc'd. for C₂₄H₂₆N₆O₃ - 446.2.


Example 38



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-methoxy-3-(1-methyl(4-piperidyl)oxy)phenyl]carboxamide

The title compound was prepared from N-(3-hydroxy-4-methoxyphenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]-carboxamide (Example 36) by a method similar to that described in Example 34 using 4-hydroxy-N-methylpiperidine. MS (ES⁺): 473 (M+H)⁺; (ES⁻): 471 (M-H). Calc'd. for C₂₆H₂₈N₆O₃ - 472.2.

Example 39



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolin-2-yl)carboxamide

Step A: Preparation of 2-amino-5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinoline


7-Nitro-2,3,4-trihydroisoquinolin-1-one (500 mg) was heated in POCl₃ (10 ml) to reflux for 8 h. The mixture was evaporated, mixed with toluene and evaporated again. The residue was dissolved in THF, H₂NNH₂ (1 ml) was slowly added to the reaction and stirred for 2 h. The reaction was evaporated, heated with HC(OEt)₃ (15 ml) at 115°C for 2 h, extracted with EtOAc and hydrogenated to give 2-amino-5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinoline.

210

Step B: Preparation of [2-(1H-indazol-6-ylamino) (3-pyridyl)]-N-(5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolin-2-yl)carboxamide

The title compound was prepared from 2-amino-5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinoline (Step A) similar to the method described in Example 24. MS (ES⁺): 473 (M+H)⁺; (ES⁻): 471 (M-H). Calc'd. for C₂₃H₁₈N₈O - 422.2.

Example 40



[2-((3-[2-(Dimethylamino)ethoxy] (1H-indazol-6-yl))amino) (3-pyridyl)]-N-[4-(tert-butyl)phenyl]carboxamide

Step A: Preparation of 6-nitro-1-Boc-2-hydroindazol-3-one


A mixture of 6-nitro-1H-2-hydroindazol-3-one (1.8 g) and Boc₂O (3 g) and DMAP (300 mg) in CH₂Cl₂ (30 ml) was stirred at RT overnight. The reaction was washed with water followed by brine. The organic layer was dried with Na₂SO₄ and evaporated under reduced pressure to give Boc-protected indazole.

Step B: Preparation of [2-((3-[2-(dimethylamino)ethoxy] (1H-indazol-6-yl))amino) (3-pyridyl)]-N-[4-(tert-butyl)phenyl]carboxamide

211

To a mixture of 6-nitro-1-Boc-2-hydroindazol-3-one (800 mg, Step A) and N,N-dimethylethanolamine (400 mg) and PPh₃ (1.1 g) in THF (20 ml) was added DEAD (0.75 ml). The reaction was stirred at RT for 2 h, diluted with EtOAc (50 ml) and washed with H₂O and brine. The organic layer was dried over Na₂SO₄, evaporated and the residue was purified by column chromatography to give dimethyl[2-(6-nitro(1-Boc-indazol-3-yloxy))ethyl]amine. It was hydrogenated in an H₂ atmosphere in the presence of Pd/C (100 mg) to give dimethyl[2-(6-amino-(1-Boc-indazol-3-yloxy))ethyl]amine. This compound (40 mg) and 2-chloronicotinic acid were heated at 130°C in pentanol overnight and coupled and deprotected simultaneously. The solvent was evaporated and the residue was used for coupling to give the title compound. MS (ES⁺): 473 (M+H)⁺; (ES⁻): 471 (M-H). Calc'd. for C₂₇H₃₂N₆O₂ - 472.3.

Example 41



N-[3,3-Dimethyl-1-(4-piperidylmethyl)indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

Step A: Preparation of tert-butyl 4-[(6-nitro-3,3-dimethylindolinyl)methyl]piperidinecarboxylate

3,3-Dimethyl-6-nitroindoline (450 mg) was dissolved in 20 mL of dichloroethane, N-boc-4-formylpiperidine (750 mg) was added to the mixture, followed by 2 g NaHB(OAc)₃ and 1 mL of glacial AcOH. The mixture was stirred at RT


overnight. Saturated NaHCO_3 solution (20 mL) was added to the reaction mixture and stirred for 1 h. The resulting mixture was separated by separation funnel, the organic layer was extracted once with saturated NaHCO_3 solution and once with brine. The resulting organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The crude material was purified by flash chromatography on silica gel with 9:1 Hexane:EtOAc to afford an orange oil. MS: 290 (M-99). Calc'd. for $\text{C}_{21}\text{H}_{31}\text{N}_3\text{O}_4$ - 389.5.

Step B: Preparation of 3,3-dimethyl-1-piperidin-4-ylmethyl-2,3-dihydro-1H-indol-6-ylamine

tert-Butyl 4-[(6-nitro-3,3-dimethylindolinyl)-methyl]piperidinecarboxylate (Step A, 900 mg) was dissolved in 10 mL MeOH, the mixture was bubbled with H_2 for 10 min. 10% Pd/C (30 mg) was added and the mixture was stirred under H_2 overnight. The mixture was filtered through Celite® and concentrated in vacuo. The crude material was purified by flash chromatography on silica gel with 1:1 Hexane:EtOAc to afford a colorless oil. MS: 360 (M+1). Calc'd. for $\text{C}_{21}\text{H}_{33}\text{N}_3\text{O}_2$ - 359.5.


Step C: Preparation of N-[3,3-dimethyl-1-(4-piperidylmethyl)indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was prepared from 3,3-dimethyl-1-piperidin-4-ylmethyl-2,3-dihydro-1H-indol-6-ylamine (Step B) similar to the method described in Example 24. MS: 496 (M+1). Calc'd. for $\text{C}_{29}\text{H}_{33}\text{N}_7\text{O}$ -495.6.

Example 42

N-[3,3-Dimethyl-1-(1-methyl-piperidin-4-ylmethyl)-2,3-dihydro-1H-indol-6-yl]-2-(1H-indazol-6-ylamino)-nicotinamide

N-[3,3-Dimethyl-1-(4-piperidylmethyl)indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide (140 mg, Example 41) was dissolved in 10 mL EtOH. Formaldehyde (10 mL, 37%) was added, followed by 100 mg of NaCNBH₃. The mixture was stirred at RT overnight, and concentrated in vacuo. The crude was extracted between saturated NaHCO₃ solution and EtOAc, the resulting organic layer was dried over MgSO₄, filtered and concentrated in vacuo to afford a yellow solid. This material was further purified by preparative HPLC to yield a white solid. MS: 510 (M+1). Calc'd. for C₃₀H₃₅N₇O - 509.6.

Example 43**2-(1H-Indazol-6-ylamino)-N-(4-phenoxyphenyl)-nicotinamide****Step A: Preparation of (2-chloro-(3-pyridyl))-N-(4-phenoxyphenyl)carboxamide:**

2-Chloronicotinoyl chloride (9.15 g, 0.052 mol) was added to a stirred solution of 4-phenoxyaniline (10.00 g, 0.054 mol) and DIEA (10.00 ml, 0.057 mol) in CH_2Cl_2 (100 ml) at RT. The mixture stirred for 48 h before removal of solvent under reduced pressure. The resulting residue was dissolved EtOAc and washed several times with saturated NaHCO_3 aqueous solution and brine, respectively. The organic layer was dried over Na_2SO_4 and evaporated to leave a solid. This material was re-crystallized from EtOAc/Hexane mixture, followed by filtration and rinsing with Et_2O to leave the desired compound as a white solid. MS m/z: 325 (M+1); 323 (M-1)

Step B: Preparation of [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-phenoxyphenyl)carboxamide:


The title compound was analogously synthesized by the method described in Example 5 from (2-chloro-(3-pyridyl))-N-

215

(4-phenoxyphenyl)carboxamide (Step A). MS: (ES+) 422 (M + 1)⁺; (ES-): 420 (M - 1)⁻. Calc'd. for C₂₅H₁₉N₅O₂ - 421.1.

The following compounds (Examples 44-63) were analogously synthesized by the method described in Example 43. Detailed intermediate preparations are described.


Example 44



[2-(1H-Indazol-5-ylamino)(3-pyridyl)]-N-(4-phenoxyphenyl)carboxamide


MS: (ES+) 422 (M + 1)⁺; (ES-): 420 (M - 1)⁻. Calc'd. for C₂₅H₁₉N₅O₂ - 421.1.

216

Example 45


[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-phenylphenyl)carboxamide

MS: (ES+) 406 (M + 1)⁺; (ES-): 404 (M - 1)⁻. Calc'd.
for C₂₅H₁₉N₅O - 405.2.

Example 46

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylsulfonyl)phenyl]carboxamide

MS: (ES+) 408 (M + 1)⁺; (ES-): 406 (M - 1)⁻. Calc'd.
for C₂₀H₁₇N₅O₃S - 407.1.

Example 47

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(1-methyl(4-piperidyl))indolin-6-yl]carboxamide

Step A: Preparation of 1-(1-methyl(4-piperidyl))-6-nitroindoline

6-Nitroindoline (5 g) was dissolved in 200 mL of dichloroethane. N-Methyl-4-piperidone (5 g) was added to the mixture, followed by $\text{NaBH}(\text{OAc})_3$ (12 g) and 1 mL of glacial AcOH. The mixture was stirred at RT overnight. A saturated NaHCO_3 (200 mL) solution was added to the reaction mixture and stirred for 1 h. The resulting mixture was separated by separation funnel. The organic layer was extracted once with saturated NaHCO_3 solution and once with brine. The resulting organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The crude material was purified by flash chromatography on silica gel with 2:1 EtOAc:MeOH to afford orange oil. MS: 262 (M+1). Calc'd. for $\text{C}_{14}\text{H}_{19}\text{N}_3\text{O}_2$ - 261.3.

Step B: Preparation of 1-(1-methyl-4-piperidyl)indoline-6-ylamine


218

1-(1-Methyl(4-piperidyl))-6-nitroindoline (3 g, Step A) was dissolved in 100 mL MeOH and the mixture was bubbled with H₂ for 10 min. 10% Pd/C (200 mg) was added and the mixture was stirred under H₂ overnight. The mixture was filtered through Celite® and concentrated in vacuo to afford light yellow oil. MS: 232 (M+1). Calc'd. for C₁₄H₂₁N₃ - 231.3.

Step C: Preparation of [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(1-methyl(4-piperidyl))indolin-6-yl]carboxamide

The title compound was analogously synthesized by the method described in Example 43 from 1-(1-methyl-4-piperidyl)indoline-6-ylamine (Step B). MS: 468 (M+1). Calc'd. for C₂₇H₂₉N₇O - 467.6.

Example 48



N-[3,3-Dimethyl-1-(1-methyl(4-piperidyl))indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

Step A: Preparation of N-(2-bromo-5-nitrophenyl)acetamide

2-Bromo-5-nitroaniline (10 g) was dissolved in CH₂Cl₂ (500 mL), DIEA (6.6 g) was added to the mixture, followed by 100 mg of DMAP. The mixture was cooled to 0°C in ice bath. Acetyl chloride (4 g in 50 mL CH₂Cl₂) was added dropwise to the reaction mixture. After the mixture was stirred at RT over 3 h, and extracted once with saturated NaHCO₃ solution

and once with brine. The resulting organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The crude material was purified by flash chromatography on silica gel with 1:1 EtOAc:Hexane to 100% EtOAc to afford a white solid. MS: 258 (M-1). Calc'd. for $\text{C}_8\text{H}_7\text{BrN}_2\text{O}_3$ - 259.1.

Step B: Preparation of N-(2-bromo-5-nitrophenyl)-N-(2-methylprop-2-enyl)acetamide

A suspension of NaH (2 g) (95% powder) in 100 mL anhydrous DMF was cooled to -78°C , and N-(2-bromo-5-nitrophenyl)acetamide (Step A, 7 g) in 50 mL dry DMF was added to the mixture under N_2 . After the mixture was warmed to 0°C , 3-bromo-2-methylpropene (7.3 g in 20 dry DMF) was added to the mixture. The mixture was stirred at RT overnight. The mixture was poured into a container of ice and extracted between saturated NaHCO_3 solution and EtOAc. The resulting organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The crude material was purified by flash chromatography on silica gel with 7:2 Hexane:EtOAc to afford a yellow gum. MS: 314 (M+1). Calc'd. for $\text{C}_{12}\text{H}_{13}\text{BrN}_2\text{O}_3$ - 313.1.

Step C: Preparation of 1-(3,3-dimethyl-6-nitro-2,3-dihydro-indol-1-yl)ethanone

N-(2-Bromo-5-nitrophenyl)-N-(2-methylprop-2-enyl)acetamide (4.5 g, Step B) was dissolved in 50 mL anhydrous DMF, 2.5 g tetraethyl-ammonium chloride, 1.2 g sodium formate, 3 g sodium acetate were added, the resulting mixture was bubbled with nitrogen gas for 10 minutes. $\text{Pd}(\text{OAc})_2$ (350 mg) was added and the mixture was heated at 80°C under N_2 overnight. After the mixture was concentrated in vacuo, it was extracted between saturated NaHCO_3 solution and EtOAc, the resulting organic layer was dried over MgSO_4 , filtered and concentrated in vacuo. The crude material was

220

purified by flash chromatography on silica gel with 2:1 Hexane:EtOAc to afford a yellow gum. MS: 235 (M+1). Calc'd. for $C_{12}H_{14}N_2O_3$ - 234.2.

Step D: Preparation of 3,3-dimethyl-6-nitroindoline

1-(3,3-Dimethyl-6-nitro-2,3-dihydro-indol-1-yl)ethanone (Step C, 1.8 g) was dissolved in 50 mL EtOH, 50 mL 12N HCl was added and the resulting mixture was heated at 70°C overnight. After the mixture was concentrated in vacuo, it was extracted between saturated $NaHCO_3$ solution and EtOAc. The resulting organic layer was dried over $MgSO_4$, filtered and concentrated in vacuo to afford a yellow solid. MS: 193 (M+1). Calc'd. for $C_{10}H_{12}N_2O_2$ - 192.2.

Step E: Preparation of 3,3-dimethyl-1-(4-methyl-piperazin-1-yl)-6-nitro-2,3-dihydro-1H-indole

3,3-Dimethyl-6-nitroindoline (0.8 g Step D) was dissolved in 50 mL of dichloroethane, N-methyl-4-piperidone (1 g) was added to the mixture, followed by 2.5 g $NaHB(OAc)_3$ and 1 mL of glacial AcOH. The mixture was stirred at RT overnight. Saturated $NaHCO_3$ solution (50 mL) was added to the mixture and stirred for 1 h. The resulting mixture was separated by separation funnel, the organic layer was extracted once with saturated $NaHCO_3$ solution and once with brine, the resulting organic layer was dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was purified by flash chromatography on silica gel with 9:1 EtOAc:MeOH to afford an orange oil. MS: 290 (M+1). Calc'd. for $C_{16}H_{23}N_3O_2$ - 289.4.

Step F Preparation of 3,3-dimethyl-1-(1-methyl(4-piperidyl))indoline-6-ylamine

3,3-Dimethyl-1-(4-methyl-piperazin-1-yl)-6-nitro-2,3-dihydro-1H-indole (Step E, 600 mg) was dissolved in 20 mL


221

MeOH, the mixture was bubbled with H₂ for 10 min. 10% Pd/C (100 mg) was added and the mixture was stirred under H₂. The mixture was filtered through Celite® and concentrated in vacuo to afford an oil. MS: 260 (M+1). Calc'd. for C₁₆H₂₅N₃ - 259.4.

Step G Preparation of N-[3,3-dimethyl-1-(1-methyl(4-piperidyl))indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was analogously synthesized by the method described in Example 43 from 3,3-dimethyl-1-(1-methyl(4-piperidyl))indoline-6-ylamine (Step F). MS: 496 (M+1). Calc'd. for C₂₉H₃₃N₇O - 495.6.

Example 49



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(1-methyl(4-piperidyl))indol-5-yl]carboxamide

Step A: Preparation of 3-(1-methyl-1,2,3,6-tetrahydro-pyridin-4-yl)-5-nitro-1H-indole

5-Nitroindole (2.6 g) was dissolved in 100 mL anhydrous MeOH, followed by 5 g N-methyl-4-piperidone and NaOMe (5 g) powder. The mixture was heated to reflux under N₂ overnight. The mixture was concentrated in vacuo, and was extracted between saturated NaHCO₃ solution and EtOAc.

222

The resulting organic layer was dried over MgSO_4 , filtered and concentrated in vacuo to afford a yellow solid. This solid was washed with 5 mL EtOAc and 2 mL MeOH to afford a bright yellow solid. MS: 258 (M+1). Calc'd. for $\text{C}_{14}\text{H}_{15}\text{N}_3\text{O}_2$ - 257.29.


Step B: Preparation of 3-(1-methyl-4-piperidyl)indole-5-ylamine:

3-(1-Methyl-1,2,3,6-tetrahydro-pyridin-4-yl)-5-nitro-1H-indole (2.7 g, Step A) was dissolved in 50 mL MeOH, the mixture was bubbled with H_2 for 10 min. 10% Pd/C (150 mg) was added and the mixture was stirred under H_2 overnight. The mixture was then filtered through Celite® and concentrated in vacuo to afford a yellow oil (1.6 g). MS: 230 (M+1). Calc'd. for $\text{C}_{14}\text{H}_{19}\text{N}_3$ - 229.3.

Step C: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[3-(1-methyl(4-piperidyl))indol-5-yl]carboxamide


The title compound was analogously synthesized by the method described in Example 43 from 3-(1-methyl-4-piperidyl)indole-5-ylamine (Step B). MS: 466 (M+1). Calc'd. for $\text{C}_{27}\text{H}_{27}\text{N}_7\text{O}$ - 465.6.

223

Example 50

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(trifluoromethyl)phenyl]carboxamide

MS (ES⁺): 513 (M+H)⁺; (ES⁻): 511. Calc'd C₂₀H₁₄F₃N₅O - 397.4.

Example 51

N-{3-[3-(Dimethylamino)propyl]-5-(trifluoromethyl)phenyl}[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

Step A: Preparation of {3-[3-amino-5-(trifluoromethyl)phenyl]propynyl}dimethylamine

A mixture of 3-bromo-5-trifluoromethylaniline (1.4 g, 5.9 mmol), 1-dimethylamino-2-propyne (1.3 mL, 0.76 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (0.26 g, 0.29 mmol) and CuI (114 mg, 0.60 mmol) in 10 mL of TEA was heated at 100°C in a sealed tube for 3 h. The resulting mixture was filtered over Celite®. The filtrate was concentrated, and the residue was purified by prep-HPLC (reverse phase) to give the aniline. MS (ES+): 243 (M+H)⁺; (ES-): 241 (M-H)⁻. Calc'd $\text{C}_{12}\text{H}_{13}\text{F}_3\text{N}_2$ - 242.24.


Step B: Preparation of {3-[3-amino-5-(trifluoromethyl)phenyl]propyl}dimethylamine

A mixture of {3-[3-amino-5-(trifluoromethyl)phenyl]propyl}dimethylamine (7 g, 29 mmol, Step A) and $\text{Pd}(\text{OH})_2$ (0.5 g) in 250 mL of MeOH was stirred under 50 psi H_2 . After 2 h, the resulting mixture was filtered over Celite®. The filtrate was concentrated, and the residue was diluted with aq. 1N HCl. The aq. layer was washed with Et_2O , made basic with aq. 5N NaOH, and extracted with CH_2Cl_2 . The organic solution was dried over Na_2SO_4 and concentrated to give the titled intermediate. MS (ES+): 386 (M+H)⁺; (ES-): 384 (M-H)⁻. Calc'd $\text{C}_{18}\text{H}_{19}\text{ClF}_3\text{N}_3\text{O}$ - 385.8.

Step C: Preparation of N-{3-[3-(Dimethylamino)propyl]-5-(trifluoromethyl)phenyl}[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was analogously synthesized by the method described in Example 43 from {3-[3-amino-5-(trifluoromethyl)phenyl]propyl}dimethylamine (Step B). MS (ES+): 483 (M+H)⁺; (ES-): 481 (M-H)⁻. Calc'd $\text{C}_{25}\text{H}_{25}\text{F}_3\text{N}_6\text{O}$ - 482.5.

225

Example 52

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[5-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-3-(trifluoromethyl)phenyl]carboxamide

Step A: Preparation of 4,4,5,5-tetramethyl-2-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-1,3,2-dioxaborolane

To a solution of LiHMDS (25 mL, 25 mmol, 1.0 M in THF) in 35 mL of THF was added 1-methyl-4-piperidinone (3.0 mL, 25 mmol) at -78°C. The resulting solution was stirred for 2 h, then Trf_2NPh (8.9 g, 25 mmol) was added. The resulting solution was warmed to RT and stirred for 2 h. The mixture was concentrated, and the residue was purified by alumina (neutral) chromatography to give 1-methyl-4-(1,2,5,6-tetrahydro)pyridyl-(trifluoromethyl) sulfonate. A mixture of above triflate (5.0 g, 20 mmol), bis(pinacolato)diboron (5.6 g, 22 mmol), potassium acetate (6.5 g, 66 mmol), PdCl_2dppf (0.44g, 0.6mmol), and $(\text{dppf})_2$ (0.33g, 0.6 mmol) in 60 mL of dioxane was heated at 80°C for 4 h. The resulting mixture was cooled to RT, diluted with Et_2O (150 mL). The ethereal solution was washed with H_2O followed by brine. The organic

layer dried over Na_2SO_4 , concentrated, and recrystallized in hexane- Et_2O to give the title intermediate.


Step B: Preparation of 5-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-3-(trifluoro-methyl)phenylamine

To a mixture of 4,4,5,5-tetramethyl-2-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-1,3,2-dioxaborolane (1.0 g, 4.4 mmol, Step A), PdCl_2pddf (0.16 g, 0.2 mmol) and K_2CO_3 (1.8g, 13.2 mmol) and 3-amino-5-bromobenzotrifluoride (0.8g, 3.3 mmol) in DMF (25 mL) was heated at 80°C for 16 h. The resulting mixture was diluted with EtOAc , washed with H_2O , dried over Na_2SO_4 , and concentrated. The residue was purified by SiO_2 chromatography to give the title intermediate. MS (ES+): 257 (M+H)⁺. Calc'd $\text{C}_{13}\text{H}_{15}\text{F}_3\text{N}_2$ - 256.3.

Step C: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[5-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-3-(trifluoromethyl)phenyl]carboxamide

The title compound was analogously synthesized by the method described in Example 43 from 5-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-3-(trifluoro-methyl)phenylamine (Step B). MS (ES+): 493 (M+H)⁺; (ES-): 491. Calc'd $\text{C}_{26}\text{H}_{23}\text{F}_3\text{N}_6\text{O}$ - 492.5.

Example 53



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(1-methyl(4-piperidyl))phenyl]carboxamide

Step A: Preparation of 4-phenylpiperidine

4-Cyano-4-phenylpiperidine hydrochloride (10.0 g, 45.0 mmol) was combined with KOH pellets and stirred vigorously under Ar at 160°C for 4 h. The reaction mix was cooled to RT and dissolved into toluene (100 ml) and H₂O (100 ml). After separation of the layers, the aqueous layer was back-extracted two times with toluene. The combined organic layer was dried over Na₂SO₄, concentrated *in vacuo*, and dried under high vacuum, yielding a white solid.

Step B: Preparation of 1-methyl-4-phenylpiperidine

To a stirring mixture at ambient temperature of 4-phenylpiperidine (5.24 g, 32.48 mmol, Step A) in CH₃CN (95 ml) was added a 37% solution of HCHO in H₂O (13 ml). To this mixture was added NaCNBH₃ (3.27 g, 51.97 mmol). AcOH was added dropwise every 10 min over the next h to maintain the reaction pH near 7. The reaction volume was then reduced *in vacuo*. The reaction mix was diluted with CH₂Cl₂ and washed with 2N NaOH and then brine. The crude was concentrated *in vacuo* and eluted through a silica gel column with 10% MeOH/CH₂Cl₂. The 1-methyl-4-phenylpiperidine was concentrated *in vacuo*, yielding a clear oil.

Step C: Preparation of 4-(1-methyl-4-piperidyl)phenylamine

To 1-methyl-4-phenylpiperidine (2.663 g, 15.19 mmol, Step B) was added carefully H₂SO₄ (15.2 ml). The reaction was cooled in an ice bath and a solution of H₂SO₄ (1.66 ml) and fuming HNO₃ (0.67 ml, 15.95 mmol) was added dropwise over 45 min. The mix was stirred at 0°C for 3 h then at RT for 1.5 h before being poured over about 90 g ice and basified with 24 g solid NaOH. The mix was extracted with CH₂Cl₂. The organic layer was washed with H₂O, dried over Na₂SO₄, and concentrated *in vacuo*. The crude was eluted on


228

a silica gel column with a MeOH/CH₂Cl₂ gradient to yield 1-methyl-4-(4-nitrophenyl)piperidine which was hydrogenated under H₂ to furnish the title compound.

Step D: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[4-(1-methyl(4-piperidyl))phenyl]carboxamide

The title compound was analogously synthesized by the method described in Example 43 from 4-(1-methyl-4-piperidyl)phenylamine (Step C). MS: 427.0 (M+1). Calc'd. for C₂₅H₂₆N₆O - 426.5.

Example 54



N-[4-(tert-Butyl)-3-(3-piperidylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

Step A: Preparation of 1-piperidylprop-2-en-1-one

To a 0°C solution of acryloyl chloride (4.576 g, 50.558 mmol) in CH₂Cl₂ (50 ml) was added dropwise and very carefully piperidine (4.305 g, 50.558 mmol). The reaction flask was vented during the exothermic addition. After the addition was completed, the white slurry was stirred at 0°C for 40 min and at RT for 1 h. The reaction was diluted with 70 ml CH₂Cl₂ and washed first with about 60 ml 2N HCl and then with about 60 ml of a mix of 2N NaOH and brine. The organic layer was dried over Na₂SO₄. The solution was evaporated by heating in a water bath at 60°C without

vacuum. Once most solvent had been evaporated off, dried the clear oil under high vacuum at RT for 30 min.

Step B: Preparation of 1-(tert-butyl)-2-bromo-4-nitrobenzene

Bromine (17.4 ml) was added dropwise over 40 min to a stirred mixture of 4-tert-butylnitrobenzene (59.5 g, 332 mmol), silver(II)sulfate (56.5 g, 181 mmol), H₂SO₄ (300 ml), and H₂O (33 ml) at RT. The mixture was stirred for a further 3 h and then poured into 0.1 M Na₂S₂O₅/H₂O (1L). The solid was filtered, washed with H₂O, Et₂O, and CH₂Cl₂. The filtrate layers were separated. The aqueous fraction was extracted with Et₂O. The combined organic layers were combined, dried over Na₂SO₄, and concentrated *in vacuo*. The yellow solid was triturated with hexanes to give a pale yellow crystalline solid.

Step C: Preparation of (2E)-3-[2-(tert-butyl)-5-nitrophenyl]-1-piperidylprop-2-en-1-one

1-(tert-Butyl)-2-bromo-4-nitrobenzene (6.885 g, 26.674 mmol, Step B), 1-piperidylprop-2-en-1-one (4.827 g, 34.677 mmol, Step A), and TEA (7.44 ml, 53.35 mmol) were dissolved into toluene (70 ml). To this solution was added Pd(OAc)₂ (60 mg, 0.267 mmol) and Pd(PPh₃)₄ (617 mg, 0.5335 mmol). The mix was degassed with N₂ and heated in a sealed vessel at 120°C for 15 h. The reaction mixture was cooled to RT, filtered, and concentrated *in vacuo*. The dark crude oil was eluted through a silica gel column with 15% to 22% EtOAc/hexanes gradient system to yield a thick amber oil as the title compound.

Step D: Preparation of 3-(5-amino-2-tert-butylphenyl)-1-piperidin-1-yl-propenone

(2E)-3-[2-(tert-Butyl)-5-nitrophenyl]-1-piperidylprop-2-en-1-one (3.22 g, 10.177 mmol, Step C) was dissolved in

dioxane (20 ml) and IpOH (40 ml). To the N₂-degassed solution was added Pd/C 10% by weight catalyst (2 g). The mix was placed in a Parr hydrogenator and stirred for 18 h under 60 psi H₂. The reaction was not complete the next day, so the reaction was continued for an additional 20 h with fresh catalyst. The mix was filtered through Celite® and concentrated in vacuo to give a foamy oil.


Step E: Preparation of 4-(tert-butyl)-3-(3-piperidylpropyl)phenylamine

3-(5-Amino-2-tert-butylphenyl)-1-piperidin-1-yl-propenone (2.312 g, 7.619 mmol, from above step) was dissolved in THF (100 ml) at RT. To this solution was added LiAlH₄ (434 mg, 11.43 mmol). After the reaction stopped exotherming, it was heated at reflux at about 80°C for 4 h. The reaction mix was cooled to 0°C and treated by dropwise addition of 0.458 ml H₂O, 0.730 ml 10% aqueous NaOH, and 1.19 ml H₂O, respectively. The mix was stirred at RT for 1 h. After 40 min about 3 g of Na₂SO₄ was added. The mix was filtered through Celite® and concentrated in vacuo. The crude was eluted through silica gel column with a gradient system of 95:5 to 90:10 CH₂Cl₂/MeOH, to yield an amber thick oil as the title compound.

Step F: Preparation of N-[4-(tert-butyl)-3-(3-piperidylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


The title compound was analogously synthesized by the method described in Example 43 from 4-(tert-butyl)-3-(3-piperidylpropyl)phenylamine (Step E). MS: 511.4 (M+1). Calc'd. for C₃₁H₃₈N₆O - 510.7.

The aniline preparations for the following Examples 55-57 are similar to Example 54.

Example 55

N-[3-((1E)-4-Pyrrolidinylbut-1-enyl)-4-(tert-butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


MS: 509.4 (M+1). Calc'd. for $C_{31}H_{36}N_6O$ - 508.7.

Example 56

N-[4-(tert-Butyl)-3-(3-pyrrolidinylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


MS: 497.2 (M+1). Calc'd. for $C_{30}H_{36}N_6O$ - 496.7.

232

Example 57

N-[4-(tert-Butyl)-3-(3-morpholin-4-ylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

MS: 513.5 (M+1). Calc'd. for C₃₀H₃₆N₆O₂ - 512.7.

Example 58

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-{3-[3-(4-methylpiperazinyl)-3-oxopropyl]phenyl}carboxamide

Step A: Preparation of 3-(3-nitrophenyl)-1-(4-methylpiperazinyl)propan-1-one

A slurry consisting of CH₂Cl₂ (15 ml), 3-nitrocinnamic acid (3.154 g, 16.329 mmol), 1-methylpiperazine (1.487 g, 14.845 mmol) and EDC (3.557 g, 18.556 mmol) were stirred at RT for 60 h. The reaction was diluted with H₂O and EtOAc. The aqueous layer was back-extracted with EtOAc. The combined organic layers were washed with 2N NaOH and then

brine, dried over Na_2SO_4 , and concentrated *in vacuo*. The crude was eluted through a silica gel column with 5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$, to yield an off-white solid, mostly trans-olefin product.


Step B: Preparation of 3-(3-aminophenyl)-1-(4-methylpiperazinyl)propan-1-one

To a nitrogen-degassed solution of nitro intermediate (3.67 g, 13.330 mmol, Step A) in MeOH (50 ml) was added 10% by weight Pd/C (500 mg). The mix was stirred under H_2 atmosphere for 18 h then filtered through Celite® and concentrated *in vacuo*, yielding a thick amber oil which eventually solidified into a dark pink solid.

Step C: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-{3-[3-(4-methylpiperazinyl)-3-oxopropyl]phenyl}carboxamide

The title compound was analogously synthesized by the method described in Example 54 from 3-(3-aminophenyl)-1-(4-methylpiperazinyl)propan-1-one (Step B). MS: 484.4 (M+1). Calc'd. for $\text{C}_{27}\text{H}_{29}\text{N}_7\text{O}_2$ - 483.6.


Example 59



[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-[3-(4-methylpiperazinyl)-3-oxopropyl]phenyl)carboxamide


234

The title compound was analogously synthesized by the method described in Example 58. MS: 484.4 (M+1). Calc'd. for $C_{27}H_{29}N_7O_2$ - 483.6.

Example 60

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-{3-[3-(4-methylpiperazinyl)propyl]phenyl}carboxamide

The title compound was analogously synthesized by the method described in Example 58 and Step E in Example 54. MS: 470 (M+1). Calc'd. for $C_{27}H_{31}N_7O$ - 469.6.

Example 61


[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-{4-[3-(4-methylpiperazinyl)propyl]phenyl}carboxamide

235

The title compound was analogously synthesized by the method described in Example 58 and Step E in Example 54.

MS: 470 (M+1). Calc'd. for $C_{27}H_{31}N_7O$ - 469.6.

Example 62




[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(2-morpholin-4-ylethyl)indol-6-yl]carboxamide

Step A: Preparation of 1-(2-morpholin-4-ylethyl)indol-6-ylamine

K_2CO_3 (5.08 g, 36.726 mmol) was added to a slurry of 6-nitroindole (1.985 g, 12.242 mmol), 4-(2-chloroethyl)morpholine hydrochloride (2.278 g, 12.242 mmol), and CH_3CN (100 ml). The mix was heated to reflux for 18 h, then cooled to RT, filtered, and concentrated *in vacuo*. The crude was eluted through a silica gel column with a gradient of 3:97 to 5:95 and finally 8:92 MeOH/ CH_2Cl_2 , to yield upon drying the desired intermediate which was hydrogenated under conditions previously described.

Step B: Preparation of [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(2-morpholin-4-ylethyl)indol-6-yl]carboxamide

The title compound was analogously synthesized by the method described in Example 54 from 1-(2-morpholin-4-ylethyl)indole-6-ylamine (Step A). MS: 482.1 (M+1). Calc'd. for $C_{27}H_{27}N_7O_2$ - 481.6.

Example 63

N-[4-(1,1-Dimethyl-3-morpholin-4-ylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

Step A: Preparation of methyl 2-methyl-2-(4-nitrophenyl)propanoate

To a stirred solution of 2-(4-nitrophenyl)propionic acid (9 g, 46 mmol, 1 eq) in MeOH (300 mL) was added HCl (4M in Dioxane, 11.5 mL, 46 mmol, 1 eq). The mixture was stirred at RT overnight and was quenched with aqueous NaHCO₃. The mixture was extracted with EtOAc. The organic layer was dried over MgSO₄ and evaporated under reduced pressure and the partial residue (4.34 g, 20.7 mmol, 1eq) at 0°C in THF (100 mL) was added NaH (1.66g, 41.5 mmol, 2 eq). Mixture was stirred at RT for 1h and CH₃I (2.58 g, 41.5 mmol, 2 eq) was added. Reaction was stirred at RT overnight and was quenched with water. Mixture was extracted with EtOAc. The organic layer was dried over MgSO₄ and evaporated under reduced pressure and used for the next step without further purification to give title compound.

Step B: Preparation of 3-methyl-3-(4-nitrophenyl)butan-1-one

To a stirred solution of methyl 2-methyl-2-(4-nitrophenyl)propanoate (5.32 g, 23.8 mmol, Step A) in THF

(200 mL) at 0°C was added a solution of 1M BH₃ in THF (25.8 mL, 45.8 mmol). The reaction was stirred at RT overnight and was quenched with MeOH. THF was evaporated under reduced pressure and the residue was diluted in EtOAc and aqueous HCl (1M) was added. The mixture was extracted with EtOAc, the organic layer was dried over MgSO₄ and evaporated under reduced pressure. Purification by flash chromatography using 40% EtOAc-hexane gave a yellow solid. To the yellow solid (2.08 g, 10.8 mmol) at 0°C in CH₂Cl₂ was added NMO (1.9 g, 16.1 mmol), molecular sieves 4Å and TPAP (76 mg, 0.2 mmol). The reaction was stirred for 1h and filtered on a silica pad. Solvent was evaporated under reduced pressure, forming the crude aldehyde which was used as is. To a suspension of methoxymethyltriphenylphosphonium chloride (6.4 g, 18.6 mmol) in THF (150 mL) was added a solution of KHMDS 0.5 M in toluene (37 mL, 18.5 mmol). The mixture was stirred for 30 min and crude aldehyde was added. The reaction was stirred at RT for 1h and quenched with H₂O. The mixture was extracted with EtOAc, dried and evaporated under reduced pressure. Et₂O was added and a precipitate formed, which was filtered on a silica pad and rinsed with 40% EtOAc-hexane. The solvent was removed and crude material was dissolved in CH₂Cl₂. A solution of TFA-water (1:1, 10 mL) was added and the reaction was stirred for 2 h at RT. Aqueous NaHCO₃ was added until pH 7 and the mixture was extracted with CH₂Cl₂. The organic layer was dried, filtered and evaporated. Crude compound was purified by flash chromatography (40% EtOAc-hexane) to give the title compound as a yellow oil.

Step C: Preparation of 4-(1,1-dimethyl-3-morpholin-4-ylpropyl)phenylamine

To a stirred solution of aldehyde (509 mg, 2.4 mmol, Step B) and morpholine (0.21 mL, 2.4 mmol) in THF (30 mL)


238

was added $\text{NaBH}(\text{OAc})_3$ (0.73 g, 3.4 mmol). The mixture was stirred at RT overnight and washed with HCl (1M). CH_2Cl_2 was added and the layers were separated. The aqueous layer was basified to pH 9 using NaOH 1M and extracted with CH_2Cl_2 . The organic layer was dried and evaporated the nitro compound. To a solution of the nitro compound (0.50g, 1.8 mmol) in THF (40 mL) was added AcOH (1.97 mmol, 34.5 mmol) followed by zinc (9.1 g, 137 mmol). The mixture was stirred for 1 h, filtered on Celite®, diluted with water and aqueous NaHCO_3 , and the THF layer was evaporated. The residue was extracted with EtOAc, dried and evaporated to give the title compound.

Step D: Preparation of N-[4-(1,1-dimethyl-3-morpholin-4-ylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide

The title compound was analogously synthesized by the method described in Example 54 from 4-(1,1-dimethyl-3-morpholin-4-ylpropyl)phenylamine (Step C). MS: 485.5 (M+1). Calc'd. for $\text{C}_{28}\text{H}_{32}\text{N}_6\text{O}_2$ - 484.6.

Example 64



2-(1H-Indazol-6-ylamino)-N-(4-{2,2,2-trifluoro-1-[2-(2-methoxy-ethoxy)-ethoxy]-1-trifluoromethyl-ethyl}-phenyl)-nicotinamide

Step A: Preparation of 4-{2,2,2-trifluoro-1-[2-(2-methoxy)ethoxy]-1-(trifluoromethyl)ethyl}phenylamine

Diethyl azodicarboxylate (366 mg, 2.1 mmol) was added drop-wise to a solution of 2-(4-aminophenyl)-1,1,1,3,3,3-hexafluoropropan-2-ol (520 mg, 2 mmol), 2-(2-methoxyethoxy)ethan-1-ol (240 mg, 2 mmol) and PPh₃ (550 mg, 2.1 mmol) in THF (10 mL). The mixture was stirred for 2 h, then partitioned between EtOAc and aqueous NaHCO₃ solution. The organic phase was washed with brine. After concentration in vacuo, the organic residue was purified by flash chromatography on silica to give the compound. MS: 362 (M+1). Calc'd. for C₁₄H₁₇F₆NO₃ - 361.29.

Step B: Preparation of 2-fluoropyridine-3-carbonyl chloride


To a solution of 2-fluoropyridine (10 g, 100 mmol) in THF (150 mL) under -78°C was added an LDA solution (2M in heptane/THF/ethylbenzene, 60 mL) dropwise. The mixture was stirred at -78°C for 3 h, then was quenched with a stream of dry CO₂. After warming to RT, the mixture was partitioned between EtOAc (100 mL) and H₂O (200 mL). The aqueous layer was acidified to pH between 3-4, and extracted with EtOAc. The organic solution was collected and washed with brine and dried over Na₂SO₄. After removing the solvent in vacuum, 2-fluoropyridine-3-carboxylic acid was obtained as a brown oil. MS: 140 (M-H). Calc'd. for C₆H₄FNO₂ - 141.10. 2-Fluoropyridine-3-carboxylic acid (7 g) was suspended in SOCl₂ (100 mL). After heating under reflux for 2 h, the mixture became homogeneous. Access SOCl₂ was removed in vacuo to afford a brown solid as desired compound.

Step C: Preparation of [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-(4-{2,2,2-trifluoro-1-[2-(2-methoxyethoxy)ethoxy]-1-(trifluoromethyl)ethyl}phenyl)carboxamide

To a suspension of 2-fluoropyridine-3-carbonyl chloride (180 mg, 0.5 mmol, Step B) and NaHCO₃ (300 mg) in CH₂Cl₂ (5 ml), a solution of 4-{2,2,2-trifluoro-1-[2-(2-methoxyethoxy)ethoxy]-1-(trifluoromethyl)ethyl}phenylamine (95 mg) was added dropwise, and the suspension was stirred at RT for 1.5 h. Solid inorganic salts were removed via filtration. The filtrate was concentrated to afford (2-fluoro(3-pyridyl))-N-(4-{2,2,2-trifluoro-1-[2-(2-methoxyethoxy)ethoxy]-1-(trifluoromethyl)ethyl}phenyl)-carboxamide as a brown solid. (2-Fluoro(3-pyridyl))-N-(4-{2,2,2-trifluoro-1-[2-(2-methoxyethoxy)ethoxy]-1-(trifluoromethyl)ethyl}-phenyl)carboxamide (240 mg, 0.5 mmol) and 6-amino-indazole (133 mg, 1 mmol) were dissolved in DMSO (2 mL). The solution was heated to 120°C for 8 h, then partitioned between CH₂Cl₂ and aqueous NaHCO₃ (sat.). The organic layer was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified via preparative HPLC to give a yellow powder as a TFA salt of the title compound. MS: 598 (M+1). Calc'd. for C₂₇H₂₅F₆N₅O₄ - 597.5.


The following compounds (Examples 65-68) were analogously synthesized by the method described in Example 64. Detailed intermediate preparations were described.

241

Example 65

[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-{4-[2,2,2-trifluoro-
1-(2-piperidylethoxy)-1-(trifluoromethyl)ethyl]phenyl}carboxamide

MS: 607 (M+1). Calc'd. for $C_{29}H_{28}F_6N_6O_2$ - 606.6.

Example 66

N-[4-(tert-Butyl)phenyl][6-fluoro-2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


Step A: Preparation of (2,6-difluoro(3-pyridyl))-N-[4-(tert-butyl)phenyl]-carboxamide

A solution of 2,6-difluoropyridine-3-carboxylic acid (prepared similar to that described for 2-fluoropyridine-3-carboxylic acid, Example 64) (3.2 g, 20 mmol), *t*-butylaniline 11 (3.0 g, 20 mmol), HOBT (2.6 g, 20 mmol), EDAC (8 g, 40 mmole), and DIEA (8 mL) in CH₂Cl₂ (80 mL) was stirred at RT for 1 h. The mixture was washed with aq. NaHCO₃ and brine. The organic solution was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified via flash chromatography on silica (Hex:EtOAc = 4:1) to give a light yellow flaky crystal as the desired product.

Step B: Preparation of N-[4-(tert-butyl)phenyl][6-fluoro-2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide


The title compound was analogously synthesized by the method described in Example 64 from (2,6-difluoro(3-pyridyl))-N-[4-(tert-butyl)phenyl]-carboxamide (Step A).
MS: 404 (M+1). Calc'd. for C₂₃H₂₂FN₅O - 403.5.

Example 67




[6-Fluoro-2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylethyl)phenyl]carboxamide

243

MS: 390 (M+1). Calc'd. for $C_{22}H_{20}FN_5O$ - 389.4.Example 68

[6-Fluoro-2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[3-(trifluoromethyl)phenyl]carboxamide

MS: 416 (M+1). Calc'd. for $C_{20}H_{13}F_4N_5O$ - 415.3.Example 69

{2-[(1-(2-Pyridyl)pyrrolidin-3-yl)amino](3-pyridyl)}-N-[3-(trifluoromethyl)phenyl]carboxamide

Step A: Preparation of 1-(2-pyridyl)pyrrolidine-3-ylamine


2-Fluoropyridine (2 g, 0.02 mol) and Boc-aminopyrrolidine (3.6 g, 0.02 mol) were heated neat at 120°C

244

for 2 h. The reaction was cooled, 4N HCl/Dioxane (100 mg) was added and the reaction was stirred at RT for 4 h. The solvent was evaporated and 2 N NaOMe/MeOH was added to adjust to basic pH. The precipitate was filtered and the filtrate was evaporated. The residue was dissolved in CH₂Cl₂, the solution was filtered and the filtrate was evaporated to give the crude material, was used in next step without further purification.

Step B: Preparation of {2-[(1-(2-pyridyl)pyrrolidin-3-yl)amino](3-pyridyl)}-N-[3-(trifluoromethyl)phenyl]-carboxamide

A mixture of (2-chloro(3-pyridyl))-N-(3-trifluoromethylphenyl)carboxamide and 1-(2-pyridyl)pyrrolidine-3-ylamine (Step A) was heated at 130°C neat for 3 h. The reaction was cooled and diluted with CH₂Cl₂ and washed with H₂O twice followed by brine. The organic layer was dried with Na₂SO₄ and evaporated under reduced pressure. The residue was purified by column chromatography with EtOAc and further mixed with MeOH and 1 N HCl/Et₂O (2 ml). The solution was evaporated to furnish the titled compound. MS (ES⁺): 428 (M+H); (ES⁻): 426 (M-H). Calc'd. for C₂₂H₂₀F₃N₅O - 427.2.


Example 70

2-(1H-Indazol-6-ylamino)-N-[3-(3-morpholin-4-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide

2-Chloro-N-[3-(3-morpholin-4-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide (2.95 g) was dissolved in a small volume of IpOH. 6-amino indazole (2.75 g) and 0.53 mL of TFA were added. The reaction mixture was heated to 155°C in an open flask for 3.5 h. After cooling to RT the residue was dissolved in 2 N HCl. The aqueous solution was extracted 2 times with CH₂Cl₂, basified to pH 12 by the addition of Na₂CO₃ and 1 N NaOH. The solution was extracted 6 times with CH₂Cl₂ to isolate the product. The combined 5 extractions were dried over Na₂SO₄, filtered and stripped. The crude product was purified by silica gel chromatography eluting with a step gradient of 3-5% MeOH:CH₂Cl₂ to yield the titled compound. MS: 525 (M+1). Calc'd for C₂₇H₂₇F₃N₆O₂ - 524.5.


246

The following compounds (Examples 71-95) were synthesized by the method described in Example 70 unless specifically described.

Example 71

2-(1H-Indazol-6-ylamino)-N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide

MS: 523.5 (M+1). Calc'd. for $C_{28}H_{29}F_3N_6O$ - 522.6.


Example 72

247

2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-piperidin-4-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide


MS: 509 (M+1). Calc'd. for $C_{27}H_{27}F_3N_6O$ - 508.5.

Example 73



2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide


Example 74



2-(1H-Indazol-6-ylamino)-N-[3-(piperidin-4-yloxy)-5-trifluoromethyl-phenyl]-nicotinamide

MS: 497 (M+1). Calc'd. for $C_{25}H_{23}F_3N_6O$ - 496.5


Example 75



2-(1H-Indazol-6-ylamino)-N-[3-(piperidin-4-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide

MS (ES+): 511 (M+H), Calc'd for $C_{26}H_{25}F_3N_6O_2$ - 510.52.


Example 76



N-(3,3-Dimethyl-1,1-dioxo-2,3-dihydro-1H-116-benzo[d]isothiazol-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide


249

MS (ES+): 449 (M+H)+; (ES-): 447 (M-H). Calc'd. for $C_{22}H_{20}N_6O_3S$ - 448.

Example 77


2-(1H-Indazol-6-ylamino)-N-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-naphthalen-2-yl)-nicotinamide

MS (ES+): 440 (M+H)+; (ES-): 438 (M-H). Calc'd. for $C_{27}H_{29}N_5O$ - 439.


Example 78

2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide

250


M+H 575; Calc'd. for $C_{28}H_{27}F_5N_6O_2$ - 574.Example 79

2-(1H-Indazol-6-ylamino)-N-[3-(1-isopropyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide


M+H 603; Calc'd. for $C_{30}H_{31}F_5N_6O_2$ - 602.Example 80

N-[3-(2-Hydroxy-3-pyrrolidin-1-yl-propoxy)-4-pentafluoroethyl-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

251


M+H 591; Calc'd. for $C_{28}H_{27}F_5N_6O_3$ - 590.Example 81

2-(1H-Indazol-6-ylamino)-N-[4-pentafluoroethyl-3-(2-piperidin-1-yl-ethoxy)-phenyl]-nicotinamide


M+H 575; Calc'd. for $C_{28}H_{27}F_5N_6O_2$ - 574.Example 82

N-[3-(2-Hydroxy-3-pyrrolidin-1-yl-propoxy)-4-pentafluoroethyl-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

252

M+H 591; Calc'd. for $C_{28}H_{27}F_5N_6O_3$ - 590.Example 83

2-(1H-Indazol-6-ylamino)-N-[4-pentafluoroethyl-3-(2S-pyrrolidin-2-ylmethoxy)-phenyl]-nicotinamide


M+H 547; Calc'd. for $C_{26}H_{23}F_5N_6O_2$ - 546.Example 84

2-(1H-Indazol-6-ylamino)-N-[4-pentafluoroethyl-3-(2R-pyrrolidin-2-ylmethoxy)-phenyl]-nicotinamide

M+H 547; Calc'd. for $C_{26}H_{23}F_5N_6O_2$ - 546.

253


Example 85



2-(1H-Indazol-6-ylamino)-N-[3-(pyrrolidin-2-ylmethoxy)-4-trifluoromethyl-phenyl]-nicotinamide

M+H 497; Calc'd. for $C_{25}H_{23}F_3N_6O_2$ - 496.


Example 86



2-(1H-Indazol-6-ylamino)-N-[3-(2-pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenyl]-nicotinamide


M+H 511; Calc'd. for $C_{26}H_{25}F_3N_6O_2$ - 510.

254

Example 87


N-(1-Acetyl-3,3-dimethyl-2,3-dihydro-1H-indol-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide

M+H 441.3; Calc'd for $C_{25}H_{24}N_6O_2$: 440.2.

Example 88


2-(1H-Indazol-6-ylamino)-N-{4-[1-methyl-1-(1-methylpiperidin-4-yl)-ethyl]-phenyl}-nicotinamide

255

MS (ES+): 469 (M+1)+, Calc'd for $C_{28}H_{32}N_6O$ - 468.26Example 89

N-(4-Acetyl-2,2-dimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide


MS (ES+): 457.5 (M+H)+; (ES-) 455.5 (M-H). Calc'd for $C_{25}H_{24}N_6O_3$ - 456.5

Example 90

2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-piperidin-4-yl)-5-trifluoromethyl-phenyl]-nicotinamide


MS (ES+): 495.1 (M+H)+; (ES-) 493.2 (M-H). Calc'd for $C_{26}H_{25}F_3N_6O$: 494.20.

256

Example 91

N-(3-Bromo-5-trifluoromethyl-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide


MS (ES⁺): 475.9 (M+H)⁺; (ES⁻) 474.0 (M-H). Calc'd for C₂₀H₁₃BrF₃N₅O: 475.30.

Example 92

2-(1H-Indazol-6-ylamino)-N-(2,2,4-trimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-nicotinamide


MS (ES⁺): 429.2 (M+H)⁺; (ES⁻) 427.4 (M-H). Calc'd for C₂₄H₂₄N₆O₂: 428.2.

257

Example 93

N-[4-tert-Butyl-3-(piperidin-4-ylmethoxy)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide


M+H 499: Calc'd for $C_{29}H_{34}N_6O_2$: 498.

Example 94

N-(2-Acetyl-4,4-dimethyl-1,2,3,4-tetrahydro-isoquinolin-7-yl)-2-(1H-indazol-6-ylamino)-nicotinamide


M+H 455.3. Calc'd for $C_{26}H_{26}N_6O_2$: 454.2.

258

Example 95

1-Boc-2-(2-tert-Butyl-5-{[2-(1H-indazol-6-ylamino)-pyridine-3-carbonyl]-amino}-phenoxy)methyl)-pyrrolidine


M+H 585; Calc'd for C₃₃H₄₀N₆O₄: 584.

Example 96

N-[4-tert-Butyl-3-(pyrrolidin-2-ylmethoxy)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

259


The title compound was deprotected by a method analogous to that described in Example 69, step a. M+H 485; Calc'd for $C_{28}H_{32}N_6O_2$: 484.

Example 97

2-(1H-Indazol-6-ylamino)-N-[3-(2S-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide


The title compound was deprotected by a method analogous to that described in Example 69, step a. MS: 497 (M+1). Calc'd. for $C_{25}H_{23}F_3N_6O_2$ - 496.5.

260

Example 98

N-(4-tert-Butyl-3-piperazin-1-yl-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide

The title compound was deprotected by a method analogous to that described in Example 69, step a. MS (ES⁺): 470 (M+H)⁺; (ES⁻): 468 (M-H). Calc'd. for C₂₇H₃₁N₇O - 469.


Example 99

N-(3,3-dimethyl-2,3-dihydro-1H-indol-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide

261


N-(1-Acetyl-3,3-dimethyl-2,3-dihydro-1H-indol-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide (500 mg) was deprotected heating with 12 N HCl (3 ml) and EtOH (3 ML) at reflux for 1 h. Dried under vacuum and purified via HPLC. M+H 399.3; Calc'd for $C_{23}H_{22}N_6O$: 398.2.

Example 100




N-(3,3-Dimethyl-1-piperidin-4-yl)-2,3-dihydro-1H-indol-6-yl)-
2-(1H-indazol-6-ylamino)-nicotinamide

The title compound was deprotected by a method analogous to that described in Example 99. MS (M+H) = 482.4; Calc'd for $C_{28}H_{31}N_7O$: 481.2.

Example 101

N-(2,2-Dimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide

The title compound was deprotected by a method analogous to that described in Example 99. MS (ES⁺): 415.3 (M+H)⁺; (ES⁻) 413.2 (M-H). Calc'd for C₂₃H₂₂N₆O₂ - 414.5.


Example 102

N-[4-tert-Butyl-3-(4-propyl-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

263

N-[4-tert-Butyl-3-(4-propyl-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide was alkylated by a method analogous to that described in Example 63, step c. MS (ES+): 512 (M+H)+; (ES-): 510 (M-H). Calc'd. for $C_{30}H_{37}N_7O$ - 511.


Example 103



N-[4-tert-Butyl-3-(4-isopropyl-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide


N-[4-tert-Butyl-3-(4-propyl-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide was alkylated by a method analogous to that described in Example 63, step c. MS (ES+): 512 (M+H)+; (ES-): 510 (M-H). Calc'd. for $C_{30}H_{37}N_7O$ - 511.

264

Example 104

2-(1H-Indazol-6-ylamino)-N-[3-(1-methylpyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide


The title compound was prepared by a method analogous to that described in Example 70. M+H 511.4. Calc'd for $C_{26}H_{25}F_3N_6O_2$: 510.5.

Example 105

N-(4,4-Dimethyl-1-oxo-1,2,3,4-tetrahydro-isoquinolin-7-yl)-2-(1H-indazol-6-ylamino)-nicotinamide


The title compound was prepared by a method analogous to that described in Example 70. (M+H) 427, Calc. for $C_{24}H_{22}N_6O_2$: 426.2.

265

Example 106

N-(3,3-Dimethyl-2,3-dihydro-benzofuran-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide

The title compound was prepared by a method analogous to that described in Example 70. MS: 400 (M+1). Calc'd. for $C_{23}H_{21}N_5O_2$ -399.4.


Example 107

266

N-[1-(2-Dimethylamino-acetyl)-3,3-dimethyl-2,3-dihydro-1H-indol-6-yl]-2-(1H-indazol-6-ylamino)-nicotinamide

The title compound was prepared by a method analogous to that described in Example 70. M+H 484.3; Calc'd for $C_{27}H_{29}N_7O_2$: 483.2.


Example 108



2-(1H-Indazol-6-ylamino)-N-[3-(4-methyl-piperazin-1-ylmethyl)-4-pentafluoroethyl-phenyl]-nicotinamide

The title compound was prepared by a method analogous to that described in Example 70. M+H 560.4. Calc'd for $C_{27}H_{26}F_5N_7O$: 559.2.

Example 109




267

2-(1H-Indazol-6-ylamino)-N-[3-(4-Boc-piperazin-1-ylmethyl)-4-pentafluoroethyl-phenyl]-nicotinamide

The title compound was prepared by a method analogous to that described in Example 70. M+H 646.4. Calc'd for $C_{31}H_{32}F_5N_7O_3$: 645.2.


Example 110



2-(1H-Indazol-6-ylamino)-N-(3-morpholin-4-ylmethyl-4-pentafluoroethyl-phenyl)-nicotinamide


The title compound was prepared by a method analogous to that described in Example 70. M+H 547.1. Calc'd for $C_{26}H_{23}F_5N_5O_2$: 546.2.

268

Example 111

2-(1H-Indazol-6-ylamino)-N-(4-pentafluoroethyl-3-piperazin-1-ylmethyl-phenyl)-nicotinamide

The title compound was prepared by a method analogous to that described in Example 69, step a. M+H 546.4. Calc'd for $C_{26}H_{24}F_5N_7O$: 545.2.


Example 112

N-[4-tert-Butyl-3-(4-Boc-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

269

The title compound was prepared by a method analogous to that described in Example 1, Step B. (ES+) 570 (M + 1)⁺; (ES-): 568 (M - 1)⁻. Calc'd. C₃₂H₃₉N₇O₃ - 569.7.


Example 113



**N-(4-tert-Butyl-3-nitro-phenyl)-2-(1H-indazol-6-ylamino)-
nicotinamide**

The title compound was prepared from N-(4-tert-butyl-3-nitro-phenyl)-2-chloro-nicotinamide by a method analogous to that described in Example 70, step E. MS (ES+): 401 (M+H), Calc'd for C₂₃H₂₄N₆O - 400.5.

Example 114




270

**N-(3-Amino-4-tert-butyl-phenyl)-2-(1H-indazol-6-ylamino)-
nicotinamide**

The title compound was prepared from N-(4-tert-butyl-3-nitro-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide by dissolving in EtOH (20 ml) and adding Sn(II)Cl₂ (12.58 g) and stirred at RT for 5 h and at 0°C overnight. The mix was warmed to RT and quenched by pouring onto ice, neutralized with NaHCO₃ and basified with 6N NaOH. An emulsion formed upon addition of EtOAc which was filtered through Celite and the aqueous layer was extracted with EtOAc. The combined organic layers were washed with H₂O, brine and dried over MgSO₄ then purified by flash chromatography (3-5% MeOH to afford the product as a orange-yellow solid.


Example 115



**N-[4-tert-Butyl-3-(2-hydroxy-ethylamino)-phenyl]-2-(1H-
indazol-6-ylamino)-nicotinamide**

The title compound was prepared from N-(3-amino-4-tert-butyl-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide by a method analogous to that described in Example 42. MS (ES⁺): 445 (M+H), Calc'd for C₂₅H₂₈N₆O₂ - 444.5.

271

Example 116

N-[4-tert-Butyl-3-(2-morpholin-4-yl-ethylamino)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

Step a N-[3-(2-Bromo-ethylamino)-4-tert-butyl-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

The title compound was prepared from N-[4-tert-butyl-3-(2-hydroxy-ethylamino)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide by dissolving in CH₂Cl₂ (5 ml) and adding CBr₄ (215 mg) and bis(diphenylphosphino)-propane (270mg). The reaction was stirred at RT for 3 h then worked up with partitioning between H₂O and CH₂Cl₂. The aqueous layer was extracted with CH₂Cl₂ (3x) and the combined organic fractions were washed with H₂O, brine and dried (MgSO₄). The crude material was purified by flash chromatography to yield a yellow foamed solid. M+H 508.


Step B N-[4-tert-Butyl-3-(2-morpholin-4-yl-ethylamino)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide

N-[3-(2-Bromo-ethylamino)-4-tert-butyl-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide was dissolved in DMF (2 ml)

272

morpholine (.09 ml) was added and stirred overnight at RT. The mixture was partitioned between sat'd NaHCO₃ and EtOAc. The aqueous layer was extracted with EtOAc and the organic layers were washed with H₂O, brine and dried (MgSO₄). The crude material was purified by flash chromatography (5% MeOH/ CH₂Cl₂) to yield a yellow foamed solid. MS (ES⁺): 514 (M+H), Calc'd for C₂₉H₃₅N₇O₂ - 513.6.


Example 117



N-[4-tert-Butyl-3-(1-Boc-piperidin-4-ylamino)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide


The title compound was prepared from N-(3-amino-4-tert-butyl-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide and Boc-piperidin-4-one by a method analogous to that described in Example 42. MS (ES⁺): 584 (M+H), Calc'd for C₃₃H₄₁N₇O₃ - 583.7.

273

Example 118

2-(1H-Indazol-6-ylamino)-N-[2-(2-morpholin-4-yl-ethyl)-
1,2,3,4-tetrahydro-isoquinolin-7-yl]-nicotinamide


498 (M+1); 497 (M-1). Calc'd. for $C_{28}H_{31}N_7O_2$ - 497.6.

Example 119

N-[4-tert-Butyl-2-(4-methyl-piperazin-1-yl)-phenyl]-2-(1H-
indazol-6-ylamino)-nicotinamide


MS (ES+): 484 (M+H)⁺; (ES-): 482 (M-H). Calc'd. for
 $C_{28}H_{33}N_7O$ - 483.3.

274

Example 120

2-(1H-Indazol-6-ylamino)-N-(2-oxo-4-trifluoromethyl-2H-chromen-7-yl)-nicotinamide


M+H 466. Calc'd for $C_{23}H_{14}F_3N_5O_3$: 465.1.

Example 121

2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-5-trifluoromethyl-phenyl]-nicotinamide


M+H, 493; M-H, 491; Calc'd for $C_{26}H_{23}F_3N_6O$: 492.

275

Example 122

2-(1H-Indazol-6-ylamino)-N-(1H-indol-7-yl)-nicotinamide


MS: 369 (M+1). Calc'd. for $C_{21}H_{16}N_6O$ -368.4.

Example 123

2-(1H-Indazol-6-ylamino)-N-(4-pentafluoroethyl-phenyl)-
nicotinamide


M+H 448. Calc'd for $C_{21}H_{14}F_5N_5O$: 447.

276

Example 124

N-[4-tert-Butyl-3-(piperidin-4-ylamino)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide


MS (ES⁺): 484 (M+H), Calc'd for C₂₈H₃₃N₇O - 483.6.

Example 125

2-(1H-Indazol-6-ylamino)-N-(3-piperazin-1-ylmethyl-5-trifluoromethyl-phenyl)-nicotinamide


M+H 496.3; Calc'd for C₂₅H₂₄F₃N₇O: 495.2.

277

Example 126


N-(4,4-Dimethyl-1,2,3,4-tetrahydro-isoquinolin-7-yl)-2-(1H-indazol-6-ylamino)-nicotinamide

M+H 413.4. Calc'd for $C_{24}H_{24}N_6O$: 412.2.

Example 127


N-(1-Boc-4,4-Dimethyl-1,2,3,4-tetrahydro-isoquinolin-7-yl)-2-(1H-indazol-6-ylamino)-nicotinamide

M+H 513. Calc'd for $C_{29}H_{32}N_6O_3$: 512.2.

Example 128

2-(1H-Indazol-6-ylamino)-N-[3-(4-Boc-piperazin-1-ylmethyl)-
5-trifluoromethyl-phenyl]-nicotinamide

M+H 596.4; Calc'd for C₃₀H₃₂F₃N₇O₃: 595.2.

Example 129


N-[3-(1-Methyl-pyrrolidin-2-ylmethoxy)-5-
trifluoromethyl-phenyl]-2-(4-oxazol-5-yl-phenylamino)-
nicotinamide

M+H 538.2. Calc'd for C₂₈H₂₆F₃N₅O₃: 537.2.

Other compounds included in this invention are set forth in Tables 1-4 below.

- 279 -


Table 1.



5	#	R ¹	R ²
	130.	2-chlorophenyl	H
	131.	4-benzimidazolyl	H
	132.	5-benzimidazolyl	H
	133.	7-benzimidazolyl	H
10	134.	2-chlorophenyl	5-F
	135.	3-isoquinolinyl	5-F
	136.	2-quinolinyl	5-F
	137.	2-benzthiazolyl	5-F
	138.	2-benzimidazolyl	5-F
15	139.	4-benzimidazolyl	5-F
	140.	5-benzimidazolyl	5-F
	141.	6-benzimidazolyl	5-F
	142.	7-benzimidazolyl	5-F
	143.	4-chlorophenyl	3-pyridyl
20	144.	4-pyridyl	H
	145.	4-pyridyl	6-CH ₃
	146.	4-chlorophenyl-	5-Cl
	147.	3,4-dichlorophenyl-	5-F
	148.	4-fluorophenyl	6-CH ₃
25	149.	3-chlorophenyl	6-F
	150.	3-fluorophenyl	6-F
	151.	3-fluoro-4-methoxyphenyl	6-CH ₃
	152.	3-fluoro-4-methylphenyl	6-Cl

- 280 -


Table 1. (cont.)



5	#	R ¹	R ²
	153.	4-phenoxyphenyl	H
	154.	3-phenoxyphenyl	H
	155.	4-biphenyl	H
	156.	4-cyclohexylphenyl	H
10	157.	2-quinolyl	H
	158.	3-isoquinolyl	H
	159.	3-quinolyl	H
	160.	1-isoquinolyl	H
	161.	5-quinolyl	H
15	162.	5-isoquinolyl	H
	163.	6-quinolyl	H
	164.	6-isoquinolyl	H
	165.	7-quinolyl	H
	166.	7-isoquinolyl	H
20	167.	4-quinolyl	H
	168.	4-isoquinolyl	H
	169.	4-pyridyl	H
	170.	4-pyrimidinyl	H
	171.	2-pyrimidinyl	H
25	172.	6-pyrimidinyl	H
	173.	4-pyridazinyl	H
	174.	5-pyridazinyl	H
	175.	4-indolyl	H
	176.	5-isoindolyl	H
30	177.	5-naphthyridinyl	H
	178.	6-quinozaliny	H

- 281 -


Table 1. (cont.)



5	#	R ¹	R ²
	179.	6-isoquinolyl	H
	180.	4-naphthyridinyl	H
	181.	5-quinozalinyl	H
	182.	4-naphthyridinyl	H
10	183.	7-tetrahydroquinolinyl	H
	184.	6-indazolyl	H
	185.	6-isoindolyl	H
	186.	5-indazolyl	H
	187.	5-isoindolyl	H
15	188.	6-benzothienyl	H
	189.	6-benzofuryl	H
	190.	5-benzothienyl	H
	191.	5-benzofuryl	H
	192.	2-benzimidazolyl	H
20	193.	2-benzoxazolyl	H
	194.	2-benzthiazolyl	H
	195.	6-benzimidazolyl	H
	196.	6-benzoxazolyl	H
	197.	6-benzthiazolyl	H
25	198.	2-quinazolinyl	H
	199.	3-(phenoxy)-6-pyridyl	H
	200.	4-(phenylcarbonyl)phenyl	H
	201.	4-(phenylamino)phenyl	H
	202.	4-cyclohexyloxyphenyl	H

- 282 -


Table 1. (cont.)



5	#	R ¹	R ²
	203.	4-(3-thienyl)phenyl	H
	204.	4-(pyrazol-3-yl)phenyl	H
	205.	4-chlorophenyl	6-Cl
	206.	4-pyridyl	6-Cl
10	207.	3-methoxyphenyl	6-Cl
	208.	4-hydroxyphenyl	6-Cl
	209.	3-hydroxyphenyl	H
	210.	2-hydroxyphenyl	H
	211.	4-chlorophenyl	6-phenyl
15	212.	4-phenoxyphenyl	6-F
	213.	4-biphenyl	6-phenyl
	214.	4-hydroxyphenyl	6-phenyl
	215.	4-cyclohexylphenyl	6-F
	216.	3-isoquinolyl	6-phenyl
20	217.	4-piperidinylmethylphenyl	H
	218.	4-morpholinylmethylphenyl	H

- 283 -

Table 2a.




5

#	R ¹
219.	4-chlorophenyl
220.	3,4-dichlorophenyl
10 221.	4-phenoxyphenyl
222.	4-biphenyl
223.	4-cyclohexylphenyl
224.	3-isoquinolyl

15


Table 2b.



#	R ¹
225.	4-chlorophenyl
226.	3,4-dichlorophenyl
227.	4-phenoxyphenyl
25 228.	4-biphenyl
229.	4-cyclohexylphenyl
230.	3-isoquinolyl

- 284 -

Table 2c.




5	#	R¹	A⁵
	231.	4-chlorophenyl	NH
	232.	3,4-dichlorophenyl	NH
10	233.	4-phenoxyphenyl	NH
	234.	4-biphenyl	NH
	235.	4-cyclohexylphenyl	NH
	236.	3-isoquinolyl	NH
	237.	4-chlorophenyl	O
15	238.	3,4-dichlorophenyl	O
	239.	4-phenoxyphenyl	O
	240.	4-biphenyl	O
	241.	4-cyclohexylphenyl	O
	242.	3-isoquinolyl	O
20	243.	3,4-dichlorophenyl	S
	244.	4-phenoxyphenyl	S
	245.	4-biphenyl	S
	246.	4-cyclohexylphenyl	S
	247.	3-isoquinolyl	S

25

- 285 -


Table 2d.



5		
	#	A ^S
	248. 4-chlorophenyl	NH
	249. 3,4-dichlorophenyl	NH
	250. 4-phenoxyphenyl	NH
10	251. 4-biphenyl	NH
	252. 4-cyclohexylphenyl	NH
	253. 3-isoquinolyl	NH
	254. 4-chlorophenyl	O
	255. 3,4-dichlorophenyl	O
15	256. 4-phenoxyphenyl	O
	257. 4-biphenyl	O
	258. 4-cyclohexylphenyl	O
	259. 3-isoquinolyl	O
	260. 3,4-dichlorophenyl	S
20	261. 4-phenoxyphenyl	S
	262. 4-biphenyl	S
	263. 4-cyclohexylphenyl	S
	264. 3-isoquinolyl	S

- 286 -

Table 2e.




5	#	R ¹	A ⁵
	265.	4-chlorophenyl	NH
	266.	3,4-dichlorophenyl	NH
10	267.	4-phenoxyphenyl	NH
	268.	4-biphenyl	NH
	269.	4-cyclohexylphenyl	NH
	270.	3-isoquinolyl	NH
	271.	4-chlorophenyl	O
15	272.	3,4-dichlorophenyl	O
	273.	4-phenoxyphenyl	O
	274.	4-biphenyl	O
	275.	4-cyclohexylphenyl	O
	276.	3-isoquinolyl	O
20	277.	3,4-dichlorophenyl	S
	278.	4-phenoxyphenyl	S
	279.	4-biphenyl	S
	280.	4-cyclohexylphenyl	S
	281.	3-isoquinolyl	S

25

- 287 -

Table 2f.




5			A ⁵
	#	R ¹	
	282.	4-chlorophenyl	NH
	283.	3,4-dichlorophenyl	NH
	284.	4-phenoxyphenyl	NH
10	285.	4-biphenyl	NH
	286.	4-cyclohexylphenyl	NH
	287.	3-isoquinolyl	NH
	288.	4-chlorophenyl	O
	289.	3,4-dichlorophenyl	O
15	290.	4-phenoxyphenyl	O
	291.	4-biphenyl	O
	292.	4-cyclohexylphenyl	O
	293.	3-isoquinolyl	O
	294.	3,4-dichlorophenyl	S
20	295.	4-phenoxyphenyl	S
	296.	4-biphenyl	S
	297.	4-cyclohexylphenyl	S
	298.	3-isoquinolyl	S

25

- 288 -

Table 2g.




5	#	R ¹	A ⁵
	299.	4-chlorophenyl	NH
	300.	3,4-dichlorophenyl	NH
	301.	4-phenoxyphenyl	NH
10	302.	4-biphenyl	NH
	303.	4-cyclohexylphenyl	NH
	304.	3-isoquinolyl	NH
	305.	4-chlorophenyl	O
	306.	3,4-dichlorophenyl	O
15	307.	4-phenoxyphenyl	O
	308.	4-biphenyl	O
	309.	4-cyclohexylphenyl	O
	310.	3-isoquinolyl	O

20

- 289 -


Table 2h.



5	#	R ¹	A ⁵
	311.	4-chlorophenyl	NCH ₃
	312.	3,4-dichlorophenyl	NCH ₃
	313.	4-phenoxyphenyl	NH
	314.	4-biphenyl	NH
10	315.	4-cyclohexylphenyl	NH
	316.	4-tertbutylphenyl	NCH ₃
	317.	4-chlorophenyl	O
	318.	3,4-dichlorophenyl	O
	319.	4-phenoxyphenyl	O
15	320.	4-biphenyl	O
	321.	4-cyclohexylphenyl	O
	322.	3-isoquinolyl	O

- 290 -


Table 3.



5	#	R ¹	R ²
	323.	4-chlorophenyl	5-Br
	324.	3-chlorophenyl	H
	325.	3-chlorophenyl	5-Br
10	326.	3-isoquinolyl	H
	327.	3-isoquinolyl	5-Br
	328.	4-phenoxyphenyl	5-Br
	329.		5-Br
	330.		H
15	331.		5-Br
	332.		5-Br
	333.	4-chlorophenyl	6-Cl
	334.	3,4-dichlorophenyl	H
	335.	4-fluorophenyl	H
20	336.	3-fluorophenyl	H
	337.	3-fluoro-4-methoxyphenyl	H
	338.	3-fluoro-4-methylphenyl	H


- 291 -

Table 3. cont.



5			R ²
	#	R ¹	
	339.	3-phenoxyphenyl	H
	340.	4-cyclohexylphenyl	H
	341.	2-quinolyl	H
10	342.	3-quinolyl	H
	343.	1-isoquinolyl	H
	344.	5-quinolyl	H
	345.	5-isoquinolyl	H
	346.	6-quinolyl	H
15	347.	6-isoquinolyl	H
	348.	7-quinolyl	H
	349.	7-isoquinolyl	H
	350.	4-quinolyl	H
	351.	4-isoquinolyl	H
20	352.	4-pyridyl	H
	353.	4-pyrimidinyl	H
	354.	2-pyrimidinyl	H
	355.	6-pyrimidinyl	H
	356.	4-pyridazinyl	H
25	357.	5-pyridazinyl	H
	358.	4-indolyl	H
	359.	5-isoindolyl	H
	360.	5-naphthyridinyl	H
	361.	6-quinozalinyl	H
30	362.	6-isoquinolyl	H
	363.	4-naphthyridinyl	H


Table 3. (cont.)



5			R ²
	#	R ¹	
	364.	5-quinozaliny1	H
	365.	4-naphthyridiny1	H
	366.	tetrahydroquinoliny1	H
10	367.	6-indazolyl	H
	368.	6-isoindolyl	H
	369.	5-indazolyl	H
	370.	5-isoindolyl	H
	371.	6-benzothienyl	H
15	372.	6-benzofury1	H
	373.	5-benzothienyl	H
	374.	5-benzofury1	H
	375.	6-benzthiazolyl	H
	376.	2-quinazoliny1	H
20	377.	3-(phenoxy)-6-pyridyl	H
	378.	4-(phenylcarbonyl)phenyl	H
	379.	4-(phenylamino)phenyl	H
	380.	cyclohexyloxyphenyl	H
	381.	4-(3-thienyl)phenyl	H
25	382.	4-(pyrazol-3-yl)phenyl	6-CH ₃
	383.	2-benzimidazolyl	H
	384.	2-benzoxazolyl	H
	385.	2-benzthiazolyl	H
	386.	6-benzimidazolyl	H
30	387.	6-benzoxazolyl	H

- 293 -


Table 3. (cont.)



5	#	R ¹	R ²
	388.		H
	389.		H
	390.		H
	391.		H
10	392.		5-Br
	393.		H
	394.		H
	395.		H

- 294 -


Table 3. (cont.)









5	#	R ¹	R ²
	396.		H
	397.		H
	398.		H
10	399.		5-Br
	400.		H
	401.		H

- 295 -


Table 3. (cont.)







5	#	R^1	R^2
			
	402.		5-Br
			
	403.		5-Br
			
	404.		5-Br
			
	405.		H
			
10	406.		H
			
	407.		H

- 296 -

Table 3. (cont.)




5	#	R ¹	R ²
			
	408.		H
			
	409.		H
			
	410.		H
			
	411.		H

10


- 297 -

Table 4.



5	#	R ¹	R ²
	412.	4-chlorophenyl	6-Cl
	413.	3,4-dichlorophenyl	5-Cl
	414.	4-fluorophenyl	H
	415.	3-chlorophenyl	H
10	416.	3-fluorophenyl	H
	417.	3-fluoro-4-methoxyphenyl	H
	418.	3-fluoro-4-methylphenyl	H
	419.	4-phenoxyphenyl	H
	420.	3-phenoxyphenyl	H
15	421.	4-biphenyl	H
	422.	4-cyclohexylphenyl	H
	423.	2-quinolyl	H
	424.	3-isoquinolyl	H
	425.	3-quinolyl	H
20	426.	1-isoquinolyl	H
	427.	5-quinolyl	H
	428.	5-isoquinolyl	H
	429.	6-quinolyl	H
	430.	6-isoquinolyl	H
25	431.	7-quinolyl	H
	432.	7-isoquinolyl	H
	433.	4-quinolyl	H
	434.	4-isoquinolyl	H
	435.	4-pyridyl	H
30	436.	4-pyrimidinyl	H


Table 4. (cont.)



5			R ²
	#	R ¹	
	437.	2-pyrimidinyl	H
	438.	6-pyrimidinyl	H
	439.	4-pyridazinyl	H
10	440.	5-pyridazinyl	H
	441.	4-indolyl	H
	442.	5-isoindolyl	H
	443.	5-naphthyridinyl	H
	444.	6-quinozalinyl	H
15	445.	6-isoquinolyl	H
	446.	4-naphthyridinyl	H
	447.	5-quinozalinyl	H
	448.	4-naphthyridinyl	H
	449.	tetrahydroquinoliny	H
20	450.	6-indazolyl	H
	451.	6-isoindolyl	H
	452.	5-indazolyl	H
	453.	5-isoindolyl	H
	454.	6-benzothienyl	H
25	455.	6-benzofuryl	H
	456.	5-benzothienyl	H
	457.	5-benzofuryl	H
	458.	2-benzimidazolyl	H
	459.	2-benzoxazolyl	H
30	460.	2-benzthiazolyl	H

- 299 -


Table 4. (cont.)



5	#	R ¹	R ²
	461.	6-benzimidazolyl	H
	462.	6-benzoxazolyl	H
	463.	6-benzthiazolyl6-benzoxazolyl	H
	464.	2-quinazolinyl6-benzoxazolyl	H
10	465.	3-(phenoxy)-6-pyridyl	H
	466.	4-(phenylcarbonyl)phenyl	H
	467.	4-(phenylamino)phenyl	H
	468.	cyclohexyloxyphenyl	H
	469.	4-(3-thienyl)phenyl	H
15	470.	4-(pyrazol-3-yl)phenyl	H
	471.	4-chlorophenyl	EtO ₂ CCH=CH-
	472.	4-chlorophenyl	5-Br

- 300 -

Example 473




5 2-(3-Oxazol-5-yl-phenylamino)-N-(3-trifluoromethyl-phenyl)-nicotinamide

ms: 447.4 [M+Na]⁺, 424.9 [M+H]⁺. Calc'd for C₂₂H₁₅F₃N₄O₂ 424.4.

10

Example 474




15 N-(4,4-Dimethyl-2-oxo-1,2,3,4-tetrahydro-quinolin-7-yl)-2-(1H-indazol-6-ylamino)-nicotinamide

[M+H]⁺ of 427.1. Calc'd for C₂₄H₂₂N₆O₂: 426.2.

- 301 -

Example 475



5

2-(2-Oxo-2,3-dihydro-1H-benzimidazol-5-ylamino)-N-(4-pentafluoroethyl-phenyl)-nicotinamide

M+H 464. Calc'd for $C_{21}H_{14}F_5N_5O_2$: 463.1.

10

Although the pharmacological properties of the compounds of Formula I-X vary with structural change, in general, activity possessed by compounds of Formula I-X may be demonstrated *in vivo*. The pharmacological properties of the compounds of this invention may be confirmed by a number of pharmacological *in vitro* assays. The exemplified pharmacological assays which follow have been carried out with the compounds according to the invention and their salts. Compounds of the present invention showed inhibition of KDR at doses less than 50 μ m.

20

- 302 -

BIOLOGICAL EVALUATION

5 HUVEC Proliferation Assay

Human Umbilical Vein Endothelial cells are purchased from Clonetics, Inc., as cryopreserved cells harvested from a pool of donors. These cells, at passage 1, are thawed and expanded in EBM-2 complete medium, until passage 2 or 3. The cells are trypsinized, washed in DMEM + 10% FBS + antibiotics, and spun at 1000 rpm for 10 min. Prior to centrifugation of the cells, a small amount is collected for a cell count. After centrifugation, the medium is discarded, and the cells are resuspended in the appropriate volume of DMEM + 10% FBS + antibiotics to achieve a concentration of 3×10^5 cells/mL. Another cell count is performed to confirm the cell concentration. The cells are diluted to 3×10^4 cells/mL in DMEM + 10% FBS + antibiotics, and 100 μ L of cells are added to a 96-well plate. The cells are incubated at 37°C for 22 h.

Prior to the completion of the incubation period, compound dilutions are prepared. Five-point, five-fold serial dilutions are prepared in DMSO, at concentrations 400-fold greater than the final concentrations desired. 2.5 μ L of each compound dilution are diluted further in a total of 1 mL DMEM + 10% FBS + antibiotics (400x dilution). Medium containing 0.25% DMSO is also prepared for the 0 μ M compound sample. At the 22-hour timepoint, the medium is removed from the cells, and 100 μ L of each compound dilution is added. The cells are incubated at 37°C for 2-3 h.

During the compound pre-incubation period, the growth factors are diluted to the appropriate concentrations. Solutions of DMEM + 10% FBS + antibiotics, containing either VEGF or bFGF at the following concentrations: 50, 10, 2, 0.4, 0.08, and 0 ng/mL are prepared. For the compound-

- 303 -

treated cells, solutions of VEGF at 550 ng/mL or bFGF at 220 ng/mL for 50 ng/mL or 20 ng/mL final concentrations, respectively, are prepared since 10 μ L of each will be added to the cells (110 μ L final volume). At the appropriate time
5 after adding the compounds, the growth factors are added. VEGF is added to one set of plates, while bFGF is added to another set of plates. For the growth factor control curves, the media on wells B4-G6 of plates 1 and 2 are replaced with media containing VEGF or bFGF at the varying
10 concentrations (50 - 0 ng/mL). The cells are incubated at 37°C for an additional 72 h.

At the completion of the 72 h incubation period, the medium is removed, and the cells are washed twice with PBS. After the second wash with PBS, the plates are tapped gently
15 to remove excess PBS, and the cells are placed at -70°C for at least 30 min. The cells are thawed and analyzed using the CyQuant fluorescent dye (Molecular Probes C-7026), following the manufacturer's recommendations. The plates are read on a Victor/Wallac 1420 workstation at 485 nm/530
20 nm (excitation/emission). Raw data are collected and analyzed using a 4-parameter fit equation in XLFit. IC₅₀ values are then determined.

The compounds of examples 3, 5-10, 13-14, 20, 22-25, 27-28, 31, 36-39, 40-43, 45-46, 48, 50-51, 54-57, 61-68, 70-
25 83, 84-90, 92-108, 110-111, 115-116, 118, 120-121, 123-126 and 406-411 inhibited VEGF-stimulated HUVEC proliferation at a level below 50 nm.

Angiogenesis Model

30

To determine the effects of the present compounds on angiogenesis *in vivo*, selective compounds are tested in the rat corneal neovascularization micropocket model or the

- 304 -

angiogenesis assay of Passaniti, Lab. Invest., 67, 519-28 (1992).

Rat Corneal Neovascularization Micropocket Model

5 **In Life Aspects:** Female Sprague Dawley rats weighing approximately 250 g were randomized into one of five treatment groups. Pretreatment with the vehicle or compound was administered orally, 24 h prior to surgery and continued
10 once a day for seven additional days. On the day of surgery, the rats were temporarily anesthetized in an Isoflurane gas chamber (delivering 2.5 liters/min oxygen + 5% Isoflurane). An othoscope was then placed inside the mouth of the animal to visualize the vocal cords. A tip-
15 blunted wire was advanced in between the vocal cords and used as a guide for the placement of an endotracheal Teflon tube (Small Parts Inc. TFE-standard Wall R-SWTT-18). A volume-controlled ventilator (Harvard Apparatus, Inc. Model 683) was connected to the endotracheal tube to deliver a
20 mixture of oxygen and 3% Isoflurane. Upon achieving deep anesthesia, the whiskers were cut short and the eye areas and eyes gently washed with Betadine soap and rinsed with sterile saline. The corneas were irrigated with one to two drops of Proparacaine HCl ophthalmic topical anesthetic
25 solution (0.5%) (Bausch and Lomb Pharmaceuticals, Tampa FL). The rat was then positioned under the dissecting microscope and the corneal surface brought into focus. A vertical incision was made on the midline of the cornea using a diamond blade knife. A pocket was created by using fine
30 scissors to separate the connective tissue layers of the stroma, tunneling towards the limbus of the eye. The distance between the apex of the pocket and the limbus was approximately 1.5 mm. After the pocket had been made, the soaked nitrocellulose disk filter (Gelman Sciences, Ann
35 Arbor MI.) was inserted under the lip of the pocket. This

- 305 -

surgical procedure was performed on both eyes. rHu-bFGF soaked disks were placed into the right eye, and the rHu-VEGF soaked disks were placed into the left eye. Vehicle soaked disks were placed in both eyes. The disk was pushed
5 into position at the desired distance from the limbal vessels. Ophthalmic antibiotic ointment was applied to the eye to prevent drying and infection. After seven days, the rats were euthanized by CO₂ asphyxiation, and the eyes enucleated. The retinal hemisphere of the eye was windowed
10 to facilitate fixation, and the eye placed into formalin overnight.

Post Mortem Aspects: After twenty-four hours in fixative, the corneal region of interest was dissected out from the eye, using fine forceps and a razorblade. The
15 retinal hemisphere was trimmed off and the lens extracted and discarded. The corneal dome was bisected and the superfluous cornea trimmed off. The iris, conjunctiva and associated limbal glands were then carefully teased away. Final cuts were made to generate a square 3x3mm containing
20 the disk, the limbus, and the entire zone of neovascularization.

Gross Image Recording: The corneal specimens were digitally photographed using a Sony CatsEye DKC5000 camera (A.G. Heinz, Irvine CA) mounted on a Nikon SMZ-U stereo
25 microscope (A.G. Heinz). The corneas were submerged in distilled water and photographed via trans-illumination at approximately 5.0 diameters magnification.

Image analysis: Numerical endpoints were generated using digital micrographs collected from the whole mount
30 corneas after trimming and were used for image analysis on the Metamorph image analysis system (Universal Imaging Corporation, West Chester PA). Three measurements were taken: Disk placement distance from the limbus, number of vessels intersecting a 2.0mm perpendicular line at the

- 306 -

midpoint of the disk placement distance, and percent blood vessel area of the diffusion determined by thresholding.

General Formulations:

- 0.1% BSA in PBS vehicle: 0.025 g of BSA was added to 25.0 ml
5 of sterile 1X phosphate buffered saline, gently shaken until fully dissolved, and filtered at 0.2 μ m. Individual 1.0 ml samples were aliquoted into 25 single use vials, and stored at -20°C until use. For the rHu-bFGF disks, a vial of this 0.1% BSA solution was allowed to thaw at room temperature.
10 Once thawed, 10 μ l of a 100 mM stock solution of DTT was added to the 1 ml BSA vial to yield a final concentration of 1 mM DTT in 0.1% BSA.

rHu-VEGF Dilutions:

- 15 Prior to the disk implant surgery, 23.8 μ l of the 0.1% BSA vehicle above was added to a 10 μ g rHu-VEGF lyophilized vial yielding a final concentration of 10 μ M.

rHu-bFGF: Stock concentration of 180 ng/ μ l:

- R&D rHu- bFGF: Added 139 μ l of the appropriate vehicle above
20 to the 25 μ g vial lyophilized vial. 13.3 μ l of the [180 ng/ μ l] stock vial and added 26.6 μ l of vehicle to yield a final concentration of 3.75 μ M concentration.

- Nitro-cellulose disk preparation:** The tip of a 20-gauge needle was cut off square and beveled with emery paper to
25 create a punch. This tip was then used to cut out \approx 0.5mm diameter disks from a nitrocellulose filter paper sheet (Gelman Sciences). Prepared disks were then placed into Eppendorf microfuge tubes containing solutions of either
0.1% BSA in PBS vehicle, 10 μ M rHu-VEGF (R&D Systems,
30 Minneapolis, MN), or 3.75 μ M rHu-bFGF (R&D Systems, Minneapolis, MN) and allowed to soak for 45-60 min before use. Each nitrocellulose filter disk absorbs approximately 0.1 μ l of solution.

- 307 -

In the rat micropocket assay, compounds of the present invention will inhibit angiogenesis at a dose of less than 50 mg/kg/day.

5

Tumor model

A431 cells (ATCC) are expanded in culture, harvested and injected subcutaneously into 5-8 week old female nude mice (CD1 nu/nu, Charles River Labs) (n=5-15). Subsequent
10 administration of compound by oral gavage (10 - 200
mpk/dose) begins anywhere from day 0 to day 29 post tumor
cell challenge and generally continues either once or twice
a day for the duration of the experiment. Progression of
tumor growth is followed by three dimensional caliper
15 measurements and recorded as a function of time. Initial
statistical analysis is done by repeated measures analysis
of variance (RMANOVA), followed by Scheffe post hoc testing
for multiple comparisons. Vehicle alone (Ora-Plus, pH 2.0)
is the negative control. Compounds of the present invention
20 are active at doses less than 150 mpk.

Rat Adjuvant Arthritis Model:

The rat adjuvant arthritis model (Pearson, Proc. Soc.
25 Exp. Biol. 91, 95-101 (1956)) is used to test the anti-
arthritic activity of compounds of the formula 1, or salts
thereof. Adjuvant Arthritis can be treated using two
different dosing schedules: either (i) starting time of
immunization with adjuvant (prophylactic dosing); or from
30 day 15 when the arthritic response is already established
(therapeutic dosing). Preferably a therapeutic dosing
schedule is used.

- 308 -

Rat Carrageenan-induced Analgesia Test

The rat carrageenan analgesia test was performed with materials, reagents and procedures essentially as described by Hargreaves, et al., (Pain, 32, 77 (1988)). Male Sprague-Dawley rats were treated as previously described for the Carrageenan Foot Pad Edema test. Three hours after the injection of the carrageenan, the rats were placed in a special plexiglass container with a transparent floor having a high intensity lamp as a radiant heat source, positionable under the floor. After an initial twenty minute period, thermal stimulation was begun on either the injected foot or on the contralateral uninjected foot. A photoelectric cell turned off the lamp and timer when light was interrupted by paw withdrawal. The time until the rat withdraws its foot was then measured. The withdrawal latency in seconds was determined for the control and drug-treated groups, and percent inhibition of the hyperalgesic foot withdrawal determined.

Formulations

Also embraced within this invention is a class of pharmaceutical compositions comprising the active compounds of Formula I-X in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The compounds and compositions of the present invention may, for example, be administered orally, mucosally, topically,

- 309 -

rectally, pulmonarily such as by inhalation spray, or parentally including intravascularly, intravenously, intraperitoneally, subcutaneously, intramuscularly intrasternally and infusion techniques, in dosage unit
5 formulations containing conventional pharmaceutically acceptable carriers, adjuvants, and vehicles.

The pharmaceutically active compounds of this invention can be processed in accordance with conventional methods of pharmacy to produce medicinal agents for
10 administration to patients, including humans and other mammals.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical
15 composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. For example, these may contain an amount of active ingredient from about 1 to 2000 mg, preferably from about 1 to 500 mg
20 or 5 to 1000 mg. A suitable daily dose for a human or other mammal may vary widely depending on the condition of the patient and other factors, but, once again, can be determined using routine methods.

The amount of compounds which are administered and the
25 dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the type of disease, the severity of the disease, the route and frequency of
30 administration, and the particular compound employed. Thus, the dosage regimen may vary widely, but can be determined routinely using standard methods. A daily dose of about 0.01 to 500 mg/kg, preferably between about 0.1 and about 50 mg/kg, and more preferably about 0.1 and about 20 mg/kg

- 310 -

body weight may be appropriate. The daily dose can be administered in one to four doses per day.

For therapeutic purposes, the active compounds of this invention are ordinarily combined with one or more
5 adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium
10 and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided
15 in a dispersion of active compound in hydroxypropylmethyl cellulose.

In the case of psoriasis and other skin conditions, it may be preferable to apply a topical preparation of compounds of this invention to the affected area two to
20 four times a day.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin (e.g., liniments, lotions, ointments, creams, or pastes) and drops suitable for
25 administration to the eye, ear, or nose. A suitable topical dose of active ingredient of a compound of the invention is 0.1 mg to 150 mg administered one to four, preferably one or two times daily. For topical administration, the active ingredient may comprise from
30 0.001% to 10% w/w, e.g., from 1% to 2% by weight of the formulation, although it may comprise as much as 10% w/w, but preferably not more than 5% w/w, and more preferably from 0.1% to 1% of the formulation.

- 311 -

When formulated in an ointment, the active ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If
5 desired, the aqueous phase of the cream base may include, for example at least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which
10 enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include DMSO and related analogs.

The compounds of this invention can also be
15 administered by a transdermal device. Preferably transdermal administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a
20 membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the
25 encapsulating agent may also function as the membrane.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or
30 an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the

- 312 -

so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers suitable
5 for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, sodium lauryl sulfate, glyceryl distearate alone or with a wax, or other materials well known in the art.

10 The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream should preferably
15 be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl
20 myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft
25 paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredients are dissolved or suspended in suitable carrier,
30 especially an aqueous solvent for the active ingredients. The active ingredients are preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% and particularly about 1.5% w/w.

- 313 -

Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules using one or more of the carriers or diluents mentioned for use in the formulations for oral administration or by using other suitable dispersing or wetting agents and suspending agents. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, tragacanth gum, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art. The active ingredient may also be administered by injection as a composition with suitable carriers including saline, dextrose, or water, or with cyclodextrin (ie. Captisol), cosolvent solubilization (ie. propylene glycol) or micellar solubilization (ie. Tween 80).

The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed, including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

For pulmonary administration, the pharmaceutical composition may be administered in the form of an aerosol or with an inhaler including dry powder aerosol.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable non-

- 314 -

irritating excipient such as cocoa butter and polyethylene glycols that are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

5 The pharmaceutical compositions may be subjected to conventional pharmaceutical operations such as sterilization and/or may contain conventional adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers, buffers etc. Tablets and pills can
10 additionally be prepared with enteric coatings. Such compositions may also comprise adjuvants, such as wetting, sweetening, flavoring, and perfuming agents.

 The foregoing is merely illustrative of the invention and is not intended to limit the invention to the disclosed
15 compounds. Variations and changes which are obvious to one skilled in the art are intended to be within the scope and nature of the invention which are defined in the appended claims.


 From the foregoing description, one skilled in the art
20 can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

 All mentioned references, patents, applications and
25 publications, are hereby incorporated by reference in their entirety, as if here written.

- 315 -


WHAT IS CLAIMED IS:

1. A compound of formula I

wherein each of A¹ and A² is independently C, CH or N;

wherein ring A is selected from

- a) 5- or 6-membered partially saturated heterocyclyl,
- 10 b) 5- or 6-membered heteroaryl,
- c) 9-, 10- or 11-membered fused partially saturated heterocyclyl,
- d) 9-, 10- or 11-membered fused heteroaryl,
- e) aryl, and
- 15 f) 4-, 5- or 6-membered cycloalkenyl;



wherein X is

wherein Z is oxygen or sulfur;

wherein R is selected from

- a) substituted or unsubstituted 4-6 membered
- 20 heterocyclyl,
- b) substituted aryl, and
- c) substituted or unsubstituted fused 9-14-membered bicyclic or tricyclic heterocyclyl;

wherein substituted R is substituted with one or more

25 substituents independently selected from halo, -OR³,

- 316 -

-SR³, -SO₂R³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -SO₂NR³R³,
 -NR³C(O)OR³, -NR³C(O)R³, cycloalkyl, optionally
 substituted 3-6 membered heterocyclyl, optionally
 substituted phenyl, nitro, alkylaminoalkoxyalkoxy,
 5 cyano, oxox, alkylaminoalkoxy, lower alkyl
 substituted with R², lower alkenyl substituted with
 R², and lower alkynyl substituted with R²;

wherein R¹ is selected from

- a) substituted or unsubstituted 6-10 membered aryl,
- 10 b) substituted or unsubstituted 4-6 membered
heterocyclyl,
- c) substituted or unsubstituted 9-14 membered bicyclic or
tricyclic heterocyclyl,
- d) cycloalkyl, and
- 15 e) cycloalkenyl,

wherein substituted R¹ is substituted with one or more
 substituents independently selected from halo, -OR³,
 -SR³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -NH(C₁-C₄
 alkylenylR¹⁴), -SO₂R³, -SO₂NR³R³, -NR³C(O)OR³, -
 20 NR³C(O)R³, -NR³C(O)NR³R³, optionally substituted
 cycloalkyl, optionally substituted 4-6 membered
 heterocyclyl, optionally substituted phenyl,
 halosulfonyl, cyano, alkylaminoalkoxy,
 alkylaminoalkoxyalkoxy, nitro, lower alkyl
 25 substituted with R², lower alkenyl substituted with
 R², and lower alkynyl substituted with R²;

wherein R² is one or more substituents independently selected
 from H, halo, -OR³, oxo, -SR³, -CO₂R³, -COR³, -CONR³R³, -
 NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, cycloalkyl,
 30 optionally substituted phenylalkylenyl, optionally
 substituted 4-6 membered heterocyclyl, optionally
 substituted heteroarylalkylenyl, optionally substituted
 phenyl, lower alkyl, cyano, lower hydroxyalkyl, lower
 carboxyalkyl, nitro, C₁₋₆-alkoxy-C₁₋₆-alkoxy, C₁₋₆-alkoxy-C₁₋

- 317 -

6-alkoxy-C₁₋₆-alkoxy, lower alkenyl, lower alkynyl, lower aminoalkyl, lower alkylaminoalkyl and lower haloalkyl; wherein R³ is independently selected from H, lower alkyl, optionally substituted phenyl, optionally substituted 3-6 membered heterocyclyl, optionally substituted C₃-C₆-cycloalkyl, optionally substituted phenylalkyl, optionally substituted 3-6 membered heterocyclalkyl, optionally substituted C₃-C₆ cycloalkylalkyl, and lower haloalkyl;

10 wherein R⁴ is selected from a direct bond, C₂₋₄-alkylenyl, C₂₋₄-alkenylenyl and C₂₋₄-alkynylenyl, where one of the CH₂ groups may be replaced with an oxygen atom or an -NH-, wherein R⁴ is optionally substituted with hydroxy;

15 wherein R⁵ is selected from H, lower alkyl, optionally substituted phenyl and optionally substituted lower aralkyl;

wherein R¹⁴ is selected from H, optionally substituted phenyl, optionally substituted 4-6 membered heterocyclyl and optionally substituted C₃-C₆ cycloalkyl;

20 and pharmaceutically acceptable derivatives thereof;

provided A is not pyridyl when X is -C(O)NH- and when R¹ is 4-[3,5-bis(trifluoromethyl)-1H-pyrazol-1-yl]phenyl when R⁵ is methyl and when R is 4-methylpiperidyl;

further provided A is not pyridyl when X is -C(O)NH-, when R⁵ is H, when R² is 6-methyl and when R is indazolyl;

25 further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl, 4-bromophenyl, 2-methylphenyl, 4-methoxyphenyl, when R⁵ is H and when R is 4-pyridyl;

further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl, when R⁵ is H and when R is 2-oxobenzopyran-4-yl;

30 further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl, 4-chlorophenyl, 3-nitrophenyl, 4-methoxyphenyl, when R⁵ is H and when R is 4-imidazolyl;

- 318 -

further provided A is not phenyl when X is -C(O)NH-, when R⁵ is H, when R^{5a} is substituted benzyl and when R is substituted triazinyl;

5 further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl or 2-chlorophenyl, when R⁵ is H and when R is 4-quinazolinyl;

further provided A is not phenyl when X is -C(O)NH-, when R¹ is phenyl, when R⁵ is H and when R is isoquinolin-1-yl;

10 further provided A is not phenyl when X is -C(O)NH-, when R¹ is 2-chlorophenyl or 4-chlorophenyl, when R⁵ is H and when R is 3-chloroisoquinolin-1-yl;

further provided A is not phenyl when X is -C(O)NH-, when R¹ is 1-ethylpiperid-3-yl or 1-ethylpiperid-4-yl, when R⁵ is H and when R is 8-trifluoromethylquinolin-4-yl;

15 further provided A is not phenyl when X is -C(O)NH-, when R¹ is 1-ethylpiperid-3-yl, when R⁵ is H and when R is 8-chloroquinolin-4-yl;

further provided A is not phenyl when X is -C(O)NH-, when R¹ is halo substituted phenyl, 1-butylpiperid-4-yl, 1-ethylpiperid-3-yl or 1-ethylpiperid-4-yl, when R⁵ is H and when R is 7-chloroquinolin-4-yl; and


20 further provided R is not unsubstituted 2-thienyl, unsubstituted 2-pyridyl or unsubstituted 3-pyridyl.

25 2. Compound of Claim 1, and pharmaceutically acceptable derivatives thereof, wherein A is selected from 5- or 6- membered partially saturated heterocyclyl.


30 3. Compound of Claim 2, wherein A is selected from dihydropyran, dihydrothienyl, dihydrofuryl, oxo-dihydrofuryl, pyrrolinyl, dihydrothiazolyl, dihydro-oxazolyl, dihydro-isothiazolyl, dihydro-isoxazolyl,

- 319 -

imidazoliny and pyrazoliny; wherein X is selected from



- wherein R is selected from substituted or unsubstituted 5-6 membered heteroaryl comprising one or more nitrogen atoms, substituted phenyl and substituted or unsubstituted 9-10 membered bicyclic or 13-14 membered tricyclic heterocyclyl; wherein substituted R is substituted with one or more substituents independently selected from halo, -OR³, -SR³, -SO₂R³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, nitro, C₁₋₄-alkylamino-C₁₋₄-alkoxy-C₁₋₄-alkoxy, cyano, C₁₋₄-alkylamino-C₁₋₄-alkoxy, C₁₋₂-alkyl substituted with R², C₂₋₃-alkenyl substituted with R², and C₂₋₃-alkynyl substituted with R²;
- wherein R¹ is selected from substituted or unsubstituted aryl selected from phenyl, naphthyl, indanyl, indenyl and tetrahydronaphthyl, substituted or unsubstituted 5-6 membered heteroaryl, C₃₋₆-cycloalkyl, and substituted or unsubstituted 9-14 membered bicyclic or tricyclic heterocyclyl; wherein substituted R¹ is substituted with one or more substituents independently selected from halo, -OR³, oxo, -SR³, -SO₂R³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -NH(C₁₋₄ alkylenylR³), -(C₁₋₄ alkylenyl)NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-alkoxy-C₁₋₆-alkoxy, halosulfonyl, optionally substituted 4-6 membered heterocyclylcarbonylalkyl, C₁₋₄-

- alkoxycarbonylamino-C₁₋₆-alkyl, , optionally substituted C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₆-alkylenyl, optionally


- 320 -

substituted 4-6 membered heterocyclyl-C₁-C₆-alkylenyl, 4-6 membered heterocyclyl-C₂-C₆-alkenylenyl, C₁₋₄-alkyl, cyano, C₁₋₄-hydroxyalkyl, nitro and C₁₋₄-haloalkyl; wherein R² is one or more substituents independently selected from H, halo, -OR³, oxo, -SR³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, C₁₋₆-alkyl, cyano, C₁₋₄-hydroxyalkyl, C₁₋₄-carboxyalkyl, nitro, C₂₋₃-alkenyl, C₂₋₃-alkynyl and C₁₋₄-haloalkyl; wherein R³ is independently selected from H, C₁₋₄-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₄-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkyl, optionally substituted C₃-C₆-cycloalkyl and C₁₋₂-haloalkyl; wherein R^{4a} is C₂₋₄-alkylenyl, where one of the CH₂ groups may be replaced with an oxygen atom or an -NH-; wherein R^{4a} is optionally substituted with hydroxy; wherein R⁵ is selected from H and C₁₋₂-alkyl; wherein R^{5a} is selected from H and C₁₋₂-alkyl; wherein R⁶ and R⁷ are independently selected from H and C₁₋₂-haloalkyl; and wherein R⁷ is selected from H, C₁₋₆-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₆-alkyl, optionally substituted cycloalkyl, optionally substituted cycloalkyl-C₁₋₆-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁-C₆-alkyl, C₁₋₄-alkoxy-C₁₋₄-alkyl and C₁₋₄-alkoxy-C₁₋₄-alkoxy-C₁₋₄-alkyl, and pharmaceutically acceptable derivatives thereof.

4. Compound of Claim 1, and pharmaceutically acceptable derivatives thereof, wherein A is selected from 5- or 6- membered heteroaryl.


- 321 -

5. Compound of Claim 4, wherein A is selected from pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, thienyl, furanyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, pyrazolyl, isoxazolyl, triazolyl and isothiazolyl; wherein X is selected from



- wherein R is selected from substituted or unsubstituted 5-6 membered heteroaryl comprising one or more nitrogen atoms, substituted phenyl and substituted or unsubstituted 9-10 membered bicyclic or 13-14 membered tricyclic heterocyclyl; wherein substituted R is substituted with one or more substituents independently selected from halo, $-\text{OR}^3$, $-\text{SR}^3$, $-\text{SO}_2\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{CONR}^3\text{R}^3$, $-\text{COR}^3$, $-\text{NR}^3\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{NR}^3\text{C}(\text{O})\text{OR}^3$, $-\text{NR}^3\text{C}(\text{O})\text{R}^3$, C_{3-6} -cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, nitro, C_{1-4} -alkylamino- C_{1-4} -alkoxy- C_{1-4} -alkoxy, cyano, C_{1-4} -alkylamino- C_{1-4} -alkoxy, C_{1-2} -alkyl substituted with R^2 , C_{2-3} -alkenyl substituted with R^2 , and C_{2-3} -alkynyl substituted with R^2 ; wherein R^1 is selected from substituted or unsubstituted aryl selected from phenyl, naphthyl, indanyl, indenyl and tetrahydronaphthyl, substituted or unsubstituted 5-6 membered heteroaryl, C_{3-6} -cycloalkyl, and substituted or unsubstituted 9-14 membered bicyclic or tricyclic heterocyclyl; wherein substituted R^1 is substituted with one or more substituents independently selected from halo, $-\text{OR}^3$, oxo, $-\text{SR}^3$, $-\text{SO}_2\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{CONR}^3\text{R}^3$, $-\text{COR}^3$, $-\text{NR}^3\text{R}^3$, $-\text{NH}(\text{C}_{1-4}\text{alkylenylR}^3)$, $-(\text{C}_{1-4}\text{alkylenyl})\text{NR}^3\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{NR}^3\text{C}(\text{O})\text{OR}^3$, $-\text{NR}^3\text{C}(\text{O})\text{R}^3$, C_{1-6} -alkylamino- C_{1-6} -alkoxy, C_{1-6} -alkylamino- C_{1-6} -alkoxy- C_{1-6} -alkoxy, halosulfonyl, optionally substituted 4-6 membered heterocyclylcarbonylalkyl, C_{1-4} -


- 322 -



- alkoxycarbonylamino-C₁₋₆-alkyl, optionally substituted C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₆-alkylenyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₆-alkylenyl, 4-6 membered heterocyclyl-C₂₋₆-alkenylenyl, C₁₋₆-alkyl, cyano, C₁₋₄-hydroxyalkyl, nitro and C₁₋₄-haloalkyl; wherein R² is one or more substituents independently selected from H, halo, -OR³, oxo, -SR³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, C₁₋₆-alkyl, cyano, C₁₋₄-hydroxyalkyl, C₁₋₄-carboxyalkyl, nitro, C₂₋₃-alkenyl, C₂₋₃-alkynyl and C₁₋₄-haloalkyl; wherein R³ is independently selected from H, C₁₋₄-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₄-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkyl, optionally substituted C₃₋₆-cycloalkyl and C₁₋₂-haloalkyl; wherein R^{4a} is C₂₋₄-alkylenyl, where one of the CH₂ groups may be replaced with an oxygen atom or an -NH-; wherein R^{4a} is optionally substituted with hydroxy; wherein R⁵ is selected from H and C₁₋₂-alkyl; wherein R^{5a} is selected from H and C₁₋₂-alkyl; wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl; and wherein R⁷ is selected from H, C₁₋₆-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₆-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₆-alkyl, C₁₋₄-alkoxy-C₁₋₄-alkyl and C₁₋₄-alkoxy-C₁₋₄-alkoxy-C₁₋₄-alkyl, and pharmaceutically acceptable derivatives thereof.


6. Compound of Claim 1 wherein A is selected from

- 323 -



wherein R^c is selected from H, methyl and optionally

5 substituted phenyl; wherein X is selected from



; wherein R is selected


from substituted or unsubstituted pyrazolyl, triazolyl, pyridyl, pyrimidinyl, and pyridazinyl, substituted phenyl, indazolyl, indolyl, isoindolyl, quinolinyl, isoquinolinyl,

10 benzotriazolyl, 2,3-dihydrobenzofuryl, 2-oxo-1,2-dihydroquinol-7-yl, naphthyridinyl and quinazolinyl; wherein substituted R is substituted with one or more substituents independently selected from halo, hydroxy, C₁₋₄-alkyl, C₁₋₂-alkoxy, optionally substituted 4-6 membered heterocyclyl-C₁-

15 ₂-alkoxy, amino, C₁₋₂-alkylamino, aminosulfonyl, -NR³C(O)OR³,

- 324 -

- NR³C(O)R³, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, nitro, C₁₋₂-alkylamino-C₁₋₂-alkoxy-C₁₋₂-alkoxy, cyano, C₁₋₂-alkylamino-C₁₋₂-alkoxy, C₁₋₂-alkylamino-C₁₋₂-alkyl, C₁₋₂-alkylamino-C₂₋₃-alkynyl, C₁₋₂-hydroxyalkyl, C₁₋₂-aminoalkyl, C₁₋₂-haloalkyl, optionally substituted 4-6 membered heterocyclyl-C₂₋₃-alkenyl, and optionally substituted 4-6 membered heterocyclyl-C₂₋₃-alkynyl; wherein R¹ is selected from substituted or unsubstituted aryl selected from phenyl, naphthyl, indanyl, indenyl and tetrahydronaphthyl, substituted or unsubstituted 5-6 membered heteroaryl, C₃₋₆-cycloalkyl, and substituted or unsubstituted 9-10 membered bicyclic or 13-14 membered tricyclic heterocyclyl; wherein substituted R¹ is substituted with one or more substituents independently selected from halo, C₁₋₆-alkyl, optionally substituted C₃₋₆-cycloalkyl, optionally substituted phenyl, optionally substituted phenyl-C₁-C₄-alkylenyl, C₁₋₂-haloalkoxy, optionally substituted phenyloxy, optionally substituted 4-6 membered heterocyclyl-C₁-C₄-alkylenyl, optionally substituted 4-6 membered heterocyclyl-C₂-C₄-alkenylenyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclxyloxy, optionally substituted 4-6 membered heterocyclylsulfonyl, optionally substituted 4-6 membered heterocyclylamino, optionally substituted 4-6 membered heterocyclylcarbonyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, oxo, cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-C₁₋₃-alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋₄-alkyl, C₁₋₄-


hydroxyalkyl,  and C₁₋₄-alkoxy; wherein R² is one

- 325 -

or more substituents independently selected from H, halo, hydroxy, C₁₋₂-alkoxy, C₁₋₂-haloalkoxy, amino, C₁₋₂-alkylamino, optionally substituted 4-6 membered heterocyclyl-C₁₋₂-alkylamino, aminosulfonyl, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, C₁₋₄-alkyl, cyano, C₁₋₂-hydroxyalkyl, C₁₋₃-carboxyalkyl, nitro, C₂₋₃-alkenyl, C₂₋₃-alkynyl and C₁₋₂-haloalkyl; wherein R³ is independently selected from H, C₁₋₄-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₄-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkyl, optionally substituted C₃-C₆ cycloalkyl and C₁₋₂-haloalkyl; wherein R^{4a} is C₂₋₃-alkylenyl where one of the CH₂ groups may be replaced with an oxygen atom or an -NH-, wherein R^{4a} is optionally substituted with hydroxy; wherein R⁵ is selected from H and C₁₋₂-alkyl; wherein R^{5a} is selected from H and C₁₋₂-alkyl; wherein R⁶ and R⁷ are independently selected from H and C₁₋₂-haloalkyl; and wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₃-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl, and pharmaceutically acceptable derivatives thereof.

25

7. Compound of Claim 6 wherein A is selected from



wherein R^c is selected from H, methyl and optionally substituted phenyl; wherein X is -C(O)-NH-; wherein R is selected from substituted or unsubstituted 4-pyridyl, 3-

30

- 326 -

pyridyl, 2-pyridyl, triazolyl, 4-pyrimidinyl, 4-pyridazinyl, optionally substituted (heterocyclyl-substituted phenyl), 5-indazolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, indolyl, isoindolyl, benzotriazolyl, 2,3-dihydrobenzofuryl, 2-oxo-
5 1,2-dihydroquinol-7-yl, quinozaliny, 4-isoquinolyl, 5-isoquinolyl, naphthyridinyl and 6-isoquinolyl; wherein substituted R is substituted with one or more substituents independently selected from chloro, fluoro, bromo, hydroxy, methoxy, ethoxy, amino, dimethylamino, diethylamino, 1-
10 methylpiperidinylmethoxy, aminosulfonyl, cyclohexyl, dimethylaminopropynyl, dimethylaminoethoxy, 3-(4-morpholinyl)propyn-1-yl, dimethylaminoethoxyethoxy, optionally substituted piperidinyl, morpholinyl, optionally substituted piperazinyl, optionally substituted phenyl,
15 methyl, ethyl, propyl, cyano, hydroxymethyl, aminomethyl, nitro and trifluoromethyl; wherein R¹ is selected from substituted or unsubstituted phenyl, indanyl, tetrahydronaphthyl, naphthyl, indazolyl, indolyl, 2,1,3-benzothiadiazolyl, cyclohexyl, isoxazolyl, pyrazolyl,
20 thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, benzothienyl, benzofuryl, benzimidazolyl, dihydro-benzimidazolyl, benzoxazolyl, benzthiazolyl,
25 isoquinolyl, quinolyl, tetrahydroquinolyl, benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, benzodioxanyl and quinazolinyl; wherein substituted R¹ is substituted with one or more substituents independently
30 selected from bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl,


- 327 -

piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl, pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl, aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy, phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy, pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and ethoxy; and wherein R² is one or more substituents independently selected from H, chloro, fluoro, bromo,

- 328 -

hydroxy, methoxy, ethoxy, trifluoromethoxy, oxo, amino, dimethylamino, aminosulfonyl, carboxymethyl, cyclopropyl, optionally substituted phenyl, methyl, ethyl, propyl, cyano, hydroxymethyl, nitro, propenyl, propynyl and trifluoromethyl
 5 and unsubstituted or substituted heteroaryl selected from thienyl, furanyl, pyridyl, imidazolyl, and pyrazolyl; and pharmaceutically acceptable derivatives thereof.

8. Compound of Claim 6 wherein A is selected from



wherein R^c is selected from H, methyl and optionally substituted phenyl; wherein X is -C(O)-NH-; wherein R is
 15 selected from substituted or unsubstituted 4-pyridyl, 3-pyridyl, 2-pyridyl, triazolyl, 4-pyrimidinyl, 4-pyridazinyl, optionally substituted (heterocyclyl-substituted phenyl), 5-indazolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, indolyl, isoindolyl, benzotriazolyl, 2,3-dihydrobenzofuryl, 2-oxo-
 20 1,2-dihydroquinol-7-yl, quinozaliny, 4-isoquinolyl, 5-isoquinolyl, naphthyridinyl and 6-isoquinolyl; wherein substituted R is substituted with one or more substituents independently selected from chloro, fluoro, bromo, hydroxy, methoxy, ethoxy, amino, dimethylamino, diethylamino, 1-
 25 methylpiperidinylmethoxy, aminosulfonyl, cyclohexyl, dimethylaminopropynyl, dimethylaminoethoxy, 3-(4-morpholinyl)propyn-1-yl, dimethylaminoethoxyethoxy, optionally substituted piperidinyl, morpholinyl, optionally


- 329 -

- substituted piperazinyl, optionally substituted phenyl, methyl, ethyl, propyl, cyano, hydroxymethyl, aminomethyl, nitro and trifluoromethyl; wherein R¹ is selected from substituted or unsubstituted phenyl, indanyl,
- 5 tetrahydronaphthyl, naphthyl, indazolyl, indolyl, 2,1,3-benzothiadiazolyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-
- 10 isoquinolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, benzothienyl, benzofuryl, benzimidazolyl, dihydro-benzimidazolyl, benzoxazolyl, benzthiazolyl, isoquinolyl, quinolyl, tetrahydroquinolyl,
- 15 benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, benzodioxanyl and quinazolinyl; wherein substituted R¹ is substituted with one or more substituents independently selected from bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl,
- 20 phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl, piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-
- 25 Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,
- 30 pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,

- 330 -

aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
 5 imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-
 10 di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
 15 phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy, pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and
 20 ethoxy; and wherein R^2 is one or more substituents independently selected from H, chloro, fluoro, bromo, hydroxy, methoxy, ethoxy, trifluoromethoxy, oxo, amino, dimethylamino, aminosulfonyl, carboxymethyl, cyclopropyl, optionally substituted phenyl, methyl, ethyl, propyl, cyano,
 25 hydroxymethyl, nitro, propenyl, propynyl and trifluoromethyl and unsubstituted or substituted heteroaryl selected from thienyl, furanyl, pyridyl, imidazolyl, and pyrazolyl; and pharmaceutically acceptable derivatives thereof.

30 9. Compound of Claim 6 wherein A is selected from



- 331 -

wherein R^c is selected from H, methyl and optionally substituted phenyl; wherein X is -C(O)-NH-; wherein R is selected from substituted or unsubstituted 4-pyridyl, 3-
5 pyridyl, 2-pyridyl, triazolyl, 4-pyrimidinyl, 4-pyridazinyl, optionally substituted (heterocyclyl-substituted phenyl), 5-indazolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, indolyl, isoindolyl, benzotriazolyl, 2,3-dihydrobenzofuryl, 2-oxo-1,2-dihydroquinol-7-yl, quinoxalyl, 4-isoquinolyl, 5-
10 isoquinolyl, naphthyridinyl and 6-isoquinolyl; wherein substituted R is substituted with one or more substituents independently selected from chloro, fluoro, bromo, hydroxy, methoxy, ethoxy, amino, dimethylamino, diethylamino, 1-methylpiperidinylmethoxy, aminosulfonyl, cyclohexyl,
15 dimethylaminopropynyl, dimethylaminoethoxy, 3-(4-morpholinyl)propyn-1-yl, dimethylaminoethoxyethoxy, optionally substituted piperidinyl, morpholinyl, optionally substituted piperazinyl, optionally substituted phenyl, methyl, ethyl, propyl, cyano, hydroxymethyl, aminomethyl,
20 nitro and trifluoromethyl; wherein R¹ is selected from substituted or unsubstituted phenyl, indanyl, tetrahydronaphthyl, naphthyl, indazolyl, indolyl, 2,1,3-benzothiadiazolyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl,
25 pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydroisoquinolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, benzothienyl, benzofuryl, benzimidazolyl, dihydro-benzimidazolyl, benzoxazolyl, benzthiazolyl, isoquinolyl, quinolyl, tetrahydroquinolyl,
30 benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-azafluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, benzodioxanyl and quinazolinyl; wherein substituted R¹ is substituted with one or more substituents independently selected from bromo, chloro, fluoro, iodo, nitro, amino,


- 332 -

- cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo,
aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl,
phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-
4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl,
5 piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-
methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-
(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-
Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-
piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-
10 4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-
ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl,
pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,
pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-
pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl,
15 pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl,
methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-
methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,
aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-
ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-
20 methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-
methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,
hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-
butyl, sec-butyl, trifluoromethyl, pentafluoroethyl,
25 nonafluorobutyl, dimethylaminopropyl, 1,1-
di(trifluoromethyl)-1-hydroxymethyl, 1,1-
di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-
di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-
30 aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy,
pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-
ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-

- 333 -


2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-ylmethoxy, isopropoxy, methoxy and ethoxy; and wherein R^2 is one or more substituents independently selected from H, chloro, fluoro, bromo, hydroxy, methoxy, ethoxy, trifluoromethoxy, oxo, amino, dimethylamino, aminosulfonyl, carboxymethyl, cyclopropyl, optionally substituted phenyl, methyl, ethyl, propyl, cyano, hydroxymethyl, nitro, propenyl, propynyl and trifluoromethyl and unsubstituted or substituted heteroaryl selected from thienyl, furanyl, pyridyl, imidazolyl, and pyrazolyl; and pharmaceutically acceptable derivatives thereof.

10. Compound of Claim 1, wherein A is selected from



15

wherein X is selected from




wherein R is selected from substituted or unsubstituted pyrazolyl, triazolyl, pyridyl, pyrimidinyl, and pyridazinyl, substituted phenyl, indazolyl, indolyl, isoindolyl, quinolinyl, isoquinolinyl, benzotriazolyl, 2,3-dihydrobenzofuryl, 2-oxo-1,2-dihydroquinol-7-yl, naphthyridinyl and quinazolinyl; wherein substituted R is substituted with one or more substituents independently selected from halo, hydroxy, C_{1-4} -alkyl, C_{1-2} -alkoxy,

25


- 334 -


- optionally substituted 4-6 membered heterocyclyl-C₁₋₂-alkoxy, amino, C₁₋₂-alkylamino, aminosulfonyl, -NR³C(O)OR³, -NR³C(O)R³, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, nitro,
- 5 C₁₋₂-alkylamino-C₁₋₂-alkoxy-C₁₋₂-alkoxy, cyano, C₁₋₂-alkylamino-C₁₋₂-alkoxy, C₁₋₂-alkylamino-C₁₋₂-alkyl, C₁₋₂-alkylamino-C₂₋₃-alkynyl, C₁₋₂-hydroxyalkyl, C₁₋₂-aminoalkyl, C₁₋₂-haloalkyl, optionally substituted 4-6 membered heterocyclyl-C₂₋₃-alkenyl, and optionally substituted 4-6 membered
- 10 heterocyclyl-C₂₋₃-alkynyl; wherein R¹ is selected from substituted or unsubstituted aryl selected from phenyl, naphthyl, indanyl, indenyl and tetrahydronaphthyl, substituted or unsubstituted 5-6 membered heteroaryl, C₃₋₆-cycloalkyl, and substituted or unsubstituted 9-10 membered
- 15 bicyclic or 13-14 membered tricyclic heterocyclyl; wherein substituted R¹ is substituted with one or more substituents independently selected from halo, C₁₋₆-alkyl, optionally substituted C₃₋₆-cycloalkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₄-alkylenyl, C₁₋₂-
- 20 haloalkoxy, optionally substituted phenyloxy, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkylenyl, optionally substituted 4-6 membered heterocyclyl-C₂₋₄-alkenylenyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered
- 25 heterocyclyloxy, optionally substituted 4-6 membered heterocyclylsulfonyl, optionally substituted 4-6 membered heterocyclylamino, optionally substituted 4-6 membered heterocyclylcarbonyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-haloalkyl, C₁₋₄-
- 30 aminoalkyl, nitro, amino, hydroxy, cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-

- 335 -



- alkoxycarbonylamino-C₁₋₄-alkyl, C₁₋₄-hydroxyalkyl, and C₁₋₄-alkoxy; wherein R² is one or more substituents independently selected from H, halo, hydroxy, C₁₋₂-alkoxy, C₁₋₂-haloalkoxy, amino, C₁₋₂-alkylamino, optionally substituted 4-6 membered heterocyclyl-C₁₋₂-alkylamino, aminosulfonyl, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, C₁₋₄-alkyl, cyano, C₁₋₂-hydroxyalkyl, C₁₋₃-carboxyalkyl, nitro, C₂₋₃-alkenyl, C₂₋₃-alkynyl and C₁₋₂-haloalkyl; wherein R³ is independently selected from H, C₁₋₄-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₄-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkyl, optionally substituted C₃₋₆ cycloalkyl and C₁₋₂-haloalkyl; wherein R^{4a} is C₂₋₃-alkylenyl where one of the CH₂ groups may be replaced with an oxygen atom or an -NH-, wherein R^{4a} is optionally substituted with hydroxy; wherein R⁵ is selected from H and C₁₋₂-alkyl; wherein R^{5a} is selected from H and C₁₋₂-alkyl; wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl; and wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₃-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl, and pharmaceutically acceptable derivatives thereof.

11. Compound of Claim 10, wherein A is , or



; wherein X is -C(O)-NH-; wherein R is selected

- 336 -

from substituted or unsubstituted 4-pyridyl, 3-pyridyl, 2-pyridyl, triazolyl, 4-pyrimidinyl, 4-pyridazinyl, optionally substituted (heterocyclyl-substituted phenyl), 5-indazolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, indolyl, isoindolyl, 5 benzotriazolyl, 2,3-dihydrobenzofuryl, 2-oxo-1,2-dihydroquinol-7-yl, quinoxalyl, 4-isoquinolyl, 5-isoquinolyl, naphthyridinyl and 6-isoquinolyl; wherein substituted R is substituted with one or more substituents independently selected from chloro, fluoro, bromo, hydroxy, 10 methoxy, ethoxy, amino, dimethylamino, diethylamino, 1-methylpiperidinylmethoxy, aminosulfonyl, cyclohexyl, dimethylaminopropynyl, dimethylaminoethoxy, 3-(4-morpholinyl)propyn-1-yl, dimethylaminoethoxyethoxy, optionally substituted piperidinyl, morpholinyl, optionally 15 substituted piperazinyl, optionally substituted phenyl, methyl, ethyl, propyl, cyano, hydroxymethyl, aminomethyl, nitro and trifluoromethyl; wherein R¹ is selected from substituted or unsubstituted phenyl, indanyl, tetrahydronaphthyl, naphthyl, indazolyl, indolyl, 2,1,3-benzothiadiaazolyl, cyclohexyl, isoxazolyl, pyrazolyl, 20 thiazolyl, thiadiaazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 2-oxo-1,2-dihydroquinol-7-yl, 1,2,3,4-tetrahydro-isoquinolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, benzothienyl, benzofuryl, benzimidazolyl, 25 dihydro-benzimidazolyl, benzoxazolyl, benzthiazolyl, isoquinolyl, quinolyl, tetrahydroquinolyl, benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, benzodioxanyl and quinazolinyl; wherein substituted R¹ is 30 substituted with one or more substituents independently selected from bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-


- 337 -

- 4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl, piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-
- 5 Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,
- 10 pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,
- 15 aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,
- 20 hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
- 25 hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy, phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy,
- 30 pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and ethoxy; and wherein R² is one or more substituents

- 338 -


independently selected from H, chloro, fluoro, bromo, hydroxy, methoxy, ethoxy, trifluoromethoxy, oxo, amino, dimethylamino, aminosulfonyl, carboxymethyl, cyclopropyl, optionally substituted phenyl, methyl, ethyl, propyl, cyano, hydroxymethyl, nitro, propenyl, propynyl, trifluoromethyl and unsubstituted or substituted heteroaryl selected from thienyl, furanyl, pyridyl, imidazolyl, and pyrazolyl; and pharmaceutically acceptable derivatives thereof.

- 10 12. Compound of Claim 1 wherein A is 9- or 10-membered fused partially saturated heterocyclyl or 9- or 10-membered fused heteroaryl; wherein X is selected from



- wherein R is selected from substituted or
- 15 unsubstituted 5-6 membered heteroaryl comprising one or more nitrogen atoms, substituted phenyl and substituted or unsubstituted 9-10 membered bicyclic or 13-14 membered tricyclic heterocyclyl; wherein substituted R is substituted with one or more substituents independently selected from
- 20 halo, $-\text{OR}^3$, $-\text{SR}^3$, $-\text{SO}_2\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{CONR}^3\text{R}^3$, $-\text{COR}^3$, $-\text{NR}^3\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{NR}^3\text{C}(\text{O})\text{OR}^3$, $-\text{NR}^3\text{C}(\text{O})\text{R}^3$, C_{3-6} -cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, nitro, C_{1-4} -alkylamino- C_{1-4} -alkoxy- C_{1-4} -alkoxy, cyano, C_{1-4} -alkylamino- C_{1-4} -alkoxy, C_{1-2} -alkyl
- 25 substituted with R^2 , C_{2-3} -alkenyl substituted with R^2 , and C_{2-3} -alkynyl substituted with R^2 ; wherein R^1 is selected from substituted or unsubstituted aryl selected from phenyl, naphthyl, indanyl, indenyl and tetrahydronaphthyl, substituted or unsubstituted 5-6 membered heteroaryl, C_{3-6} -
- 30 cycloalkyl, and substituted or unsubstituted 9-14 membered bicyclic or tricyclic heterocyclyl; wherein substituted R^1

- 339 -

- is substituted with one or more substituents independently selected from halo, $-OR^3$, oxo, $-SR^3$, $-SO_2R^3$, $-CO_2R^3$, $-CONR^3R^3$, $-COR^3$, $-NR^3R^3$, $-NH(C_1-C_4 \text{ alkylene}R^3)$, $-(C_1-C_4 \text{ alkylene})NR^3R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$, C_1-C_6 -alkylamino- C_1-C_6 -alkoxy, C_1-C_6 -alkylamino- C_1-C_6 -alkoxy- C_1-C_6 -alkoxy, halosulfonyl, optionally substituted 4-6 membered heterocyclylcarbonylalkyl, C_1-4 -alkoxycarbonylamino- C_1-6 -alkyl, , optionally substituted C_{3-6} -cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, optionally substituted phenyl- C_{1-6} -alkylene, optionally substituted 4-6 membered heterocyclyl- C_1-C_6 -alkylene, 4-6 membered heterocyclyl- C_2-C_6 -alkylene, C_1-6 -alkyl, cyano, C_1-4 -hydroxyalkyl, nitro and C_1-4 -haloalkyl; wherein R^2 is one or more substituents independently selected from H, halo, $-OR^3$, oxo, $-SR^3$, $-CO_2R^3$, $-CONR^3R^3$, $-COR^3$, $-NR^3R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$, C_{3-6} -cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, C_{1-6} -alkyl, cyano, C_1-4 -hydroxyalkyl, C_1-4 -carboxyalkyl, nitro, C_{2-3} -alkenyl, C_{2-3} -alkynyl and C_1-4 -haloalkyl; wherein R^3 is independently selected from H, C_{1-4} -alkyl, optionally substituted phenyl, optionally substituted phenyl- C_{1-4} -alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl- C_{1-4} -alkyl, optionally substituted C_3-C_6 cycloalkyl and C_{1-2} -haloalkyl; wherein R^{4a} is C_{2-4} -alkylene, where one of the CH_2 groups may be replaced with an oxygen atom or an $-NH-$; wherein R^{4a} is optionally substituted with hydroxy; wherein R^5 is selected from H and C_{1-2} -alkyl; wherein R^{5a} is selected from H and C_{1-2} -alkyl; wherein R^6 and R^7 are independently selected from H and C_{1-2} -haloalkyl; and wherein R^7 is selected from H, C_{1-6} -alkyl, optionally substituted phenyl, optionally substituted phenyl- C_{1-6} -alkyl, optionally substituted 4-6 membered


- 340 -

heterocyclyl, optionally substituted 4-6 membered heterocyclyl- C_1 - C_6 -alkyl, C_{1-4} -alkoxy- C_{1-4} -alkyl and C_{1-4} -alkoxy- C_{1-4} -alkoxy- C_{1-4} -alkyl, and pharmaceutically acceptable derivatives thereof.

5

13. Compound of Claim 12 wherein A is selected from benzothienyl, benzofuryl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indazolyl, indolyl, isoindolyl, quinolyl, isoquinolyl, naphthpyridinyl, tetrahydroquinolyl,
 10 quinoxalinyl and quinazolinyl;
 and pharmaceutically acceptable salts thereof.


14. Compound of Claim 1, wherein A is 5- or 6-membered cycloalkenyl; wherein X is selected from



wherein R is selected from substituted or unsubstituted 5-6 membered heteroaryl comprising one or more nitrogen atoms, substituted phenyl and substituted or unsubstituted 9-10 membered bicyclic or 13-14 membered
 20 tricyclic heterocyclyl; wherein substituted R is substituted with one or more substituents independently selected from halo, $-OR^3$, $-SR^3$, $-SO_2R^3$, $-CO_2R^3$, $-CONR^3R^3$, $-COR^3$, $-NR^3R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$, C_{3-6} -cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally
 25 substituted phenyl, nitro, C_{1-4} -alkylamino- C_{1-4} -alkoxy- C_{1-4} -alkoxy, cyano, C_{1-4} -alkylamino- C_{1-4} -alkoxy, C_{1-2} -alkyl substituted with R^2 , C_{2-3} -alkenyl substituted with R^2 , and C_{2-3} -alkynyl substituted with R^2 ; wherein R^1 is selected from substituted or unsubstituted aryl selected from phenyl,
 30 naphthyl, indanyl, indenyl and tetrahydronaphthyl, substituted or unsubstituted 5-6 membered heteroaryl, C_{3-6} -

- 341 -


cycloalkyl, and substituted or unsubstituted 9-14 membered bicyclic or tricyclic heterocyclyl; wherein substituted R^1 is substituted with one or more substituents independently selected from halo, $-OR^3$, oxo, $-SR^3$, $-SO_2R^3$, $-CO_2R^3$, $-CONR^3R^3$,
 5 $-COR^3$, $-NR^3R^3$, $-NH(C_1-C_4 \text{ alkylenyl}R^3)$, $-(C_1-C_4 \text{ alkylenyl})NR^3R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$, C_1-C_6 -alkylamino- C_1-C_6 -alkoxy, C_1-C_6 -alkylamino- C_1-C_6 -alkoxy- C_1-C_6 -alkoxy, halosulfonyl, optionally substituted 4-6 membered heterocyclylcarbonylalkyl, C_{1-4} -alkoxycarbonylamino- C_{1-6} -

- 10 alkyl, , optionally substituted C_{3-6} -cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, optionally substituted phenyl- C_{1-6} -alkylenyl, optionally substituted 4-6 membered heterocyclyl- C_{1-6} -alkylenyl, 4-6 membered heterocyclyl- C_{2-6} -alkenylenyl,
 15 C_{1-4} -alkyl, cyano, C_{1-4} -hydroxyalkyl, nitro and C_{1-4} -haloalkyl; wherein R^2 is one or more substituents independently selected from H, halo, $-OR^3$, oxo, $-SR^3$, $-CO_2R^3$, $-CONR^3R^3$, $-COR^3$, $-NR^3R^3$, $-SO_2NR^3R^3$, $-NR^3C(O)OR^3$, $-NR^3C(O)R^3$, C_{3-6} -cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally
 20 substituted phenyl, C_{1-6} -alkyl, cyano, C_{1-4} -hydroxyalkyl, C_{1-4} -carboxyalkyl, nitro, C_{2-3} -alkenyl, C_{2-3} -alkynyl and C_{1-4} -haloalkyl; wherein R^3 is independently selected from H, C_{1-4} -alkyl, optionally substituted phenyl, optionally substituted phenyl- C_{1-4} -alkyl, optionally substituted 4-6 membered
 25 heterocyclyl, optionally substituted 4-6 membered heterocyclyl- C_{1-4} -alkyl, optionally substituted C_3-C_6 cycloalkyl and C_{1-2} -haloalkyl; wherein R^{4a} is C_{2-4} -alkylenyl, where one of the CH_2 groups may be replaced with an oxygen atom or an $-NH-$; wherein R^{4a} is optionally substituted with
 30 hydroxy; wherein R^5 is selected from H and C_{1-2} -alkyl; wherein R^{5a} is selected from H and C_{1-2} -alkyl; wherein R^e and R^f are independently selected from H and C_{1-2} -haloalkyl; and wherein R^7 is selected from H, C_{1-6} -alkyl, optionally substituted

- 342 -

phenyl, optionally substituted phenyl-C₁₋₆-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₆-alkyl, C₁₋₄-alkoxy-C₁₋₄-alkyl and C₁₋₄-alkoxy-C₁₋₄-alkoxy-C₁₋₄-alkyl, and
 5 pharmaceutically acceptable derivatives thereof.


15. Compound of Claim 1, wherein A is phenyl; wherein X is selected from



10 wherein R is selected from substituted or unsubstituted 5-6 membered heteroaryl comprising one or more nitrogen atoms, substituted phenyl and substituted or unsubstituted 9-10 membered bicyclic or 13-14 membered tricyclic heterocyclyl; wherein substituted R is substituted
 15 with one or more substituents independently selected from halo, -OR³, -SR³, -SO₂R³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₃₋₆-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, nitro, C₁₋₄-alkylamino-C₁₋₄-alkoxy-C₁₋₄-
 20 alkoxy, cyano, C₁₋₄-alkylamino-C₁₋₄-alkoxy, C₁₋₂-alkyl substituted with R², C₂₋₃-alkenyl substituted with R², and C₂₋₃-alkynyl substituted with R²; wherein R¹ is selected from substituted or unsubstituted aryl selected from phenyl, naphthyl, indanyl, indenyl and tetrahydronaphthyl,
 25 substituted or unsubstituted 5-6 membered heteroaryl, C₃₋₆-cycloalkyl, and substituted or unsubstituted 9-14 membered bicyclic or tricyclic heterocyclyl; wherein substituted R¹ is substituted with one or more substituents independently selected from halo, -OR³, oxo, -SR³, -SO₂R³, -CO₂R³, -CONR³R³,
 30 -COR³, -NR³R³, -NH(C₁₋₄ alkylene)R³, -(C₁₋₄ alkylene)NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₁₋₆-alkylamino-C₁₋₆-

- 343 -

alkoxy, C₁-C₆-alkylamino-C₁-C₆-alkoxy-C₁-C₆-alkoxy,
 halosulfonyl, optionally substituted 4-6 membered
 heterocyclylcarbonylalkyl, C₁-4-alkoxycarbonylamino-C₁-6-

alkyl, , optionally substituted C₃-6-cycloalkyl,

- 5 optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, optionally substituted phenyl-C₁-6-alkylenyl, optionally substituted 4-6 membered heterocyclyl-C₁-C₆-alkylenyl, 4-6 membered heterocyclyl-C₂-C₆-alkenylenyl, C₁-6-alkyl, cyano, C₁-4-hydroxyalkyl, nitro and C₁-4-haloalkyl;
- 10 wherein R² is one or more substituents independently selected from H, halo, -OR³, oxo, -SR³, -CO₂R³, -CONR³R³, -COR³, -NR³R³, -SO₂NR³R³, -NR³C(O)OR³, -NR³C(O)R³, C₃-6-cycloalkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted phenyl, C₁-6-alkyl, cyano, C₁-4-hydroxyalkyl, C₁-4-
- 15 carboxyalkyl, nitro, C₂-3-alkenyl, C₂-3-alkynyl and C₁-4-haloalkyl; wherein R³ is independently selected from H, C₁-4-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁-4-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered
- 20 heterocyclyl-C₁-4-alkyl, optionally substituted C₃-C₆-cycloalkyl and C₁-2-haloalkyl; wherein R^{4a} is C₂-4-alkylenyl, where one of the CH₂ groups may be replaced with an oxygen atom or an -NH-; wherein R^{4a} is optionally substituted with hydroxy; wherein R⁵ is selected from H and C₁-2-alkyl; wherein
- 25 R^{5a} is selected from H and C₁-2-alkyl; wherein R^e and R^f are independently selected from H and C₁-2-haloalkyl; and wherein R⁷ is selected from H, C₁-6-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁-6-alkyl, optionally substituted 4-6 membered heterocyclyl, optionally
- 30 substituted 4-6 membered heterocyclyl-C₁-C₆-alkyl, C₁-4-alkoxy-C₁-4-alkyl and C₁-4-alkoxy-C₁-4-alkoxy-C₁-4-alkyl, and pharmaceutically acceptable derivatives thereof.

- 344 -

16. Compound of Claim 1 and pharmaceutically acceptable salts thereof selected from
- N-(4-Chlorophenyl) [2-(6-quinolylamino) (3-pyridyl)] carboxamide;
- 5 N-(4-Chlorophenyl) [2-(5-isoquinolylamino) (3-pyridyl)] carboxamide;
- N-(4-Chlorophenyl) [2-(1H-indazol-5-ylamino) (3-pyridyl)] carboxamide;
- N-(4-Chlorophenyl) [2-(1H-indazol-6-ylamino) (3-pyridyl)] carboxamide;
- 10 2-(1H-Indazol-6-ylamino)-N-(4-isopropyl-phenyl) nicotinamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)] -N-[3-(methylethyl) phenyl] carboxamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)] -N-[4-(methylpropyl) phenyl] carboxamide;
- 15 N-[4-(tert-Butyl) phenyl] [2-(1H-indazol-6-ylamino) (3-pyridyl)] carboxamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)] -N-[3-(trifluoromethyl) phenyl] carboxamide;
- 20 N-[3-(tert-Butyl) phenyl] [2-(1H-indazol-6-ylamino) (3-pyridyl)] carboxamide;
- [2-(Benzotriazol-6-ylamino) (3-pyridyl)] -N-[4-(tert-butyl) phenyl] carboxamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)] -N-(3-phenylpyrazol-5-yl) carboxamide;
- 25 N-(4-Chloro-3-sulfamoylphenyl) [2-(1H-indazol-6-ylamino) (3-pyridyl)] carboxamide;
- [2-(1H-Indazol-6-ylamino) (3-pyridyl)] -N-(4-methyl-2-oxo-1,2-dihydroquinol-7-yl) carboxamide;
- 30 N-[4-(Methylethyl) phenyl] {2-[(4-methyl-2-oxo(7-hydroquinolyl)) amino] (3-pyridyl)} carboxamide;
- N-[5-(tert-Butyl) isoxazol-3-yl] [2-(1H-indazol-6-ylamino) (3-pyridyl)] carboxamide;

- 345 -

- N-[5-(tert-Butyl)-1-methylpyrazol-3-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- N-[4-(tert-Butyl)(1,3-thiazol-2-yl)][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 5 N-[5-(tert-Butyl)(1,3,4-thiadiazol-2-yl)][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(1,1,2,2,3,3,4,4,4-nonafluorobutyl)phenyl]carboxamide;
- {2-[(1-Methyl(1H-indazol-6-yl))amino](3-pyridyl)}-N-[4-(methylethyl)phenyl]carboxamide;
- 10 N-[4-(tert-Butyl)phenyl]{2-[(7-bromo(1H-indazol-6-yl))amino](3-pyridyl)}carboxamide;
- 2-(1H-Indazol-6-ylamino)-N-[4-tert-butyl-3-(1,2,3,6-tetrahydropyridin-4-yl)phenyl]nicotinamide;
- 15 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]carboxamide;
- N-[5-(tert-Butyl)-2-methoxyphenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 20 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-{6-[4-(trifluoromethyl)piperidyl](3-pyridyl)}carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(1-oxo(7-2,3,4-trihydroisoquinolyl))carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylethoxy)phenyl]carboxamide;
- 25 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-{4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl}carboxamide;
- N-(4-{(1S)-1-[(Methylethyl)amino]ethyl}phenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 30 N-[4-(tert-Butyl)-3-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(4-methylpiperazinyl)phenyl]carboxamide;

- 346 -

- N-[4-(tert-Butyl)-2-(4-methylpiperazinyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- N-[2-[2-(Dimethylamino)ethoxy]-5-(tert-butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 5 N-[3-[2-(Dimethylamino)ethoxy]phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- N-(3-Hydroxy-4-methoxyphenyl)[2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- N-[3-[2-(Dimethylamino)ethoxy]-4-methoxyphenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 10 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-methoxy-3-(1-methyl(4-piperidyl)oxy)phenyl]carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolin-2-yl)carboxamide;
- 15 [2-((3-[2-(Dimethylamino)ethoxy](1H-indazol-6-yl))amino)(3-pyridyl)]-N-[4-(tert-butyl)phenyl]carboxamide;
- N-[3,3-Dimethyl-1-(4-piperidylmethyl)indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- N-[3,3-Dimethyl-1-(1-methyl-piperidin-4-ylmethyl)-2,3-dihydro-1H-indol-6-yl]-2-(1H-indazol-6-ylamino)-
- 20 nicotinamide;
- 2-(1H-Indazol-6-ylamino)-N-(4-phenoxy-phenyl)-nicotinamide;
- [2-(1H-Indazol-5-ylamino)(3-pyridyl)]-N-(4-phenoxyphenyl)carboxamide;
- 25 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-(4-phenylphenyl)carboxamide;
- [2-(1H-indazol-6-ylamino)(3-pyridyl)]-N-[4-(methylsulfonyl)phenyl]carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(1-methyl(4-piperidyl))indolin-6-yl]carboxamide;
- 30 N-[3,3-Dimethyl-1-(1-methyl(4-piperidyl))indolin-6-yl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-(1-methyl(4-piperidyl))indol-5-yl]carboxamide;


- 347 -

- [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(trifluoromethyl)phenyl]carboxamide;
N-[3-[3-(Dimethylamino)propyl]-5-(trifluoromethyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 5 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[5-(1-methyl(4-1,2,5,6-tetrahydropyridyl))-3-(trifluoromethyl)phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-(1-methyl(4-piperidyl))phenyl]carboxamide;
- 10 N-[4-(tert-Butyl)-3-(3-piperidylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
N-[3-((1E)-4-Pyrrolidinylbut-1-enyl)-4-(tert-butyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
- 15 N-[4-(tert-Butyl)-3-(3-pyrrolidinylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
N-[4-(tert-Butyl)-3-(3-morpholin-4-ylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-[3-(4-
- 20 methylpiperazinyl)-3-oxopropyl]phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-[3-(4-methylpiperazinyl)-3-oxopropyl]phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[3-[3-(4-methylpiperazinyl)propyl]phenyl]carboxamide;
- 25 [2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[4-[3-(4-methylpiperazinyl)propyl]phenyl]carboxamide;
[2-(1H-Indazol-6-ylamino)(3-pyridyl)]-N-[1-(2-morpholin-4-ylethyl)indol-6-yl]carboxamide;
- 30 N-[4-(1,1-Dimethyl-3-morpholin-4-ylpropyl)phenyl][2-(1H-indazol-6-ylamino)(3-pyridyl)]carboxamide;
2-(1H-Indazol-6-ylamino)-N-(4-(2,2,2-trifluoro-1-[2-(2-methoxy-ethoxy)-ethoxy]-1-trifluoromethyl-ethyl)-phenyl)-nicotinamide;

- 348 -

[2-(1H-Indazol-6-ylamino) (3-pyridyl)]-N-{4-[2,2,2-trifluoro-
 1-(2-piperidylethoxy)-1-
 (trifluoromethyl)ethyl]phenyl}carboxamide;
 N-[4-(*tert*-Butyl)phenyl][6-fluoro-2-(1H-indazol-6-
 5 ylamino) (3-pyridyl)]carboxamide;
 [6-Fluoro-2-(1H-indazol-6-ylamino) (3-pyridyl)]-N-[4-
 (methylethyl)phenyl]carboxamide;
 [6-Fluoro-2-(1H-indazol-6-ylamino) (3-pyridyl)]-N-[3-
 (trifluoromethyl)phenyl]carboxamide; and
 10 {2-[(1-(2-Pyridyl)pyrrolidin-3-yl)amino] (3-pyridyl)}-N-[3-
 (trifluoromethyl)phenyl]carboxamide.

17. A compound of Claim 1 having Formula II



II


15 wherein R is selected from unsubstituted or substituted 9-
 or 10-membered fused nitrogen-containing heteroaryl,
 wherein R is substituted with one or more substituents
 20 selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
 haloalkyl, C₁₋₆-alkoxy, optionally substituted
 heterocyclalkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
 C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
 alkoxy-C₁₋₆-alkoxy, and optionally substituted
 25 heterocyclalkyl-C₂₋₄-alkynyl;
 wherein R¹ is selected from unsubstituted or substituted
 aryl,
 cycloalkyl,
 5-6 membered heteroaryl and

- 349 -

9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclyl,

wherein substituted R¹ is substituted with one or more

substituents selected from halo, C₁₋₆-alkyl, optionally
5 substituted C₃₋₆-cycloalkyl, optionally substituted
phenyl, optionally substituted phenyl-C₁-C₄-alkylenyl,
C₁₋₂-haloalkoxy, optionally substituted phenyloxy,
optionally substituted 4-6 membered heterocyclyl-C₁-C₄-
alkyl, optionally substituted 4-6 membered
10 heterocyclyl-C₂-C₄-alkenyl, optionally substituted 4-6
membered heterocyclyl, optionally substituted 4-6
membered heterocycliloxy, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclylsulfonyl,
15 optionally substituted 4-6 membered heterocyclylamino,
optionally substituted 4-6 membered
heterocyclylcarbonyl, optionally substituted 4-6
membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
20 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋


4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy;

25 wherein R² is one or more substituents independently
selected from

H,
halo,
hydroxy,
30 amino,
C₁₋₆-alkyl,
C₁₋₆-haloalkyl,
C₁₋₆-alkoxy,

- 350 -

- C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 cyano,
 5 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,
 10 C₁₋₆-carboxyalkyl,
 4-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 heterocyclyl;
 15 wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and



; and

- wherein R^e and R^f are independently selected from H and C₁₋₂-
 haloalkyl; and
 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally
 20 substituted phenyl, optionally substituted phenyl-C₁₋₃-
 alkyl, 4-6 membered heterocyclyl, optionally
 substituted 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-
 alkoxy-C₁₋₂-alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;
 and pharmaceutically acceptable derivatives thereof.


25

18. Compound of Claim 17 wherein R is selected from
 indazolyl, where R is unsubstituted or substituted with one
 or more substituents selected from chloro, fluoro, amino,
 hydroxy, methyl, ethyl, propyl, trifluoromethyl,
 30 dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
 dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein R¹ is
 selected from phenyl, tetrahydronaphthyl, indanyl, indenyl,
 naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl,

- 351 -


thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, quinoxalyl, benzo[d]isothiazolyl, 5 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolinyl, indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl, benzothienyl, benzofuryl, benzimidazolyl, dihydro-benzimidazolyl, benzoxazolyl and benzthiazolyl, where R¹ is 10 unsubstituted or substituted with one or more substituents selected from bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-15 4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl, piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-20 piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl, pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-25 pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl, aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-30 ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-

- 352 -

- butyl, sec-butyl, trifluoromethyl, pentafluoroethyl,
 nonafluorobutyl, dimethylaminopropyl, 1,1-
 di(trifluoromethyl)-1-hydroxymethyl, 1,1-
 di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-
 5 di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
 hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-
 aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
 isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
 phenoxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy,
 10 pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-
 ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-
 2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-
 ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and
 ethoxy; wherein R² is selected from H, chloro, fluoro,
 15 bromo, amino, hydroxy, methyl, ethyl, propyl, oxo,
 dimethylamino, aminosulfonyl, cyclopropyl, cyano,
 hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy,
 ethoxy, trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 20 substituted phenyl and unsubstituted or substituted
 heteroaryl selected from thienyl,
 furanyl, pyridyl, imidazolyl, and pyrazolyl; and
 wherein R⁴ is selected from a direct bond, ethyl, butyl, and

 ; and pharmaceutically acceptable derivatives
 25 thereof.

19. A compound of Claim 1 having Formula III

- 353 -




III

- wherein R is selected from unsubstituted or substituted 9-
or 10-membered fused nitrogen-containing heteroaryl,
5 where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
haloalkyl, C₁₋₆-alkoxy, optionally substituted
heterocyclalkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl,
C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-
10 alkoxy-C₁₋₆-alkoxy, and optionally substituted
heterocyclalk-C₂₋₄-alkynyl;
wherein R¹ is selected from unsubstituted or substituted
aryl,
cycloalkyl,
15 5-6 membered heteroaryl and
9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclalkyl,
wherein substituted R¹ is substituted with one or more
substituents selected from halo, C₁₋₆-alkyl, optionally
20 substituted C₃₋₆-cycloalkyl, optionally substituted
phenyl, optionally substituted phenyl-C_{1-C4}-alkylenyl,
C₁₋₂-haloalkoxy, optionally substituted phenyloxy,
optionally substituted 4-6 membered heterocyclalk-C_{1-C4}-
alkylenyl, optionally substituted 4-6 membered
25 heterocyclalk-C_{2-C4}-alkenylenyl, optionally substituted
4-6 membered heterocyclalkyl, optionally substituted 4-6
membered heterocyclalkoxy, optionally substituted 4-6
membered heterocyclalk-C₁₋₄-alkoxy, optionally
substituted 4-6 membered heterocyclalksulfonyl,

- 354 -


optionally substituted 4-6 membered heterocyclylamino,
 optionally substituted 4-6 membered
 heterocyclylcarbonyl, optionally substituted 4-6
 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 5 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
 C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
 alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
 alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋

10 ₄-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy;
 wherein R² is one or more substituents independently
 selected from

H,
 halo,
 15 hydroxy,
 amino,
 C₁₋₆-alkyl,
 C₁₋₆-haloalkyl,
 C₁₋₆-alkoxy,
 20 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 cyano,
 C₁₋₂-hydroxyalkyl,
 25 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 30 4-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 heterocyclyl;

- 355 -

wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and



; and

wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl; and

- 5 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₃-alkyl, 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;
 10 and pharmaceutically acceptable derivatives thereof.

20. Compound of Claim 19 wherein R is selected from indazolyl, where R is unsubstituted or substituted with one or more substituents selected from chloro, fluoro, amino,
 15 hydroxy, methyl, ethyl, propyl, trifluoromethyl, dimethylaminopropynyl, 1-methylpiperidinylmethoxy, dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein R¹ is selected from phenyl, tetrahydronaphthyl, indanyl, indenyl, naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl,
 20 thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, quinoxalyl, benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-
 25 1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolyl, indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl, benzothienyl, benzofuryl, dihydro-benzimidazolyl, benzimidazolyl, benzoxazolyl and benzthiazolyl, where R¹ is unsubstituted or substituted with one or more substituents
 30 selected from bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-

- 356 -

4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl, piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-

5 Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,

10 pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,

15 aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl), imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,

20 hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-

25 hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy, phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy,


30 pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and ethoxy; wherein R² is selected from H, chloro, fluoro,

- 357 -


bromo, amino, hydroxy, methyl, ethyl, propyl, oxo,
dimethylamino, aminosulfonyl, cyclopropyl, cyano,
hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy,
ethoxy, trifluoromethoxy, carboxymethyl,

5 morpholinylethylamino, propynyl, unsubstituted or
substituted phenyl and unsubstituted or substituted
heteroaryl selected from thienyl,

furanyl, pyridyl, imidazolyl, and pyrazolyl; and
wherein R^4 is selected from a direct bond, ethyl, butyl, and

10  ; and pharmaceutically acceptable derivatives
thereof.

21. A compound of Claim 1 having Formula IV



wherein A^3 is selected from CR^2 and N;

wherein A^4 is selected from CR^2 and N; provided one of A^3 and
 A^4 is not CR^2 ;

20 wherein R is selected from unsubstituted or
substituted 9- or 10-membered fused nitrogen-
containing heteroaryl,

where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C_{1-6} -alkyl, C_{1-6} -
25 C_{1-6} -haloalkyl, C_{1-6} -alkoxy, optionally substituted
heterocyclalkoxy, C_{1-6} -alkylamino- C_{2-4} -alkynyl,
 C_{1-6} -alkylamino- C_{1-6} -alkoxy, C_{1-6} -alkylamino- C_{1-6} -

- 358 -


alkoxy-C₁₋₆-alkoxy, and optionally substituted
heterocyclyl-C₂₋₄-alkynyl;

wherein R¹ is selected from unsubstituted or substituted
aryl,

5 cycloalkyl,
 5-6 membered heteroaryl and
 9-10 membered bicyclic and 13-14 membered
 tricyclic heterocyclyl,

wherein substituted R¹ is substituted with one or more

10 substituents selected from halo, C₁₋₆-alkyl, optionally
 substituted C₃₋₆-cycloalkyl, optionally substituted
 phenyl, optionally substituted phenyl-C_{1-C4}-alkylenyl,
 C₁₋₂-haloalkoxy, optionally substituted phenyloxy,
 optionally substituted 4-6 membered heterocyclyl-C_{1-C4}-
15 alkylenyl, optionally substituted 4-6 membered
 heterocyclyl-C_{2-C4}-alkenylenyl, optionally substituted
 4-6 membered heterocyclyl, optionally substituted 4-6
 membered heterocycliloxy, optionally substituted 4-6
 membered heterocyclyl-C₁₋₄-alkoxy, optionally
20 substituted 4-6 membered heterocyclylsulfonyl,
 optionally substituted 4-6 membered heterocyclylamino,
 optionally substituted 4-6 membered
 heterocyclylcarbonyl, optionally substituted 4-6
 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
25 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy,
 cyano, aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl,
 C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-
 alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-
 alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋


30 4-alkyl, C₁₋₄-hydroxyalkyl,  and C₁₋₄-alkoxy;

wherein R² is one or more substituents independently
selected from

H,

- 359 -

- halo,
hydroxy,
amino,
C₁₋₆-alkyl,
5 C₁₋₆-haloalkyl,
C₁₋₆-alkoxy,
C₁₋₂-alkylamino,
aminosulfonyl,
C₃₋₆-cycloalkyl,
10 cyano,
C₁₋₂-hydroxyalkyl,
nitro,
C₂₋₃-alkenyl,
C₂₋₃-alkynyl,
15 C₁₋₆-haloalkoxy,
C₁₋₆-carboxyalkyl,
5-6-membered heterocyclyl-C₁₋₆-alkylamino,
unsubstituted or substituted phenyl and
unsubstituted or substituted 4-6 membered
20 heterocyclyl;
wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and



- ; and
wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl; and
25 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₃-alkyl, 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;
30 and pharmaceutically acceptable derivatives thereof.

22. Compound of Claim 21 wherein R is selected from indazolyl, where R is unsubstituted or substituted with one

- 360 -

or more substituents selected from chloro, fluoro, amino, hydroxy, methyl, ethyl, propyl, trifluoromethyl, dimethylaminopropynyl, 1-methylpiperidinylmethoxy, dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein R¹ is

5 selected from phenyl, tetrahydronaphthyl, indanyl, indenyl, naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-

10 indolyl, naphthyridinyl, quinoxalinyl, benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolinyl, indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl, benzothienyl, benzofuryl, benzimidazolyl, dihydro-

15 benzimidazolyl, benzoxazolyl and benzthiazolyl, where R¹ is unsubstituted or substituted with one or more substituents selected from bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl,

20 phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl, piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-


25 Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,

30 pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,

- 361 -


- aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
- 5 imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-
- 10 di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
- 15 phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy, pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and
- 20 ethoxy; wherein R^2 is selected from H, chloro, fluoro, bromo, amino, hydroxy, methyl, ethyl, propyl, oxo, dimethylamino, aminosulfonyl, cyclopropyl, cyano, hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy, ethoxy, trifluoromethoxy, carboxymethyl,
- 25 morpholinylethylamino, propynyl, unsubstituted or substituted phenyl and unsubstituted or substituted heteroaryl selected from thienyl,

furanyl, pyridyl, imidazolyl, and pyrazolyl; and wherein R^4 is selected from a direct bond, ethyl, butyl, and

- 30  ; and pharmaceutically acceptable derivatives thereof.

23. A compound of Claim 1 having the formula V


- 362 -



- wherein A⁵ is selected from S, O and NR⁶;
- 5 wherein R is selected from unsubstituted or substituted 9-
 or 10-membered fused nitrogen-containing heteroaryl,
 wherein R is substituted with one or more substituents
 selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
 haloalkyl, C₁₋₆-alkoxy, optionally substituted
 10 heterocyclalkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl, C₁₋₆-
 alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-alkoxy-C₁₋₆-
 alkoxy, and optionally substituted heterocycl-C₂₋₄-
 alkynyl;
- wherein R¹ is selected from unsubstituted or substituted
 15 aryl,
 cycloalkyl,
 5-6 membered heteroaryl and
 9-10 membered bicyclic and 13-14 membered
 tricyclic heterocycl,
- 20 wherein substituted R¹ is substituted with one or more
 substituents selected from halo, C₁₋₆-alkyl, optionally
 substituted C₃₋₆-cycloalkyl, optionally substituted
 phenyl, optionally substituted phenyl-C_{1-C4}-alkylenyl, C₁-
 ₂-haloalkoxy, optionally substituted phenyloxy,
 25 optionally substituted 4-6 membered heterocycl-C_{1-C4}-
 alkylenyl, optionally substituted 4-6 membered
 heterocycl-C_{2-C4}-alkenylenyl, optionally substituted 4-6
 membered heterocycl, optionally substituted 4-6
 membered heterocycloxy, optionally substituted 4-6

- 363 -

membered heterocyclyl-C₁₋₄-alkoxy, optionally substituted
 4-6 membered heterocyclylsulfonyl, optionally substituted
 4-6 membered heterocyclylamino, optionally substituted 4-
 6 membered heterocyclylcarbonyl, optionally substituted
 5 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, cyano,
 aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-
 alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-
 C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-
 10 alkoxy carbonyl, C₁₋₄-alkoxy carbonylamino-C₁₋₄-alkyl, C₁₋₄-




wherein R² is one or more substituents independently
 selected from

H,
 15 halo,
 hydroxy,
 amino,
 C₁₋₆-alkyl,
 C₁₋₆-haloalkyl,
 20 C₁₋₆-alkoxy,
 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 cyano,
 25 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,
 30 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and

- 364 -

unsubstituted or substituted 4-6 membered
heterocyclyl;

wherein R^d is selected from a direct bond, C₁₋₄-alkyl, and



5 wherein R^e and R^f are independently selected from H and C₁₋₂-
haloalkyl;

wherein R^g is H or C₁₋₆-alkyl; and

wherein R^h is selected from H, C₁₋₃-alkyl, optionally

substituted phenyl, optionally substituted phenyl-C₁₋₃-
10 alkyl, 4-6 membered heterocyclyl, optionally substituted
4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-
alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;
and pharmaceutically acceptable derivatives thereof.

15 24. Compound of Claim 23 wherein R is selected from
indazolyl, where R is unsubstituted or substituted with one
or more substituents selected from chloro, fluoro, amino,
hydroxy, methyl, ethyl, propyl, trifluoromethyl,
dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
20 dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein Rⁱ is
selected from phenyl, tetrahydronaphthyl, indanyl, indenyl,
naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl,
1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl,
25 isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-
indolyl, naphthyridinyl, quinoxalinyl, benzo[d]isothiazolyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-
1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolinyl,
indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl,
30 benzothienyl, benzofuryl, benzimidazolyl, dihydro-
benzimidazolyl, benzoxazolyl and benzthiazolyl, where Rⁱ is
unsubstituted or substituted with one or more substituents
selected from bromo, chloro, fluoro, iodo, nitro, amino,


- 365 -

cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo,
aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl,
phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-
4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl,
5 piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-
methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-
(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-
Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-
piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-
10 4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-
ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl,
pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,
pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-
pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl,
15 pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl,
methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-
methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,
aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-
ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-
20 methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-
methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,
hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-
butyl, sec-butyl, trifluoromethyl, pentafluoroethyl,
25 nonafluorobutyl, dimethylaminopropyl, 1,1-
di(trifluoromethyl)-1-hydroxymethyl, 1,1-
di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-
di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-
30 aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy,
pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-
ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-

- 366 -

2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and ethoxy; wherein R² is selected from H, chloro, fluoro, bromo, amino, hydroxy, methyl, ethyl, propyl, oxo, dimethylamino, aminosulfonyl, cyclopropyl, cyano, hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy, ethoxy, trifluoromethoxy, carboxymethyl, morpholinylethylamino, propynyl, unsubstituted or substituted phenyl and unsubstituted or substituted heteroaryl selected from thienyl,


furanyl, pyridyl, imidazolyl, and pyrazolyl; wherein R⁴ is selected from a direct bond, ethyl, butyl, and



wherein R⁶ is H or methyl;

and pharmaceutically acceptable derivatives thereof.

25. A compound of Claim 1 having the formula



VI

20

wherein A⁵ is selected from S, O and NR⁶;

wherein R is selected from unsubstituted or substituted 9-

or 10-membered fused nitrogen-containing heteroaryl,

where R is substituted with one or more substituents

selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-haloalkyl, C₁₋₆-alkoxy, optionally substituted heterocyclalkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl, C₁₋₆-alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-alkoxy-C₁₋₆-

- 367 -


alkoxy, and optionally substituted heterocyclyl-C₂₋₄-alkynyl;

wherein R¹ is selected from unsubstituted or substituted aryl,

5 cycloalkyl,
 5-6 membered heteroaryl and
 9-10 membered bicyclic and 13-14 membered
 tricyclic heterocyclyl,

wherein substituted R¹ is substituted with one or more

10 substituents selected from halo, C₁₋₆-alkyl, optionally substituted C₃₋₆-cycloalkyl, optionally substituted phenyl, optionally substituted phenyl-C_{1-C4}-alkylenyl, C₁₋₂-haloalkoxy, optionally substituted phenyloxy, optionally substituted 4-6 membered heterocyclyl-C_{1-C4}-alkylenyl, optionally substituted 4-6 membered heterocyclyl-C_{2-C4}-alkenylenyl, optionally substituted 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocycliloxy, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkoxy, optionally substituted 20 4-6 membered heterocyclylsulfonyl, optionally substituted 4-6 membered heterocyclylamino, optionally substituted 4-6 membered heterocyclylcarbonyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, cyano, 25 aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋₄-alkyl, C₁₋₄-

hydroxyalkyl,  and C₁₋₄-alkoxy;


30 wherein R² is one or more substituents independently selected from

H,
halo,

- 368 -

- hydroxy,
 amino,
 C₁₋₆-alkyl,
 C₁₋₆-haloalkyl,
 5 C₁₋₆-alkoxy,
 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 cyano,
 10 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,
 15 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 heterocyclyl;

- 20 wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and



wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl;

wherein R⁶ is H or C₁₋₆-alkyl; and

- 25 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally substituted phenyl, optionally substituted phenyl-C₁₋₃-alkyl, 4-6 membered heterocyclyl, optionally substituted 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;
 30 and pharmaceutically acceptable derivatives thereof.

26. Compound of Claim 25 wherein R is selected from indazolyl, where R is unsubstituted or substituted with one

- 369 -

or more substituents selected from chloro, fluoro, amino, hydroxy, methyl, ethyl, propyl, trifluoromethyl, dimethylaminopropynyl, 1-methylpiperidinylmethoxy, dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein R¹ is

5 selected from phenyl, tetrahydronaphthyl, indanyl, indenyl, naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-

10 indolyl, naphthyridinyl, quinoxalinyl, benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolinyl, indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl, benzothienyl, benzofuryl, benzimidazolyl, dihydro-

15 benzimidazolyl, benzoxazolyl and benzthiazolyl, where R¹ is unsubstituted or substituted with one or more substituents selected from bromo, chloro, fluoro, iodo, nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl,

20 phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl, piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-

25 Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,


30 pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl, methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,

- 370 -

- aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
- 5 imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl, hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, trifluoromethyl, pentafluoroethyl, nonafluorobutyl, dimethylaminopropyl, 1,1-di(trifluoromethyl)-1-hydroxymethyl, 1,1-
- 10 di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
- 15 phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy, pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and
- 20 ethoxy; wherein R^2 is selected from H, chloro, fluoro, bromo, amino, hydroxy, methyl, ethyl, propyl, oxo, dimethylamino, aminosulfonyl, cyclopropyl, cyano, hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy, ethoxy, trifluoromethoxy, carboxymethyl,
- 25 morpholinylethylamino, propynyl, unsubstituted or substituted phenyl and unsubstituted or substituted heteroaryl selected from thienyl,

furanyl, pyridyl, imidazolyl, and pyrazolyl;

wherein R^4 is selected from a direct bond, ethyl, butyl, and




; and

wherein R^6 is H or methyl;

and pharmaceutically acceptable derivatives thereof.

- 371 -


27. A compound of Claim 1 having the formula



- 5 wherein A⁵ is selected from S, O and NR⁶;
 wherein R is selected from unsubstituted or substituted 9-
 or 10-membered fused nitrogen-containing heteroaryl,
 where R is substituted with one or more substituents
 selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
 haloalkyl, C₁₋₆-alkoxy, optionally substituted
 heterocyclalkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl, C₁₋₆-
 alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-alkoxy-C₁₋₆-
 alkoxy, and optionally substituted heterocycl-C₂₋₄-
 alkynyl;
 wherein R¹ is selected from unsubstituted or substituted
 aryl,
 cycloalkyl,
 5-6 membered heteroaryl and
 9-10 membered bicyclic and 13-14 membered
 tricyclic heterocycl,
 wherein substituted R¹ is substituted with one or more
 substituents selected from halo, C₁₋₆-alkyl, optionally
 substituted C₃₋₆-cycloalkyl, optionally substituted
 phenyl, optionally substituted phenyl-C₁₋₄-alkylenyl, C₁-
 2-haloalkoxy, optionally substituted phenyloxy,
 optionally substituted 4-6 membered heterocycl-C₁₋₄-
 alkylenyl, optionally substituted 4-6 membered
 heterocycl-C₂₋₄-alkenylenyl, optionally substituted 4-6
 membered heterocycl, optionally substituted 4-6
 membered heterocycl, optionally substituted 4-6
 membered heterocycl, optionally substituted 4-6

- 372 -

membered heterocyclyl-C₁₋₄-alkoxy, optionally substituted
 4-6 membered heterocyclylsulfonyl, optionally substituted
 4-6 membered heterocyclylamino, optionally substituted 4-
 6 membered heterocyclylcarbonyl, optionally substituted
 5 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, cyano,
 aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-
 alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-
 C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-
 10 alkoxy carbonyl, C₁₋₄-alkoxy carbonylamino-C₁₋₄-alkyl, C₁₋₄-




wherein R² is one or more substituents independently
 selected from

H,
 15 halo,
 hydroxy,
 amino,
 C₁₋₆-alkyl,
 C₁₋₆-haloalkyl,
 20 C₁₋₆-alkoxy,
 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 cyano,
 25 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,
 30 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and

- 373 -

unsubstituted or substituted 4-6 membered
heterocyclyl;

wherein R^d is selected from a direct bond, C₁₋₄-alkyl, and



; and

- 5 wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl;

wherein R^g is H or C₁₋₆-alkyl; and

wherein R^h is selected from H, C₁₋₃-alkyl, optionally

- substituted phenyl, optionally substituted phenyl-C₁₋₃-
10 alkyl, 4-6 membered heterocyclyl, optionally substituted
4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-
alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;
and pharmaceutically acceptable derivatives thereof.


- 15 28. Compound of Claim 27 wherein R is selected from
indazolyl, where R is unsubstituted or substituted with one
or more substituents selected from chloro, fluoro, amino,
hydroxy, methyl, ethyl, propyl, trifluoromethyl,
dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
20 dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein Rⁱ is
selected from phenyl, tetrahydronaphthyl, indanyl, indenyl,
naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl,
thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl,
1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl,
25 isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-
indolyl, naphthyridinyl, quinoxalinyl, benzo[d]isothiazolyl,
2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-
1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolinyl,
indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl,
30 benzothienyl, benzofuryl, benzimidazolyl, dihydro-
benzimidazolyl, benzoxazolyl and benzthiazolyl, where Rⁱ is
unsubstituted or substituted with one or more substituents
selected from bromo, chloro, fluoro, iodo, nitro, amino,

- 374 -

cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo,
aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl,
phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-
4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl,
5 piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-
methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-
(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-
Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-
piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-
10 4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-
ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl,
pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl,
pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-
pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl,
15 pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl,
methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-
methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,
aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-
ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-
20 methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-
methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,
hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-
butyl, sec-butyl, trifluoromethyl, pentafluoroethyl,
25 nonafluorobutyl, dimethylaminopropyl, 1,1-
di(trifluoromethyl)-1-hydroxymethyl, 1,1-
di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-
di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-
30 aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy,
pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-
ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-


- 375 -

- 2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and ethoxy; wherein R^2 is selected from H, chloro, fluoro, bromo, amino, hydroxy, methyl, ethyl, propyl, oxo, dimethylamino, aminosulfonyl, cyclopropyl, cyano, hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy, ethoxy, trifluoromethoxy, carboxymethyl, morpholinylethylamino, propynyl, unsubstituted or substituted phenyl and unsubstituted or substituted heteroaryl selected from thienyl, furanyl, pyridyl, imidazolyl, and pyrazolyl; wherein R^4 is selected from a direct bond, ethyl, butyl, and



- wherein R^6 is H or methyl; and pharmaceutically acceptable derivatives thereof.

29. Compound of Claim 1 of the formulas



- wherein A^5 is selected from S, O and NR^6 ; wherein R is selected from unsubstituted or substituted 9- or 10-membered fused nitrogen-containing heteroaryl, where R is substituted with one or more substituents selected from halo, amino, hydroxy, C_{1-6} -alkyl, C_{1-6} -haloalkyl, C_{1-6} -alkoxy, optionally substituted heterocyclylalkoxy, C_{1-6} -alkylamino- C_{2-4} -alkynyl, C_{1-6} -

- 376 -

alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-alkoxy-C₁₋₆-alkoxy, and optionally substituted heterocyclyl-C₂₋₄-alkynyl;

wherein R¹ is selected from unsubstituted or substituted

5 aryl,

cycloalkyl,

5-6 membered heteroaryl and

9-10 membered bicyclic and 13-14 membered

tricyclic heterocyclyl,

10 wherein substituted R¹ is substituted with one or more

substituents selected from halo, C₁₋₆-alkyl, optionally

substituted C₃₋₆-cycloalkyl, optionally substituted

phenyl, optionally substituted phenyl-C₁₋₄-alkylenyl, C₁₋

2-haloalkoxy, optionally substituted phenyloxy,

15 optionally substituted 4-6 membered heterocyclyl-C₁₋₄-

alkylenyl, optionally substituted 4-6 membered

heterocyclyl-C₂₋₄-alkenylenyl, optionally substituted 4-6

membered heterocyclyl, optionally substituted 4-6

membered heterocycliloxy, optionally substituted 4-6

20 membered heterocyclyl-C₁₋₄-alkoxy, optionally substituted

4-6 membered heterocyclylsulfonyl, optionally substituted

4-6 membered heterocyclylamino, optionally substituted 4-

6 membered heterocyclylcarbonyl, optionally substituted

4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-


25 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, cyano,

aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-

alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-

C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-

alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋₄-alkyl, C₁₋₄-

30 hydroxyalkyl,  and C₁₋₄-alkoxy;

wherein R² is one or more substituents independently


selected from

H,

- 377 -

halo,
hydroxy,
amino,
C₁₋₆-alkyl,
5 C₁₋₆-haloalkyl,
C₁₋₆-alkoxy,
C₁₋₂-alkylamino,
aminosulfonyl,
C₃₋₆-cycloalkyl,
10 cyano,
C₁₋₂-hydroxyalkyl,
nitro,
C₂₋₃-alkenyl,
C₂₋₃-alkynyl,
15 C₁₋₆-haloalkoxy,
C₁₋₆-carboxyalkyl,
5-6-membered heterocyclyl-C₁₋₆-alkylamino,
unsubstituted or substituted phenyl and
unsubstituted or substituted 4-6 membered
20 heterocyclyl;

wherein R^d is selected from a direct bond, C₁₋₄-alkyl, and



; and

wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl;


25 wherein R^g is H or C₁₋₆-alkyl; and
wherein R^h is selected from H, C₁₋₃-alkyl, optionally
substituted phenyl, optionally substituted phenyl-C₁₋₃-
alkyl, 4-6 membered heterocyclyl, optionally substituted
4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-
30 alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;
and pharmaceutically acceptable derivatives thereof.

- 378 -

30. Compound of Claim 29 wherein R is selected from indazolyl, where R is unsubstituted or substituted with one or more substituents selected from chloro, fluoro, amino, hydroxy, methyl, ethyl, propyl, trifluoromethyl,
- 5 dimethylaminopropynyl, 1-methylpiperidinylmethoxy, dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein R¹ is selected from phenyl, tetrahydronaphthyl, indanyl, indenyl, naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl,
- 10 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, quinoxalinyl, benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolinyl,
- 15 indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl, benzothienyl, benzofuryl, benzimidazolyl, dihydro-benzimidazolyl, benzoxazolyl and benzthiazolyl, where R¹ is unsubstituted or substituted with one or more substituents selected from bromo, chloro, fluoro, iodo, nitro, amino,
- 20 cyano, aminoethyl, Boc-aminoethyl, hydroxy, oxo, aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl, phenyl, phenylmethyl, morpholinylmethyl, 1-methylpiperazin-4-ylmethyl, 1-methylpiperazin-4-ylpropyl, morpholinylpropyl, piperidin-1-ylmethyl, 1-methylpiperidin-4-ylmethyl, 2-
- 25 methyl-2-(1-methylpiperidin-4-yl)ethyl, morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl, piperidin-4-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-1-ylethyl, 1-Boc-piperidin-4-ylethyl, piperidin-4-ylmethyl, 1-Boc-piperidin-4-ylmethyl, piperidin-4-ylpropyl, 1-Boc-piperidin-4-
- 30 ylpropyl, piperidin-1-ylpropyl, pyrrolidin-1-ylpropyl, pyrrolidin-2-ylpropyl, 1-Boc-pyrrolidin-2-ylpropyl, pyrrolidin-1-ylmethyl, pyrrolidin-2-ylmethyl, 1-Boc-pyrrolidin-2-ylmethyl, pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl,

- 379 -

- methylcarbonyl, Boc, piperidin-1-ylmethylcarbonyl, 4-
 methylpiperazin-1-ylcarbonylethyl, methoxycarbonyl,
 aminomethylcarbonyl, dimethylaminomethylcarbonyl, 3-
 ethoxycarbonyl-2-methyl-fur-5-yl, 4-methylpiperazin-1-yl, 4-
 5 methyl-1-piperidyl, 1-Boc-4-piperidyl, piperidin-4-yl, 1-
 methylpiperidin-4-yl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
 imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,
 hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-
 butyl, sec-butyl, trifluoromethyl, pentafluoroethyl,
 10 nonafluorobutyl, dimethylaminopropyl, 1,1-
 di(trifluoromethyl)-1-hydroxymethyl, 1,1-
 di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-
 di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
 hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-
 15 aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
 isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
 phenyloxy, azetidin-3-ylmethoxy, 1-Boc-azetidin-3-ylmethoxy,
 pyrrol-2-ylmethoxy, 1-Boc-pyrrol-2-ylmethoxy, pyrrol-1-
 ylmethoxy, 1-methyl-pyrrol-2-ylmethoxy, 1-isopropyl-pyrrol-
 20 2-ylmethoxy, 1-Boc-piperdin-4-ylmethoxy, piperdin-4-
 ylmethoxy, 1-methylpiperdin-4-yloxy, isopropoxy, methoxy and
 ethoxy; wherein R² is selected from H, chloro, fluoro,
 bromo, amino, hydroxy, methyl, ethyl, propyl, oxo,
 dimethylamino, aminosulfonyl, cyclopropyl, cyano,
 25 hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy,
 ethoxy, trifluoromethoxy, carboxymethyl,
 morpholinylethylamino, propynyl, unsubstituted or
 substituted phenyl and unsubstituted or substituted
 heteroaryl selected from thienyl,
 30 furanyl, pyridyl, imidazolyl, and pyrazolyl;
 wherein R⁴ is selected from a direct bond, ethyl, butyl, and




; and

wherein R⁶ is H or methyl;

- 380 -

and pharmaceutically acceptable derivatives thereof.

31. Compound of Claim 1 of the formula



IX

wherein A⁵ is selected from S, O and NR⁶;

wherein R is selected from

unsubstituted or substituted 9- or 10-membered fused
nitrogen-containing heteroaryl,

where R is substituted with one or more substituents
selected from halo, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-
haloalkyl, C₁₋₆-alkoxy, optionally substituted
heterocyclalkoxy, C₁₋₆-alkylamino-C₂₋₄-alkynyl, C₁₋₆-
alkylamino-C₁₋₆-alkoxy, C₁₋₆-alkylamino-C₁₋₆-alkoxy-C₁₋₆-
alkoxy, and optionally substituted heterocyclalkyl-C₂₋₄-
alkynyl;

wherein R¹ is selected from unsubstituted or substituted
aryl,

cycloalkyl,

5-6 membered heteroaryl and


9-10 membered bicyclic and 13-14 membered
tricyclic heterocyclalkyl,

wherein substituted R¹ is substituted with one or more

substituents selected from halo, C₁₋₆-alkyl, optionally
substituted C₃₋₆-cycloalkyl, optionally substituted
phenyl, optionally substituted phenyl-C₁₋₄-alkylenyl, C₁-
haloalkoxy, optionally substituted phenyloxy,
optionally substituted 4-6 membered heterocyclalkyl-C₁₋₄-

- 381 -

alkylenyl, optionally substituted 4-6 membered
 heterocyclyl-C₂-C₄-alkenylenyl, optionally substituted 4-6
 membered heterocyclyl, optionally substituted 4-6
 membered heterocyclyloxy, optionally substituted 4-6
 5 membered heterocyclyl-C₁₋₄-alkoxy, optionally substituted
 4-6 membered heterocyclylsulfonyl, optionally substituted
 4-6 membered heterocyclylamino, optionally substituted 4-
 6 membered heterocyclylcarbonyl, optionally substituted
 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 10 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, cyano,
 aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-
 alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-
 C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-
 alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋₄-alkyl, C₁₋₄-

15 hydroxyalkyl,  and C₁₋₄-alkoxy;


wherein R² is one or more substituents independently
 selected from

H,
 halo,
 20 hydroxy,
 amino,
 C₁₋₆-alkyl,
 C₁₋₆-haloalkyl,
 C₁₋₆-alkoxy,
 25 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 cyano,
 C₁₋₂-hydroxyalkyl,
 30 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 C₁₋₆-haloalkoxy,

- 382 -

C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 5 heterocyclyl;

wherein R^d is selected from a direct bond, C₁₋₄-alkyl, and



wherein R^e and R^f are independently selected from H and C₁₋₂-
 haloalkyl;

10 wherein R^g is H or C₁₋₆-alkyl; and

wherein R^h is selected from H, C₁₋₃-alkyl, optionally
 substituted phenyl, optionally substituted phenyl-C₁₋₃-
 alkyl, 4-6 membered heterocyclyl, optionally substituted
 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-

15 alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl;

and pharmaceutically acceptable derivatives thereof.

32. Compound of Claim 31 wherein R is selected from
 indazolyl, where R is unsubstituted or substituted with one
 20 or more substituents selected from chloro, fluoro, amino,
 hydroxy, methyl, ethyl, propyl, trifluoromethyl,
 dimethylaminopropynyl, 1-methylpiperidinylmethoxy,
 dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein Rⁱ is
 selected from phenyl, tetrahydronaphthyl, indanyl, indenyl,
 25 naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl,
 thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl,
 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl,
 isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-
 indolyl, naphthyridinyl, quinoxalinyl, benzo[d]isothiazolyl,
 30 benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-
 fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl,
 tetrahydroquinolinyl, indazolyl, 2,1,3-benzothiadiazolyl,
 benzodioxanyl, benzothienyl, benzofuryl, benzimidazolyl,

- 383 -

dihydro-benzimidazolyl, benzoxazolyl and benzthiazolyl,
where R¹ is unsubstituted or substituted with one or more
substituents selected from bromo, chloro, fluoro, iodo,
nitro, amino, cyano, aminoethyl, Boc-aminoethyl, hydroxy,
5 aminosulfonyl, 4-methylpiperazinylsulfonyl, cyclohexyl,
phenyl, phenylmethyl, morpholinylmethyl,
methylpiperazinylmethyl, methylpiperazinylpropyl,
morpholinylpropyl, methylpiperidinylmethyl,
morpholinylethyl, 1-(4-morpholinyl)-2,2-dimethylpropyl,
10 piperidinylethyl, piperidinylmethyl, piperidinylpropyl,
pyrrolidinylpropyl, pyrrolidinylpropenyl,
pyrrolidinylbutenyl, fluorosulfonyl, methylsulfonyl,
methylcarbonyl, piperidinylmethylcarbonyl,
methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-
15 ethoxycarbonyl-2-methyl-fur-5-yl, methylpiperazinyl,
methylpiperidyl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,
hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-
butyl, sec-butyl, trifluoromethyl, pentafluoroethyl,
20 nonafluorobutyl, dimethylaminopropyl, 1,1-
di(trifluoromethyl)-1-hydroxymethyl, 1,1-
di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-
di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-
25 aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
phenyloxy, 1-methylpiperidin-4-yloxy, isopropoxy, methoxy and
ethoxy; wherein R² is selected from H, chloro, fluoro,
bromo, amino, hydroxy, methyl, ethyl, propyl, oxo,
30 dimethylamino, aminosulfonyl, cyclopropyl, cyano,
hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy,
ethoxy, trifluoromethoxy, carboxymethyl,
morpholinylethylamino, propynyl, unsubstituted or


- 384 -

substituted phenyl and unsubstituted or substituted
heteroaryl selected from thienyl,

furanyl, pyridyl, imidazolyl, and pyrazolyl;

wherein R^4 is selected from a direct bond, ethyl, butyl, and

5




; and

wherein R^6 is H or methyl;

and pharmaceutically acceptable derivatives thereof.

33. Compound of Claim 1 of the formula

10



wherein A^5 is selected from S, O and NR^6 ;

wherein A^6 is selected from CR^2 and N;

15

wherein R is selected from

unsubstituted or substituted 9- or 10-membered fused
nitrogen-containing heteroaryl,

where R is substituted with one or more substituents

20

selected from halo, amino, hydroxy, C_{1-6} -alkyl, C_{1-6} -
haloalkyl, C_{1-6} -alkoxy, optionally substituted
heterocyclalkoxy, C_{1-6} -alkylamino- C_{2-4} -alkynyl, C_{1-6} -
alkylamino- C_{1-6} -alkoxy, C_{1-6} -alkylamino- C_{1-6} -alkoxy- C_{1-6} -
alkoxy, and optionally substituted heterocycl- C_{2-4} -
alkynyl;


25

wherein R^1 is selected from unsubstituted or substituted
aryl,

- 385 -

cycloalkyl,
 5-6 membered heteroaryl and
 9-10 membered bicyclic and 13-14 membered
 tricyclic heterocyclyl,

- 5 wherein substituted R¹ is substituted with one or more
 substituents selected from halo, C₁₋₆-alkyl, optionally
 substituted C₃₋₆-cycloalkyl, optionally substituted
 phenyl, optionally substituted phenyl-C₁₋₄-alkylenyl, C₁-
 2-haloalkoxy, optionally substituted phenyloxy,
 10 optionally substituted 4-6 membered heterocyclyl-C₁₋₄-
 alkylenyl, optionally substituted 4-6 membered
 heterocyclyl-C₂₋₄-alkenylenyl, optionally substituted 4-6
 membered heterocyclyl, optionally substituted 4-6
 membered heterocycliloxy, optionally substituted 4-6
 15 membered heterocyclyl-C₁₋₄-alkoxy, optionally substituted
 4-6 membered heterocyclylsulfonyl, optionally substituted
 4-6 membered heterocyclylamino, optionally substituted 4-
 6 membered heterocyclylcarbonyl, optionally substituted
 4-6 membered heterocyclyl-C₁₋₄-alkylcarbonyl, C₁₋₂-
 20 haloalkyl, C₁₋₄-aminoalkyl, nitro, amino, hydroxy, cyano,
 aminosulfonyl, C₁₋₂-alkylsulfonyl, halosulfonyl, C₁₋₄-
 alkylcarbonyl, C₁₋₃-alkylamino-C₁₋₃-alkyl, C₁₋₃-alkylamino-
 C₁₋₃-alkoxy, C₁₋₃-alkylamino-C₁₋₃-alkoxy-C₁₋₃-alkoxy, C₁₋₄-
 alkoxycarbonyl, C₁₋₄-alkoxycarbonylamino-C₁₋₄-alkyl, C₁₋₄-

- 25 hydroxyalkyl,  and C₁₋₄-alkoxy;


wherein R² is one or more substituents independently
 selected from

- H,
 halo,
 30 hydroxy,
 amino,
 C₁₋₆-alkyl,
 C₁₋₆-haloalkyl,

- 386 -


C₁₋₆-alkoxy,
 C₁₋₂-alkylamino,
 aminosulfonyl,
 C₃₋₆-cycloalkyl,
 5 cyano,
 C₁₋₂-hydroxyalkyl,
 nitro,
 C₂₋₃-alkenyl,
 C₂₋₃-alkynyl,
 10 C₁₋₆-haloalkoxy,
 C₁₋₆-carboxyalkyl,
 5-6-membered heterocyclyl-C₁₋₆-alkylamino,
 unsubstituted or substituted phenyl and
 unsubstituted or substituted 4-6 membered
 15 heterocyclyl;

wherein R⁴ is selected from a direct bond, C₁₋₄-alkyl, and








wherein R^e and R^f are independently selected from H and C₁₋₂-haloalkyl;

20 wherein R⁶ is H or C₁₋₆-alkyl;
 wherein R⁷ is selected from H, C₁₋₃-alkyl, optionally
 substituted phenyl, optionally substituted phenyl-C₁₋₃-
 alkyl, 4-6 membered heterocyclyl, optionally substituted
 4-6 membered heterocyclyl-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₂-
 25 alkyl and C₁₋₃-alkoxy-C₁₋₃-alkoxy-C₁₋₃-alkyl; and
 wherein



- 387 -



34. Compound of Claim 33 wherein R is selected from indazolyl, where R is unsubstituted or substituted with one or more substituents selected from chloro, fluoro, amino, hydroxy, methyl, ethyl, propyl, trifluoromethyl, dimethylaminopropynyl, 1-methylpiperidinylmethoxy, dimethylaminoethoxyethoxy, methoxy and ethoxy; wherein R^1 is selected from phenyl, tetrahydronaphthyl, indanyl, indenyl, naphthyl, cyclohexyl, isoxazolyl, pyrazolyl, thiazolyl, thiadiazolyl, thienyl, pyridyl, pyrimidinyl, pyridazinyl, 1,2-dihydroquinolyl, 1,2,3,4-tetrahydro-isoquinolyl, isoquinolyl, quinolyl, indolyl, isoindolyl, 2,3-dihydro-1H-indolyl, naphthyridinyl, quinoxalinyl, benzo[d]isothiazolyl, 2,3,4,4a,9,9a-hexahydro-1H-3-aza-fluorenyl, 5,6,7-trihydro-1,2,4-triazolo[3,4-a]isoquinolyl, tetrahydroquinolyl,

15
20
25

- 388 -


indazolyl, 2,1,3-benzothiadiazolyl, benzodioxanyl,
benzothienyl, benzofuryl, benzimidazolyl, dihydro-
benzimidazolyl, benzoxazolyl and benzthiazolyl, where R¹ is
unsubstituted or substituted with one or more substituents
5 selected from bromo, chloro, fluoro, iodo, nitro, amino,
cyano, aminoethyl, Boc-aminoethyl, hydroxy, aminosulfonyl,
4-methylpiperazinylsulfonyl, cyclohexyl, phenyl,
phenylmethyl, morpholinylmethyl, methylpiperazinylmethyl,
methylpiperazinylpropyl, morpholinylpropyl,
10 methylpiperidinylmethyl, morpholinylethyl, 1-(4-
morpholinyl)-2,2-dimethylpropyl, piperidinylethyl,
piperidinylmethyl, piperidinylpropyl, pyrrolidinylpropyl,
pyrrolidinylpropenyl, pyrrolidinylbutenyl, fluorosulfonyl,
methylsulfonyl, methylcarbonyl, piperidinylmethylcarbonyl,
15 methylpiperazinylcarbonylethyl, methoxycarbonyl, 3-
ethoxycarbonyl-2-methyl-fur-5-yl, methylpiperazinyl,
methylpiperidyl, 1-methyl-(1,2,3,6-tetrahydropyridyl),
imidazolyl, morpholinyl, 4-trifluoromethyl-1-piperidinyl,
hydroxybutyl, methyl, ethyl, propyl, isopropyl, butyl, tert-
20 butyl, sec-butyl, trifluoromethyl, pentafluoroethyl,
nonafluorobutyl, dimethylaminopropyl, 1,1-
di(trifluoromethyl)-1-hydroxymethyl, 1,1-
di(trifluoromethyl)-1-(piperidinylethoxy)methyl, 1,1-
di(trifluoromethyl)-1-(methoxyethoxyethoxy)methyl, 1-
25 hydroxyethyl, 2-hydroxyethyl, trifluoromethoxy, 1-
aminoethyl, 2-aminoethyl, 1-(N-isopropylamino)ethyl, 2-(N-
isopropylamino)ethyl, dimethylaminoethoxy, 4-chlorophenoxy,
phenyloxy, 1-methylpiperidin-4-yloxy, isopropoxy, methoxy and
ethoxy; wherein R² is selected from H, chloro, fluoro,
30 bromo, amino, hydroxy, methyl, ethyl, propyl, oxo,
dimethylamino, aminosulfonyl, cyclopropyl, cyano,
hydroxymethyl, nitro, propenyl, trifluoromethyl, methoxy,
ethoxy, trifluoromethoxy, carboxymethyl,
morpholinylethylamino, propynyl, unsubstituted or

- 389 -

substituted phenyl and unsubstituted or substituted
heteroaryl selected from thienyl,

furanyl, pyridyl, imidazolyl, and pyrazolyl;

wherein R⁴ is selected from a direct bond, ethyl, butyl, and



; and

wherein R⁶ is H or methyl;

and pharmaceutically acceptable derivatives thereof.

35. A pharmaceutical composition comprising a
10 pharmaceutically-acceptable carrier and a compound as in any
of Claims 1-34.

36. A method of treating cancer in a subject, said
method comprising administering an effective amount of a
15 compound as in any of Claims 1-34.

37. The method of Claim 36 comprising a combination
with a compound selected from antibiotic-type agents,
alkylating agents, antimetabolite agents, hormonal agents,
20 immunological agents, interferon-type agents and
miscellaneous agents.

38. A method of treating angiogenesis in a subject,
said method comprising administering an effective amount of
25 a compound as in any of Claims 1-34.

39. A compound as in any of Claims 1-34 for use in a
method of therapeutic treatment for the human or animal
body.

30 40. A method of treating KDR-related disorders in a
mammal, said method comprising administering an effective
amount of a compound of Claims 1-34.

- 390 -

41. A method of treating proliferation-related disorders in a mammal, said method comprising administering an effective amount of a compound of any of Claim 1-34.

5

42. Method of Claim 41 wherein the disorder is inflammation or an inflammation-related disorder.

43. Compound of Claim 1 and pharmaceutically acceptable salts thereof selected from

10 2-(1H-Indazol-6-ylamino)-N-[3-(3-morpholin-4-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide;

2-(1H-Indazol-6-ylamino)-N-[3-(3-piperidin-1-yl-propyl)-5-trifluoromethyl-phenyl]-nicotinamide;

15 2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-piperidin-4-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide;

2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide;

2-(1H-Indazol-6-ylamino)-N-[3-(piperidin-4-yloxy)-5-trifluoromethyl-phenyl]-nicotinamide;

20 2-(1H-Indazol-6-ylamino)-N-[3-(piperidin-4-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide;

N-(3,3-Dimethyl-1,1-dioxo-2,3-dihydro-1H-116-benzo[d]isothiazol-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;

25 2-(1H-Indazol-6-ylamino)-N-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-naphthalen-2-yl)-nicotinamide;

2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide;

30 2-(1H-Indazol-6-ylamino)-N-[3-(1-isopropyl-piperidin-4-ylmethoxy)-4-pentafluoroethyl-phenyl]-nicotinamide;

N-[3-(2-Hydroxy-3-pyrrolidin-1-yl-propoxy)-4-pentafluoroethyl-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide;

- 391 -

- 2-(1H-Indazol-6-ylamino)-N-[4-pentafluoroethyl-3-(2-piperidin-1-yl-ethoxy)-phenyl]-nicotinamide;
N-[3-(2-Hydroxy-3-pyrrolidin-1-yl-propoxy)-4-pentafluoroethyl-phenyl]-2-(1H-indazol-6-ylamino)-
5 nicotinamide;
2-(1H-Indazol-6-ylamino)-N-[4-pentafluoroethyl-3-(pyrrolidin-2-ylmethoxy)-phenyl]-nicotinamide;
2-(1H-Indazol-6-ylamino)-N-[4-pentafluoroethyl-3-(pyrrolidin-2-ylmethoxy)-phenyl]-nicotinamide;
10 2-(1H-Indazol-6-ylamino)-N-[3-(pyrrolidin-2-ylmethoxy)-4-trifluoromethyl-phenyl]-nicotinamide;
2-(1H-Indazol-6-ylamino)-N-[3-(2-pyrrolidin-1-yl-ethoxy)-4-trifluoromethyl-phenyl]-nicotinamide;
N-(1-Acetyl-3,3-dimethyl-2,3-dihydro-1H-indol-6-yl)-2-(1H-
15 indazol-6-ylamino)-nicotinamide;
2-(1H-Indazol-6-ylamino)-N-{4-[1-methyl-1-(1-methyl-piperidin-4-yl)-ethyl]-phenyl}-nicotinamide;
N-(4-Acetyl-2,2-dimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;
20 2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-piperidin-4-yl)-5-trifluoromethyl-phenyl]-nicotinamide;
N-(3-Bromo-5-trifluoromethyl-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide;
2-(1H-Indazol-6-ylamino)-N-(2,2,4-trimethyl-3,4-dihydro-2H-
25 benzo[1,4]oxazin-6-yl)-nicotinamide;
N-[4-tert-Butyl-3-(pyrrolidin-2-ylmethoxy)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide;
N-(7-Acetyl-5,5-dimethyl-5,6,7,8-tetrahydro-naphthalen-2-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;
30 1-Boc-2-(2-tert-Butyl-5-{[2-(1H-indazol-6-ylamino)-pyridine-3-carbonyl]-amino}-phenoxy-methyl)-pyrrolidine;
N-[4-tert-Butyl-3-(piperidin-4-ylmethoxy)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide;

- 392 -

- 2-(1H-Indazol-6-ylamino)-N-[3-(pyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide;
N-(4-tert-Butyl-3-piperazin-1-yl-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide;
5 N-(3,3-dimethyl-2,3-dihydro-1H-indol-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;
N-(3,3-Dimethyl-1-piperidin-4-yl-2,3-dihydro-1H-indol-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;
N-(2,2-Dimethyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;
10 N-[4-tert-Butyl-3-(4-propyl-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide;
N-[4-tert-Butyl-3-(4-isopropyl-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide;
15 2-(1H-Indazol-6-ylamino)-N-[3-(1-methylpyrrolidin-2-ylmethoxy)-5-trifluoromethyl-phenyl]-nicotinamide;
N-(4,4-Dimethyl-1-oxo-1,2,3,4-tetrahydro-isoquinolin-7-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;
N-(3,3-Dimethyl-2,3-dihydro-benzofuran-6-yl)-2-(1H-indazol-6-ylamino)-nicotinamide;
20 N-[1-(2-Dimethylamino-acetyl)-3,3-dimethyl-2,3-dihydro-1H-indol-6-yl]-2-(1H-indazol-6-ylamino)-nicotinamide;
2-(1H-Indazol-6-ylamino)-N-[3-(4-methyl-piperazin-1-ylmethyl)-4-pentafluoroethyl-phenyl]-nicotinamide;
25 2-(1H-Indazol-6-ylamino)-N-[3-(4-Boc-piperazin-1-ylmethyl)-4-pentafluoroethyl-phenyl]-nicotinamide;
2-(1H-Indazol-6-ylamino)-N-(3-morpholin-4-ylmethyl-4-pentafluoroethyl-phenyl)-nicotinamide;
2-(1H-Indazol-6-ylamino)-N-(4-pentafluoroethyl-3-piperazin-1-ylmethyl-phenyl)-nicotinamide;
30 N-[4-tert-Butyl-3-(4-Boc-piperazin-1-yl)-phenyl]-2-(1H-indazol-6-ylamino)-nicotinamide;
N-(4-tert-Butyl-3-nitro-phenyl)-2-(1H-indazol-6-ylamino)-nicotinamide;

- 393 -

- N-(3-Amino-4-tert-butyl-phenyl)-2-(1H-indazol-6-ylamino)-
nicotinamide;
- N-[4-tert-Butyl-3-(2-hydroxy-ethylamino)-phenyl]-2-(1H-
indazol-6-ylamino)-nicotinamide;
- 5 N-[4-tert-Butyl-3-(2-morpholin-4-yl-ethylamino)-phenyl]-2-
(1H-indazol-6-ylamino)-nicotinamide;
- N-[4-tert-Butyl-3-(1-Boc-piperidin-4-ylamino)-phenyl]-2-(1H-
indazol-6-ylamino)-nicotinamide;
- 2-(1H-Indazol-6-ylamino)-N-[2-(2-morpholin-4-yl-ethyl)-
1,2,3,4-tetrahydro-isoquinolin-7-yl]-nicotinamide;
- 10 N-[4-tert-Butyl-2-(4-methyl-piperazin-1-yl)-phenyl]-2-(1H-
indazol-6-ylamino)-nicotinamide;
- 2-(1H-Indazol-6-ylamino)-N-(2-oxo-4-trifluoromethyl-2H-
chromen-7-yl)-nicotinamide;
- 15 2-(1H-Indazol-6-ylamino)-N-[3-(1-methyl-1,2,3,6-tetrahydro-
pyridin-4-yl)-5-trifluoromethyl-phenyl]-nicotinamide;
- 2-(1H-Indazol-6-ylamino)-N-(1H-indol-7-yl)-nicotinamide;
- 2-(1H-Indazol-6-ylamino)-N-(4-pentafluoroethyl-phenyl)-
nicotinamide;
- 20 N-[4-tert-Butyl-3-(piperidin-4-ylamino)-phenyl]-2-(1H-
indazol-6-ylamino)-nicotinamide;
- 2-(1H-Indazol-6-ylamino)-N-(3-piperazin-1-ylmethyl-5-
trifluoromethyl-phenyl)-nicotinamide; and
- N-(4,4-Dimethyl-1,2,3,4-tetrahydro-isoquinolin-7-yl)-2-(1H-
25 indazol-6-ylamino)-nicotinamide.

44. Use of a compound of any of Claims 1-34 for
preparing a medicament for the treatment of cancer.

- 30 45. Use of a compound of any of Claims 1-34 for
preparing a medicament for the treatment of angiogenesis.

46. Use of a compound of any of Claims 1-34 for
preparing a medicament for the treatment of cell
35 proliferation.

- 394 -

47. A compound of any of Claims 1-34 and pharmaceutically acceptable derivatives thereof; for use as an active therapeutic substance.


5

48. Compound of Claim 47 for its anti-neoplasia use.

49. Compound of Claim 47 for its use in the treatment of angiogenesis.

10

50. A process for preparing compounds of Claim 1 comprising treatment of a compound of the formula



15

with R¹R⁴-NH₂, in the presence of base and an inert solvent; followed by coupling with a primary or secondary amine HNR⁵R;

20 where LG is halo, X^a is halo, ring A, A¹, A² and R² are as defined in Claim 1.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number
WO 02/068406 A3

(51) International Patent Classification⁷: **A61K 31/00**,
A61P 19/02, 35/00, C07D 231/56, 401/12, 401/14, 405/14,
407/12, 409/14, 413/14, 417/14, 471/04

(21) International Application Number: PCT/US02/03064

(22) International Filing Date: 11 January 2002 (11.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/261,882 12 January 2001 (12.01.2001) US
60/323,808 19 September 2001 (19.09.2001) US
10/046,622 10 January 2002 (10.01.2002) US

(71) Applicant: AMGEN INC. [US/US]; One Amgen Center
Drive, Thousand Oaks, CA 91320-1799 (US).

(72) Inventors: CHEN, Guoqing; 515 Oakbury Court, Thou-
sand Oaks, CA 91360 (US). ADAMS, Jeffrey; 2915
Venezia Lane, Thousand Oaks, CA 91362 (US). BEMIS,
Jean; 256 Appleton Street, Arlington, MA 02476 (US).
CROGHAN, Michael; 1108 Dover Lane, Ventura, CA
93001 (US). DIPIETRO, Lucian; 37 Centennial Avenue,
Gloucester, MA 01930 (US). DOMINGUEZ, Celia;
963 Cedarcliff Court, Thousand Oaks, CA 91362 (US).
ELBAUM, Daniel; 25 Sherrin Road, Newton, MA 02462
(US). GERMAIN, Julie; 28 Woods Avenue, Somerville,
MA 02144 (US). HUANG, Qi; 12308 Willow Forest
Drive, Moorpark, CA 93021 (US). KIM, Joseph, L.;
20 Green Way, Wayland, MA 01778 (US). OUYANG,
Xiaohu; 132-20 Maple Avenue, Flushing, NY 11355 (US).
PATEL, Vinod, F.; 3 Mossy Lane, Acton, MA 01720
(US). SMITH, Leon, M.; 33 Julie Court, Sommerset, NJ
08873 (US). TASKER, Andrew; 561 Granite Hills Street,

Simi Valley, CA 93065 (US). XI, Ning; 565 Timberwood
Avenue, Thousand Oaks, CA 91360 (US). XU, Shimin;
1890 West Hillcrest Drive, Number 406, Newbury Park,
CA 91320 (US). YUAN, Chester, Chenguang; 1416
Geyser Court, Newbury Park, CA 91320 (US). KIM,
Tae-Seong; 1580 Dover Avenue, Thousand Oaks, CA
91360 (US).

(74) Agents: ODRE, Steven, M. et al.; Amgen Inc., One Am-
gen Center Drive, M/S 27-4-A, Thousand Oaks, CA 91320-
1799 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:
24 April 2003

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED AMINE DERIVATIVES AND THEIR USE FOR THE TREATMENT OF ANGIOGENESIS



(I)

(57) Abstract: Selected amines of formula (I) are effective for prophylaxis and treatment of diseases, such as angiogenesis mediated diseases. The invention encompasses novel compounds, analogs, prodrugs and pharmaceutically acceptable salts thereof, pharmaceutical compositions and methods for prophylaxis and treatment of diseases and other maladies or conditions involving, cancer and the like. The subject invention also relates to processes for making such compounds as well as to intermediates useful in such processes.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 02/03064

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/00 A61P19/02 A61P35/00 C07D231/56 C07D401/12
 C07D401/14 C07D405/14 C07D407/12 C07D409/14 C07D413/14
 C07D417/14 C07D471/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 62778 A (BRISTOL MYERS SQUIBB CO) 26 October 2000 (2000-10-26) Claims 1,7-11,15-16,23,34; formulas (I) and (II); ex. 91,319-322,376-580 ---	17-30, 35-50
P,X	WO 01 78648 A (DONG WHA PHARM IND CO LTD ;KIM NAM DOO (KR); LEE JIN SOO (KR); LEE) 25 October 2001 (2001-10-25) Claims; formula 1; ex. 2-6,11,14-21 ---	17-30, 35-50
P,X	WO 01 55114 A (NOVARTIS ERFIND VERWALT GMBH ;MANLEY PAUL WILLIAM (CH); NOVARTIS A) 2 August 2001 (2001-08-02) Claims; formula (I); p. 8, 1st paragraph; p. 14, 1st paragraph --- -/--	17-30, 35-50

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

30 August 2002

Date of mailing of the international search report

31. 01. 03

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Rivat, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/03064

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 15118 A (THOMAS ANDREW PETER ;ZENECA LTD (GB); BROWN DEARG SUTHERLAND (GB);) 23 May 1996 (1996-05-23) Claims; formula (I); p. 42, l. 15-33; ex. 11, 15(27aa) ---	17-30, 35-50
Y	WO 00 27819 A (MANLEY PAUL WILLIAM ;NOVARTIS AKTIENGESELLSCHAFT (CH); BRUEGGEN JO) 18 May 2000 (2000-05-18) Claims; formula (I); p. 7, l. 26-31; p. 25, l. 15-p. 26, l. 12 ---	17-30, 35-50
A	WO 00 47212 A (PLE PATRICK ;HENNEQUIN LAURENT FRANCOIS AND (FR); ZENECA PHARMA SA) 17 August 2000 (2000-08-17) Claims 1-5,8,21-22; formula (I) ---	17-30, 35-50
A	WO 00 71532 A (NOE MARK CARL ;PFIZER PROD INC (US); GANT THOMAS GEORGE (US)) 30 November 2000 (2000-11-30) Claims 1, 8-20; formula 1; p. 8, l. 34-p. 9, l. 20 -----	17-30, 35-50

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/03064

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 36-4, 48-49 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1-13, 16, 31-34
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1 (part), 2-13, 16-50

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-13,16,31-34

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty . So many documents were retrieved that it is impossible to determine which parts of the claim(s) may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claim(s) is impossible. Consequently, the search has been restricted to the compounds given in formulas II-VI, VIIa-VIIb and VIIIa-VIIIb (claims 17-30), i.e. wherein A is a 5- or 6-membered heterocycle, R is a 9- or 10-membered fused nitrogen-containing heteroaryl, Z is O and R5 and R5a represent hydrogen, which definitions are also representative of a great amount of the examples disclosed in the present application.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 02/03064

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1(part),2-13,16-50

Heterocyclic compounds substituted in 1- and 2-positions by an heterocyclo-amino group and an N-cyclo-aminocarbonyl group and their use as VEGF ligands in the treatment of angiogenesis

2. Claims: 1(part),15,35-50

Aryl compounds substituted in 1- and 2-positions by an heterocyclo-amino group and an N-cyclo-aminocarbonyl group and their use as VEGF ligands in the treatment of angiogenesis

3. Claims: 1(part),14,35-50

Cycloalkene compounds substituted in 1- and 2-positions by an heterocyclo-amino group and an N-cyclo-aminocarbonyl group and their use as VEGF ligands in the treatment of angiogenesis

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/03064

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0062778 A	26-10-2000	AU 4233800 A BR 0009721 A CN 1348370 T CZ 20013677 A EP 1169038 A HU 0202708 A NO 20014970 A NZ 513639 A TR 200102969 T	02-11-2000 13-02-2002 08-05-2002 12-06-2002 09-01-2002 28-12-2002 10-12-2001 28-09-2001 21-08-2002
WO 0178648 A	25-10-2001	AU 4890301 A	30-10-2001
WO 0155114 A	02-08-2001	AU 2849901 A BR 0107805 A CZ 20022582 A EP 1259487 A NO 20023218 A	07-08-2001 22-10-2002 16-10-2002 27-11-2002 16-09-2002
WO 9615118 A	23-05-1996	AT 175962 T AU 703328 B AU 3813095 A CA 2200871 A DE 69507505 D DE 69507505 T DK 790986 T EP 0790986 A ES 2128092 T FI 971970 A JP 10508616 T NO 972152 A NZ 294917 A US 5821246 A ZA 9509572 A	15-02-1999 25-03-1999 06-06-1996 23-05-1996 04-03-1999 02-06-1999 13-09-1999 27-08-1997 01-05-1999 07-05-1997 25-08-1998 12-05-1997 25-02-1999 13-10-1998 13-05-1996
WO 0027819 A	18-05-2000	DE 19910396 A AU 1045400 A BG 105588 A BR 9915553 A CN 1325384 T CZ 20011631 A EE 200100258 A EP 1129074 A HU 0104425 A JP 2002529452 T NO 20012245 A PL 348349 A SK 6072001 A TR 200101307 T AU 1381100 A BR 9915210 A CA 2346898 A CN 1331680 T CZ 20011615 A WO 0027820 A EP 1129075 A HU 0104188 A JP 2002529453 T NO 20011894 A	07-09-2000 29-05-2000 30-04-2002 14-08-2001 05-12-2001 17-10-2001 16-12-2002 05-09-2001 28-03-2002 10-09-2002 10-07-2001 20-05-2002 07-01-2002 21-05-2002 29-05-2000 24-07-2001 18-05-2000 16-01-2002 15-08-2001 18-05-2000 05-09-2001 28-03-2002 10-09-2002 04-07-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PL 1/US 02/03064

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0027819 A		PL 347589 A	08-04-2002
		SK 6282001 A	11-09-2001
		TR 200101237 T	21-08-2001
		US 2002019414 A	14-02-2002
WO 0047212 A	17-08-2000	AU 2447500 A	29-08-2000
		BR 0008128 A	13-02-2002
		CN 1346271 T	24-04-2002
		CZ 20012889 A	14-11-2001
		EE 200100409 A	16-12-2002
		EP 1154774 A	21-11-2001
		HU 0104964 A	29-04-2002
		JP 2002536414 T	29-10-2002
		NO 20013882 A	09-10-2001
		SK 11402001 A	07-01-2002
		TR 200102314 T	21-01-2002
WO 0071532 A	30-11-2000	AU 4137400 A	12-12-2000
		BR 0010746 A	13-02-2002
		EP 1187826 A	20-03-2002
		US 6380214 B	30-04-2002